Case reports of suspected adverse drug reactions

Case reports were dismissed too quickly

EDITOR—The main difference between the replications by Loke et al of Venning's studies from the 1980s and the original is the evidence that is accepted as further follow-up.¹ Loke et al mention that 56 of 63 case reports on adverse reactions were cited at least once but accept only nine instances as valid follow-up: attempts at proof of mechanisms or controlled studies. However, unexpected adverse effects are by definition due to an unknown mechanism, and many established adverse effects still escape elucidation after years. Moreover, further accumulation of anecdotal reports can be sufficient evidence. A 10 year tally of 22 drugs that were withdrawn from the market showed that 18 were prompted by case reports and only four by a controlled study; for at least 13 the case reports proved sufficient reason for withdrawal.²

A highly unusual event that is repeatedly reported in a recently marketed drug, in the absence of publicity, points to an exceedingly high relative risk, a good argument for cause and effect.³ Moreover, these risks will be picked up spontaneously.^{4,5}

Apart from the nine instances of follow-up that were accepted by Loke et al, 15 datasheets were amended and seven BNP drug monographs changed. So, even with some overlap, close to half of the case reports were somehow sufficiently noteworthy to have had consequences. Given the small sample size, this is not far from Venning's original. It is a pity that Loke et al do not make all follow-up more explicit for all 56 instances that followed the 63 reports.

Their overall conclusion that lack of follow-up means that case reports are of limited value is a non-sequitur. It shows that a more consistent scheme of hypothesis-testing studies by industry, government, or third parties is needed instead of the current haphazard approach to drug safety. Such publications may alert and help clinicians in recognising potential adverse events in their patients. Often there is no other way of discovering something unusual. The record of systematic—that is, controlled—postmarketing surveillance in detecting

unexpected adverse effects remains surprisingly poor.

Jan P Vandenbroucke *professor of clinical epidemiology*
Leiden University Medical Center, 2300 RC
Leiden, Netherlands
J.P.Vandenbroucke@lumc.nl

Competing interests: None declared.

- 1 Loke YK, Price D, Derry S, Aronson JK. Case reports of suspected adverse drug reactions—systematic literature survey of follow-up. *BMJ* 2006;332:335-9. (11 February.)
- 2 Arnaiz JA, Carne X, Riba N, Codina C, Ribas J, Trilla A. The use of evidence in pharmacovigilance. Case reports as the reference source for drug withdrawals. *Eur J Clin Pharmacol* 2001;57:89-91.
- 3 Cornfield J, Haenszel W, Hammond EC, Lilienfeld AM, Shimkin MB, Wynder EL. Smoking and lung cancer: recent evidence and a discussion of some questions. *J Natl Cancer Inst* 1959;22:173-203.
- 4 Vandenbroucke JP. In defense of case reports and case series. *Ann Intern Med* 2001;134:330-4.
- 5 Ottavanger JP, Stricker BH. Loss of taste and terbinafine. *Lancet* 1992;340:728.

Case reports generate signals efficiently

EDITOR—Loke et al judge the value of case reports of suspected adverse drug reactions by the outcome measures of further formal validation and the incorporation into the product information.¹ They conclude that such reports are of limited value.

The main sources of information or tools that are available for detecting and evaluating suspected adverse

drug reactions are spontaneous reports, formal epidemiological studies and controlled clinical trials, and meta-analyses. Misunderstandings about the value of drug safety activities occur when inappropriate tasks or unrealistic expectations are assigned to these tools. However, used and interpreted appropriately, all of these methods have made major contributions to drug safety.²⁻⁴

According to Loke et al, 83% of reports had not been subjected to further evaluation. However, the first question is whether most clinically relevant suspected adverse drug reactions are initially communicated by case reports, and whether this process is efficient—sensitivity combined with low cost is the primary concern with spontaneous reports.

In contrast to many reports of limited quality and relevance that end up being buried in spontaneous reports databases, published case reports of suspected adverse drug reactions undergo peer review for

quality and relevance and are accessible to anyone caring about drug safety. They increase specific as well as general awareness of adverse drug effects and therefore their diagnosis in clinical practice; contribute to mechanistic hypotheses; and provide important information for the design and conduct of formal safety studies. They do all this at no cost and are therefore highly efficient.

Yes, in rare instances case reports do more harm than good, but further evaluations are able to identify such false signals or otherwise confirm and quantify safety problems.⁵ Published case reports remain a valid and efficient source for signal generation and are of great value for drug safety.

Stefan Russmann *assistant professor of epidemiology*
Boston Collaborative Drug Surveillance Program,
Boston University, Lexington, MA 02421, USA
russmann@bu.edu

Competing interests: None declared.

- 1 Loke YK, Price D, Derry S, Aronson JK. Case reports of suspected adverse drug reactions—systematic literature survey of follow-up. *BMJ* 2006;332:335-9. (11 February.)
- 2 Arnaiz JA, Carne X, Riba N, Codina C, Ribas J, Trilla A. The use of evidence in pharmacovigilance. Case reports as the reference source for drug withdrawals. *Eur J Clin Pharmacol* 2001;57:89-91.
- 3 Bresalier RS, Sandler RS, Quan H, Bolognese JA, Oxenius B, Horgan K, et al. Cardiovascular events associated with rofecoxib in a colorectal adenoma chemoprevention trial. *N Engl J Med* 2005;352:1092-102.
- 4 Jick H, Kaye JA, Jick SS. Antidepressants and the risk of suicidal behaviors. *JAMA* 2004;292:338-43.
- 5 Jick H, Kaye JA. Epidemiology and possible causes of autism. *Pharmacotherapy* 2003;23:1524-30.

Pain is an issue in renal impairment

EDITOR—Conway et al start their lesson of the week (that unrecognised accumulation of opiates in patients with renal failure may be life threatening) by saying that opiates and their metabolites accumulate in renal failure.¹ Not all opioids behave in the same way: some, such as fentanyl, have no active metabolites and others, such as hydromorphone, have minor active metabolites that do not accumulate substantially in renal failure. Morphine, pethidine, and dextro-propoxyphene all have active metabolites that accumulate in renal failure.² Codeine is methylmorphine and is metabolised in the same way as morphine, so it too can result in accumulation of active metabolites.²

In case 1 the authors imply that the deterioration in renal function was caused by the opioids, whereas the opioid toxicity is likely to have been a consequence of a deterioration in renal function, commonly seen perioperatively. We wonder whether the patient would have responded to the first

PHOTOS.COM

dose of naloxone had the intravenous cannula been patent.

Case 2 shows the care required when managing pain in patients with end stage renal failure. However, Davison found that pain is a problem in more than half of patients receiving haemodialysis and is often unrecognised and so not managed.³

Handbooks and guidelines already exist for managing pain in patients with renal impairment.^{4,5} Left unchallenged, this case report will add to the existing incorrect myths that surround opioid analgesics and prevent their proper use, denying patients with renal impairment adequate pain control. Unfortunately, unlike naloxone for opioid toxicity, no antidote exists for the uncontrolled pain resulting from opioidphobia.

Colette Reid research fellow in palliative medicine
colette.reid@bristol.ac.uk

Jane Gibbins specialist registrar, locum appointment training in palliative medicine

Geoffrey Hanks professor of palliative medicine
Department of Palliative Medicine, Bristol
Oncology Centre, Bristol BS2 8ED

Competing interests: None declared

- 1 Conway BR, Fogarty DG, Nelson WE, Doherty CC. Opiate toxicity in patients with renal failure. *BMJ* 2006;332:345-6. (11 February.)
- 2 Hanks GW, Hawkins C. Agreeing a gold standard in the management of cancer pain: the role of opioids. In: Hillier R, Finlay I, Welsh J, Miles A, eds. *The effective management of cancer pain*. London: Aesculapius Medical Press 2000:57-77.
- 3 Davison SN. Pain in hemodialysis patients: prevalence, cause, severity, and management. *Am J Kidney Dis* 2003;42:1239-47.
- 4 Chambers EJ, Brown EA, Germain M. *Supportive care for the renal patient*. 1st ed. Oxford: Oxford University Press, 2004.
- 5 Kurella M, Bennett WM, Chertow GM. Analgesia in patients with ESRD: a review of the available evidence. *Am J Kidney Dis* 2003;42:217-28.

Proteinuria in chronic renal disease was not mentioned

EDITOR—The 10 minute consultation on proteinuria by Haynes and Haynes did not mention recently developed multiprofessional UK guidelines on chronic kidney disease that include detailed, evidence based guidance on investigation, management, and referral of all presenting features of chronic kidney disease, including proteinuria.¹ These have been available for many months in draft form on the Renal Association's website; a concise version has recently been published.² They can be found through the National Electronic Library for Health's guideline finder (<http://libraries.nelh.nhs.uk/guidelinesFinder/>); and a full publication is in press at the Royal College of Physicians of London. Extensive decision support based on the guidelines is available at www.renal.org/eGFR/eguide.html, where the full guideline document can also be downloaded.

The most important difference—and potential cause of confusion—is in the recommendation for estimation of excretory kidney function by using the Cockcroft and Gault formula. The UK guidelines endorse the four variable MDRD formula, which does not require knowledge of body weight, and gives an estimated glomerular

filtration rate that is normalised to body surface area (as is routinely done for isotopic glomerular filtration rate measurements). The use of this formula is also endorsed by the Department of Health for England and will come into routine use in laboratories throughout the country very soon as part of the implementation of the national service framework for renal services.

Chronic kidney disease is common, and in most patients the management is simple. We encourage all those involved in the management of patients with chronic kidney disease to use the UK guidelines, which have been designed to be consistent wherever possible with other guidelines, including those in other national service frameworks and guidelines from other UK bodies.

Charlie Tomson consultant nephrologist
Southmead Hospital, Bristol BS10 5NB
ctomson@doctors.org.uk

The other five authors of this letter are Steve Blades, general practitioner, Newcastle upon Tyne; Richard Burden, consultant nephrologist, City Hospital, Nottingham; Julian Dennis, general practitioner, Newark; Edmund Lamb, clinical biochemist, and Paul Stevens, consultant nephrologist, Kent and Canterbury Hospital, Canterbury.

Competing interests: All authors were members of the working party that drew up the UK guidelines on the identification, management, and referral of adults with chronic kidney disease.

- 1 Haynes J, Haynes R. Proteinuria. *BMJ* 2006;332:284. (4 February.)
- 2 Burden R, Tomson C. Identification, management, and referral of adults with chronic kidney disease: concise guidelines. *Clin Med* 2005;5:635-42.

More on JAMA's policy on industry sponsored studies

EDITOR—Fontanarosa and DeAngelis refer to “numerous errors and misconceptions” in our editorial,¹ but they mention only two, one of which was attributable to editing by the *BMJ*.^{2,3} The other was our presumption that under JAMA's new policy, authors from industry must hire an academic statistician before submitting a paper. Fontanarosa and DeAngelis claim that the JAMA policy actually allows submissions from industry without an academic statistician. JAMA, however, will simply refuse to accept them until the authors purchase an academic statistical review.

If a JAMA paper must have an academic statistical author to be acceptable, we assumed that investigators from the private sector would not submit a paper to JAMA without such an author. Why would JAMA review such submissions if they are unacceptable a priori? Does this stance imply that the academic review is cursory, and that the academic statistician would not join the original investigators as an author? Or is JAMA suggesting that investigators from industry should submit papers with an incomplete list of authors, and then add another author during the review process?

Fontanarosa and DeAngelis describe our criticism of their policies as “blatant and

unbalanced” and they take umbrage “that a journal like the *BMJ* would criticise another journal's policy on such a pressing issue without properly vetting the material for accuracy.” We sent JAMA a draft of our essay; they did not respond. Furthermore, should not journal editors welcome criticism of their own work as warmly as they encourage critical appraisal of the articles they publish, even if the forum is outside of their control? Indeed, previous JAMA editors published criticism of editorial policies at other journals.⁴

Citing examples of industry malfeasance, of which we agree there are many, does little to justify an inequitable policy. In which examples was the malfeasance the responsibility of the industry statistician? Which would have been prevented by having the analysis reviewed by an academic statistician? As Senn noted,⁵ “independent” academic authors have their own biases, and they do not usually have to undergo the intense scrutiny of regulatory authorities.

Kenneth J Rothman vice president, epidemiology research

RTI Health Solutions, RTI International, 200 Park
Offices Drive, Research Triangle Park, NC 27709,
USA

Stephen Evans professor of pharmacoepidemiology
Medical Statistics Unit, London School of Hygiene
and Tropical Medicine, London WC1 7HT
stephen.evans@lshtm.ac.uk

Competing interests: KJR is an employee of RTI Health Solutions, an independent non-profit research organisation that does work for government agencies and pharmaceutical companies. SE has no competing interests to declare.

- 1 Fontanarosa PB, DeAngelis CD. JAMA's policy on industry sponsored studies. *BMJ* 2006;332:177. (21 January.)
- 2 Rothman KJ, Evans S. Response. Corrections and a rebuttal to Vandenbroucke. *bmj.com* 2005. <http://bmj.bmjjournals.com/cgi/letters/331/7529/1350#123723>
- 3 Corrections and clarifications. Extra scrutiny for industry funded trials. *BMJ* 2006;332:151. (21 January.)
- 4 Rothman KJ. Conflict of interest—the new McCarthyism in science. *JAMA* 1993;269:2782-4.
- 5 Senn SJ. JAMA's policy does not go far enough. *BMJ* 2006;332:305. (4 February.)

*The editorial's title was changed inhouse after the return of the proofs. Unfortunately, the correction was not highlighted in the letter from JAMA's editors, although both were published in the same issue (21 January).

Compensation for non-negligent harm in trials remains shaky

EDITOR—Last year in the *BMJ* I showed that a patient's agreement to take part in a clinical trial is a legal contract which consumer law requires to be expressed in plain language.^{1,2} In a letter to me on behalf of the chief medical officer (for which I am grateful), the Department of Health has now agreed that where there are contracts between patients and trials sponsors, consumer law is applicable.

I had cited words in my article as not being plain language that the Department of Health's Central Office for Research Ethics Committees (COREC)³ was advocating for use in information to potential patient

volunteers for clinical trials. The words were: "Broadly speaking... the sponsor, *without legal commitment, should* compensate you [for non-negligent harm], without you having to prove that it is at fault" (my italics).⁴

The Department of Health told me that it was always open to suggestions for improvement and would ask COREC to review whether the words could leave a lay person in any doubt that there is no automatic right to compensation.

On 22 November 2005 the Department of Health and COREC published replacement words: "[the sponsor] *will* pay compensation for [non-negligent harm]. Any payment would be without legal commitment (please ask if you wish for more information on this)" (my italic).⁵ Clearly these new words also are not plain language: the certainty of compensation given in the first sentence is negated by the legal jiggery-pokery of the second.

Unforeseen risks are inherent in medical research. In clinical trials patients will inevitably sometimes suffer serious non-negligent harm (for which the courts will not award compensation when there is no prior legally binding commitment to compensate).

As a matter of law, the Department of Health and COREC may not recruit patients for trials by concealing the fact that present compensation arrangements for non-negligent harm are flimsy and unreliable. The Office of Fair Trading has legal powers to ask a court for an injunction to prevent such misrepresentation.² The department should now obey the law. If it simply will not, then the Office of Fair Trading should now enforce the law.

Desmond R Laurence *professor emeritus, University College London*
London NW3 1ST
d.laurence@virgin.net

Competing interests: None declared.

- 1 Laurence DR. Participants in research. *BMJ* 2005;331:110.
- 2 Office of Fair Trading. *Unfair terms in contracts—what is an unfair term?* www.offt.gov.uk/Consumer/Unfair+terms+in+contracts/what+are+unfair+terms.htm (accessed 5 Dec 2005).
- 3 Central Office for Research Ethics Committees (COREC). *Annual report 2004*. www.corec.org.uk/recs/about/docs/COREC_Annual_Report_2004.doc (accessed 6 Dec 2005).
- 4 Central Office for Research Ethics Committees. *Guidelines for researchers*. Para 15. What if something goes wrong? February 2001. www.corec.org.uk/applicants/help/docs/Guidance_on_Patient_Information_Sheets_and_Consent_Forms.doc (accessed 5 Jun 2005).
- 5 Central Office for Research Ethics Committees. *Guidelines for researchers*. Consultation version 22 November 2005; para 19. What if there is a problem? Harm. www.corec.org.uk/applicants/help/docs/Guidance_on_Patient_Information_Sheets_and_Consent_Forms.doc (accessed 6 Dec 2005).

Referrals between consultants

Public should rise to the defence

EDITOR—Goodman's complaint about the deferral of consultant to consultant referrals is timely.¹ Only a fortnight ago a consultant at my local hospital insisted on speaking with me about a case that was considered to need urgent attention by another department in the same hospital. My primary care trust has apparently recently decreed that there should be no consultant to consultant referrals forthwith, except in the case of suspected cancer.

This is inefficient and does not seem ethical. Certainly, my patient needed an outpatient appointment then and should not wait until April 2006, the earliest "urgent" appointment that I could manage, even with telephoning the secretary and faxing my referral. Thankfully, the first consultant has managed to get the needed urgent appointment, but only after speaking with me and haggling with the second team.

There has been no forewarning of this new state of affairs: the trust did not see fit to write to local general practitioners. We have not had a chance to discuss it or lobby for something different, if not preserve the status quo. If ever there was an example of how managers with no clinical knowledge can wield so much power over the fate and wellbeing of the population, this is it. The public should know and demand that this silliness stops.

David M Lewis *general practitioner*
Tudor Surgery, Watford WD24 7PH
david.lewis@gp-E82034.nhs.uk

Competing interests: DML is a general practitioner who is directly affected by this situation.

- 1 Goodman M. Defending referrals between consultants. *BMJ* 2006;332:371. (11 February.)

System has potential for error

EDITOR—Goodman draws attention to a problem that frustrates general practitioners and hospital doctors alike: it is ludicrous that one professional cannot readily refer to another in the hospital system.¹

Much anecdotal evidence shows that this is a major problem. Deliberately introducing delay into such a referral system increases the risk of problems arising in the interim. Perhaps the time has come for a formal study of the effects of obstructing efficient clinical referral. If hospital risk managers are not considering this, then why not?

The rising practice of asking general practitioners to make a referral to a second consultant at the request of a first consultant introduces an extra step into the referral process, which increases the potential for error and also delay. Many general practitioners are extremely efficient, but is it fair to them, or patients, to erect an unnecessary hurdle to referral that a consultant considers necessary? I believe not. Giving general practitioners unnecessary tasks diverts their clinical time from appropriate clinical activity.

Goodman mentions exercise electrocardiography. In the 1990s—when fundholding existed and importance was attached to general practitioners' referrals—I had control over exercise electrocardiography lists as a senior house officer in two teaching hospitals. Accepting a referral from a hospital colleague was no problem, and providing a prompt service, with immediate reporting, was a source of professional satisfaction.

Certainly hospital doctors should assert their professional status; it is, after all, in the best interests of patients.

Peter Gooderham *tutor*
Cardiff Law School, Cardiff CF10 3XJ
GooderhamEP@Cardiff.ac.uk

Competing interests: PG has an academic interest in preventing medical error and is a retired medical practitioner.

- 1 Goodman M. Defending referrals between consultants. *BMJ* 2006;332:371. (11 February.)

Consultant referrals are funded

EDITOR—Goodman correctly says that consultant to consultant referrals are not part of the monitoring of waiting time standards for trusts and therefore may be given less priority by trust managers.¹ However, he is mistaken in his assertion that consultant to consultant referrals are not funded and that this may also influence how managers behave.

Under payment by results, all outpatient attendances are charged at tariff. First outpatient appointments have a higher tariff than follow-up attendances, but the source of the referral is not relevant. Primary care trusts therefore fund trusts for consultant to consultant referrals in exactly the same way as they do for general practitioners' referrals. It is this issue that is driving some primary care trusts to try to get a better grip on referrals between consultants.

There will always be perverse incentives in a bureaucracy as complex as the NHS, but it is important to understand the rules of the system before ascribing possible motives to others.

Tim E P Davies *director of public health*
South Worcestershire Primary Care Trust,
Worcester WR4 9RW
tep.davies@btinternet.com

Competing interests: None declared.

- 1 Goodman M. Defending referrals between consultants. *BMJ* 2006;332:371. (11 February.)

Prognostic assessment will be important in flu pandemic

EDITOR—As highlighted by Bonten and Prins, secondary pneumonia will cause considerable problems in a flu pandemic.¹ Symptoms and signs, however, have low predictive value in diagnosing pneumonia.² As patients with secondary pneumonia are likely to be sicker, prognostic assessment could support clinicians' judgment.

One potential predictive tool is CURB65, which has been validated in community-acquired pneumonia and is recommended by the British Thoracic Society.³ This uses four bedside criteria (confusion, respiratory rate, blood pressure, and age) and one laboratory criterion (urea) to determine a patient's risk of death. In primary care the urea component can be dropped with similar performance.² During a pandemic, CURB65 could be used to target interventions such as neuraminidase inhibitors, antibiotics, and hospital assessment. Its use in flu infection warrants further investigation.

The British Thoracic Society recommends amoxicillin for patients with community acquired pneumonia at low risk of death (CURB65 = 0-1) and amoxicillin plus a macrolide in those at moderate risk (CURB65 = 2).³ Given that these cover the

most likely cause of secondary infection (*Streptococcus pneumoniae*) and that co-infection with atypical pathogens is recognised,³ their use should be retained in a pandemic.

A regimen for severe pneumonia (CURB65=3-5) is more difficult. The society currently recommends intravenous coamoxiclav or a second or third generation cephalosporin plus a macrolide. Intravenous levofloxacin plus benzylpenicillin is an alternative.² Some hospitals use the latter to limit *Clostridium difficile* associated diarrhoea. In North America respiratory fluoroquinolones are commonly used. Although all have antipneumococcal activity, fluoroquinolones are not reliable in staphylococcal infections. In east Yorkshire 30% of methicillin sensitive *Staphylococcus aureus* and 98% of methicillin resistant (MRSA) isolates are resistant.

In areas with community acquired MRSA, the optimal regimen is unclear. Glycopeptides poorly penetrate lung tissue.⁴ Oxazolidinones may be more efficacious and are easier to use.⁵ Hospital antibiotic committees will therefore need to think carefully before making recommendations for pneumonia in a flu pandemic.

Gavin D Barlow consultant in infectious diseases and medicine

Department of Infection and Tropical Medicine,
Hull and East Yorkshire Hospitals NHS Trust,
Castle Hill Hospital, Cottingham, North
Humberside HU16 5JQ
Gavin.Barlow@hey.nhs.uk

Competing interests: None declared.

- 1 Bonten MJM, Prins JM. Antibiotics in pandemic flu will be essential for treating, but not preventing, bacterial pneumonia. *BMJ* 2006;332:248-9. (4 February)
- 2 British Thoracic Society. BTS Guidelines for the management of community acquired pneumonia in adults. *Thorax* 2001;56(suppl 4):1-64.
- 3 Lim WS, Macfarlane JT, Boswell TCJ, Harrison TG, Rose D, Leinonen M, et al. Study of community acquired pneumonia aetiology (SCAPA) in adults admitted to hospital: implications for management guidelines. *Thorax* 2001;56:296-301.
- 4 Cruciani M, Gatti G, Lazzarini L, Furlan G, Broccali G, Malena M, et al. Penetration of vancomycin into human lung tissue. *J Antimicrob Chemother* 1996;38:865-9.
- 5 Kollef MH, Rello J, Cammarata SK, Croos-Dabrera RV, Wunderink RG. Clinical cure and survival in Gram-positive ventilator-associated pneumonia: retrospective analysis of two double-blind studies comparing linezolid with vancomycin. *Intensive Care Med* 2004;30:388-94.

Best research

New strategy is warmly welcomed

EDITOR—Surely the goal of research in the NHS should be to achieve synergy, not opposition between research of the highest international quality and maintaining and improving public health and the care of present and future patients. This requires tackling the needs of several constituencies in a way that can benefit them all. It also requires allocation of resource in a way that will most effectively achieve this goal, given that resource is always finite.

Research policy also has to achieve collaboration with those players outside the NHS notably, but not exclusively, the pharmaceutical and medical devices industries, without whom successful use of research in many domains of practice cannot be

achieved. Such relations should not be focused on “freebies” or drug lunches but on the recognition that only ethical and effective relations with commerce can provide in quantity for the NHS, and other healthcare systems worldwide, the further new vaccines, pharmaceuticals, and instrumentation that have been so effective at improving health care over the past generation.

In the Greater Manchester research alliance, we are endeavouring to achieve just this synergy of patients who volunteer to engage in clinical studies and trials with clinicians and academics supported by NHS management and the commercial sector.

We are optimistic that the new Department of Health's strategy, “Best Research for Best Health,” will support us and others in our aims. It has identified the right challenges and boldly established a coherent group of appropriate solutions, together with funding streams ringfenced for research. Its central themes of transparent allocation of resource in response to quality and of support for research staff to develop and to achieve their goals unencumbered by the layers of bureaucracy reported by Galbraith et al¹ will, we believe, provide the most important step change in research for and with patients since the inception of the NHS research and development programme some 15 years ago.

We do not recognise the new strategy in the editorial by Pollock et al.² By contrast, we welcome it warmly.

Robert Boyd Greater Manchester director of NHS research and development
University of Manchester, Manchester M13 9PL
rboyd@doctors.org.uk

Mike Burrows chief executive
Salford Primary Care Trust

Competing interests: The North West is a region that might benefit from the new strategy.

- 1 Galbraith N, Hawley C, De-Souza V. Research governance: Research governance approval is putting people off research. *BMJ* 2006;332:238. (28 January)
- 2 Pollock A, McNally N, Kerrison S. Best research. *BMJ* 2006;332:247-8. (4 February)

Research originating in primary care needs support

EDITOR—Pollock et al clearly summarise the problems inherent in the UK Department of Health's new research strategy.^{1,2} The proposed reorganisation of primary care research networks is also a concern: pharmaceutical industry representatives may have membership of key strategic groups in reorganised primary care networks and the predominant research paradigm will become the large randomised controlled trial.

One of the clear strengths of the current networks is the broad range of studies and methods that can be identified in their research portfolios. This range of topics represents the diversity of work in primary care as well as the welcome opportunity over recent years for many to investigate an area of their practice that has fired their curiosity and enthusiasm. There are potential advantages to the NHS as a whole in practitioners becoming involved in their own research, albeit on a small scale, since this is likely to lead to individuals possessing the critical skills

required to practise evidence based medicine. The presence of this cohort of research experienced practitioners is also a resource for other colleagues in primary care.

The proposed total budget for the portion of the Department of Health's funding available to open competition will be only £25m. This will mean that obtaining funding for small scale projects will be an uphill struggle.

The practice of good primary health care depends on a broad range of knowledge—for example, from the beneficial effects of medicines to why patients do not take them. A straw poll of “papers that changed my practice” among primary care practitioners is not likely to produce a list of reports of randomised controlled trials. Unfortunately the centrally prescribed clinical areas to be tackled in the new research networks, together with commercial pressures, are likely to deliver an unhealthy balance of trials of therapeutic agents. The unique contribution of primary care research is in danger of being diminished.

Jim Sikorski chair
j.sikorski@iop.kcl.ac.uk

Susan Robinson research fellow
Martin Edwards GP researcher

Lewisham Research Unit, London SE26 4TH

Competing interests: The correspondents are members, and have received funding from, Lewisham Research Unit, a Department of Health funded primary care research group.

- 1 Pollock A, McNally N, Kerrison S. Best research. *BMJ* 2006;332:247-8. (4 February)
- 2 Department of Health. *Best research for best health: a new National Health research strategy*. London: DH, 2006. www.dh.gov.uk/assetRoot/04/12/71/52/04127152.pdf (accessed 16 Feb 2006).

Tobacco: are we all to blame?

EDITOR—Gilmore et al have written a clear commentary on the activities of British American Tobacco (BAT), and, indirectly, the investment bank Schroders.¹ Almost all of us who have savings for retirement or other long term needs will have invested in these companies through collective investment vehicles such as pension funds or individual savings accounts (ISAs).

Equally, if we are paid by the NHS our pay will come in part from monies collected as taxation from the company profits of companies such as Schroders and BAT. So none of us is exempt from our share of the collective guilt to be associated with the investment policies described in this paper.

Jonathan H Burton editor, *Work Based Learning in Primary Care*
London Deanery, London WC1B 5DN
jburton@londondeanery.ac.uk

Competing interests: JHB holds savings in investment trusts that have in the past invested or still invest in British American Tobacco, among other international companies, and has worked as a consultant in a primary care development project in Uzbekistan.

- 1 Gilmore AB, Collin J, McKee M. British American Tobacco's erosion of health legislation in Uzbekistan. *BMJ* 2006;332:355-8. (11 February)