Published in final edited form as:

J Obstet Gynecol Neonatal Nurs. 2005 ; 34(5): 595–605.

Perinatal Loss in Low-income African American Parents: The Lived Experience

Karen Kavanaugh, PhD, RN and

University of Illinois at Chicago College of Nursing, Associate Professor and Co-Project Director, Advanced Practice Palliative Nurse Training Program, University of Illinois at Chicago College of Nursing, Department of Maternal-Child Nursing, 845 South Damen, Room 848, Chicago, Illinois 60612-7350

Patricia Hershberger, MSN, RNC, APRN, BC, FNP

Assistant Professor, Valparaiso University, Valparaiso, Indiana, Doctoral Candidate, University of Illinois at Chicago, Chicago, Illinois

Abstract

Objective: To examine the experience of low-income, African-American parents surrounding perinatal loss and to describe how other life stressors influenced the parents' responses and caring needs.

Design: Descriptive, using a phenomenologic approach.

Setting: All data were collected in person. Interviews were held in parents' homes or, at the request of three parents, in an office in the university between 5 and 21 weeks after the loss.

Participants: A total of 23 parents (17 mothers and 6 of their partners) were interviewed following a perinatal loss due to pregnancy (16 weeks gestation or later) or a neonatal death (first 28 days of life). Follow-up interviews were held with 21 of these parents.

Results: Revealed four themes: (a) recognizing problems and responding to the loss, (b) dealing with stressful life events, (c) creating and cherishing memories of their infant, and (d) living with the loss.

Conclusions: The results of this study reveal experiences not previously reported, and provide a beginning understanding of the loss experience in this group of parents. Health care professionals should be aware of the presence of additional stressful events that parents may be experiencing and intervene appropriately to provide culturally competent care in a sensitive manner.

Keywords

African Americans; parental grief; perinatal loss; phenomenological research

Numerous studies have been conducted on perinatal loss but mainly with an exclusive (Crowther, 1995; deMontigny, Beaudet, & Dumas, 1999; Madden, 1994; Theut et al., 1989) or primarily middle class Caucasian population of parents (DiMarco, Menke, & McNamara, 2001; Swanson, 1999; Thomas & Striegel, 1994–5). Furthermore, only a few investigations included parents with a wide range of socioeconomic levels, and the majority of these were Caucasian (Lang & Gottlieb, 1993; Mekosh-Rosenbaum & Lasker, 1995; Stinson, Lasker, Lohmann, & Toedter, 1992). However, these investigators did not report differences due to economic variables. The experiences of African American parents and loss were recently examined (Van, 2001; Van & Meleis, 2003) but these studies included only mothers who were primarily middle to upper income. Therefore, little attention has been given to the experience of perinatal loss in low-income, African Americans.

Studying the responses of African American parents is critical because of the high incidence of perinatal loss among blacks, and the other stressful life situations that African Americans often experience concurrently. Perinatal data for blacks demonstrate that the incidence of perinatal loss in blacks is more than twice as high as whites (Arias, MacDorman, Strobino, & Guyer, 2003). Furthermore, a recent study of African American and white women in the Detroit area demonstrated that African American women who lived in the city had significantly more acute life events, such as the death of someone close, than white women (Schulz et al., 2000). In that study, socioeconomic status made a significant contribution to variations in health status. Also, in clinical observations of mourning in low-income families, one-third of whom were black, Fulmer (1987) reported frequent and intense negative environmental pressures which gave the families more occasion to mourn but depleted their capacity to do so. Therefore, because of the lack of research in this area, the purpose of this study was to examine the experience of low-income, African American parents surrounding perinatal loss due to stillbirth (fetal death of at least 16 weeks gestation) or neonatal death (live-born infant death of less than 28 days), and to describe how other life stressors influence the parents' response and caring needs.

Low-income, Childbearing African American Women and Stress

In a study of urban, African-American, single mothers, Brodsky (1999) described both protective and risk factors in the women. 'Making it' was the overall process used by mothers to respond to the stress, which was brought on by their neighborhood, relationships with family and friends, and lack of money. An essential characteristic of 'making it' was cultivating a sense of appreciation for even small successes.

Natality and mortality data for blacks demonstrate that black women have worse pregnancy outcome in comparison to white women (Arias et al., 2003). Poor pregnancy outcome, as defined by a high percentage of low birth weight and preterm infants and high infant mortality, has been attributed to a number of factors, including stress and poor prenatal care. Even though the percentage of black women who began prenatal care in their first trimester of pregnancy has increased in recent years, racial disparities still exist (88.7% for white women; 75.2% for black women) (Arias et al.). Morgan (1996) asserted that decreased use of prenatal care by African American women has been related to care that was not culturally congruent. In this study (Morgan) professional prenatal care was seen as necessary, but there was a distrust of non-caring professionals, and barriers to such care. Culturally congruent care meant protection, presence, and sharing. Social structural factors that greatly influenced health were spirituality, kinship, and economics.

The importance of spirituality and social support have been documented for childbearing and childrearing African American women (Brodsky, 1999; Coffman & Ray, 2002; Wilson, & Miles, 2001). Brodsky described the components and process of resilience among urban, African-American, single parents. Spirituality was an important resource for most women. Furthermore, support from friends came largely through church based friendships. Spirituality has also been described as a source of support for African Americans who are grieving (Clements et al., 2003) and as a powerful coping strategy for African American mothers of a seriously ill infant (Wilson, & Miles).

Perinatal Loss and African American Parents

Several early investigators, using structured tools for data collection, found no relationship between race and intensity of grief response (Best, 1981; Lake, Johnson, Murphy, & Knuppel, 1987). However, the conclusions of another investigator (Bradshaw, 1986) raised the possibility that using a structured tool could have been an inadequate measure of outcome for the black subjects for the previous studies. In her study with 26 low-income women of various

racial backgrounds, Bradshaw found no differences in the grief response of women who had received an intervention strategy. She recommended further descriptive research with women of low socioeconomic status because of the lack of information on how other stressors could influence their emotional health and impact their perception of the loss. A more recent study of Caucasian, African American, and Hispanic American parents (DiMarco, Menke, & McNamara, 2001) has demonstrated ethnic differences in response to perinatal loss. These investigators reported that significantly more Caucasians attended a support group than other ethnic groups and that Caucasian participants had lower detachment scores (defined as being withdrawn and avoiding others) than the ethnic minority participants.

Experiences of African American mothers following a perinatal loss have been described recently (Van, 2001; Van & Meleis, 2003). In the first study, the healing processes used by 10 African American women after a pregnancy loss were examined (Van). The first strategy was 'putting it aside' whereby women avoided thoughts about the loss. The second strategy, 'there was a purpose' was an attempt to find meaning in the loss. The third strategy was 'heal yourself,' which were actions women used to get better. The final strategy 'he's in a good place' was based on religion and spirituality. In the second report, coping strategies of 20 African American women after a pregnancy loss were examined (Van & Meleis). The findings were categorized into coping with: (a) personal reactions and responses, (b) reactions of others, (c) memories of the baby, and (d) subsequent pregnancies. Personal reactions included: "I talked", "I haven't dealt with it", "I prayed", and "going inside myself." Reactions of others, such as coworkers, were often unsupportive, and mothers coped by excusing, praying for, or overlooking these behaviors. Some mothers had received mementoes of their infants or had planned a funeral service and these were a source of comfort. Finally, either planning for or experiencing a subsequent pregnancy was a source of fear or anxiety.

First callout Studying the responses of African American parents is critical because of the high incidence of perinatal loss among blacks, and the other stressful life situations that low-income, African Americans often experience.

Methods

A phenomenological approach was used in this research. The goal of phenomenology is to describe the meaning of human experience from the individual's perspective (Cohen & Omery, 1994; Giorgi, 1986).

Participants

For this study, a sample of 23 parents (17 families consisting of 17 mothers and 6 of their partners) was recruited from three hospitals that provide high-risk perinatal care, and also via an advertisement in a newspaper. The newspaper ad was used to recruit subjects from a larger variety of hospitals. However, only one mother was recruited by this method. The inclusion criteria were: (a) a parent who had recently (within four months) experienced a pregnancy loss at 16 weeks gestation or greater or a neonatal death; (b) African American; (c) low-income, as defined by the mother being a recipient of Medicaid; and (d) at least 18 years of age. The sample for this proposed study was larger than the typical sample size for a phenomenological study. A larger sample size was planned for the proposed study due to potential attrition of participants between the initial and subsequent interviews and the goal to explore the impact of other stressors on the participant's response to the loss.

Eleven parents reported having other living children. Six parents had prior perinatal losses; for three parents, the current perinatal loss was their fourth loss, and for three parents, the loss was their second loss. Two additional parents had elective abortions because of family pressure. For the 17 families, 11 experienced pregnancy loss (stillbirths) between 17 and 37 weeks

gestation ($M = 26.27$) and six experienced neonatal deaths of infants weighing between 486 and 3208 grams. These neonatal deaths occurred between one and 28 days of age and included one twin. Additional demographic information on the parents is in Table 1.

Data Collection

The study was approved by the Institutional Review Boards at the participating institutions. The investigator only contacted those parents who gave their permission through a hospital staff person or who responded to the advertisement. If the parent agreed to participate in the study, an appointment was made for the initial interview. Before the actual interview began, written, informed consents were obtained. The investigator (first author) used her expertise with interviewing bereaved parents to assess for psychological distress. She then used strategies, throughout the data collection process, to minimize psychological risk to participants (Kavanaugh & Ayres, 1998). For example, the investigator began each data collection session by establishing a relationship with the parent. One very successful strategy was to look at their baby's mementoes before starting the interview.

All interviews were tape recorded and were conducted separately with each parent, privately and in person by the investigator (first author). With the exception of three parents who requested to be interviewed in an office in the university, data collection occurred in parents' homes. During the initial interview, the interviewer elicited the parent's description of the experience using an open-ended format and began with the broad question, "What has it been like for you to lose your baby?" Then, attentive listening, which is critical for intuiting in a phenomenologic study (Swanson-Kauffman & Schonwald, 1988), and probes were used to assist parents to talk freely about their experience. Each interview concluded with the question, "What was it like for you to participate in this interview today?" At the end of the initial interview, each parent completed a sociodemographic questionnaire. Initial interviews were conducted with all 23 parents between 5 and 17 weeks after the loss and follow-up interviews with 21 parents who could be reached. Therefore, a total of 44 interviews were conducted. The follow-up interview was conducted to add or clarify any information obtained during the initial interview. The follow-up interview was held as soon as the investigator reviewed the initial interview and was able to schedule the interview with the participant. The time between the initial and follow-up interview ranged between two and five weeks.

Each parent received \$25.00 for participating in each interview, a personalized thank you letter and sympathy card, and either a Memory Box™ for displaying their baby pictures and storing mementoes, or if they already had a container for mementoes, several perinatal loss books, which they selected from a catalog. Without exception, parents immediately placed their baby's pictures and other mementoes in the box. Parents appreciated these incentives, but valued the experience of being able to talk to an attentive listener who was knowledgeable about perinatal loss. Therefore, parents reported that participating in the study was beneficial. Parents identified the following benefits: emotional relief (4 fathers, 7 mothers), a unique opportunity to talk (3 fathers, 12 mothers), an opportunity to help others (2 fathers, 1 mother), a better understanding of their experience (2 fathers), and evidence that someone cared about them (2 fathers). Even though some interviews were only 30–40 minutes in length, parents explained that they had talked more than they ever did about their loss. One mother whose interview was the shortest at 30 minutes, said, "It was really nice because I've been holding a lot of things in...So it was kind of easy to get a lot of stuff off my chest. (sic) Because I had just had so much bottled up because there was nobody really to talk to."

Data Analysis

The interviews were transcribed, typed verbatim, and then checked against the original taped interviews for accuracy. The approach for data analysis was adapted from the steps described

by Colaizzi (1978). Initially, all transcripts were read to develop an overall understanding of the experience. Then, significant statements were identified, coded and grouped into broad topical areas. Next, case summaries were created for each parent subject using the coded data and the demographic forms. Codes were then clustered into themes which were common to all participants. This step was accomplished by a process called 'free writing,' which captures the investigator's intuited experience. Then, the themes were validated with the original interviews. A description of each theme and subtheme was generated; finally, the exhaustive description of the phenomenon was formulated.

A number of measures that address issues of trustworthiness of qualitative data, as described by Lincoln and Guba (1985), and applied to phenomenological research (Beck, 1994), were performed. First, peer debriefing, which is interaction of the investigator with other professionals, was used. Consultants, which included nurse researchers and perinatal nurse specialists with clinical expertise with the population of parents, were used during the study. Second, thick descriptive data were collected, which is the process of providing a detailed description of the participants to allow for transferability of findings. Third, the investigators established an audit trail, which is a mechanism of maintaining extensive notes which reflect the investigators' analytical thought process during the course of the study. Finally, member check was performed. For a phenomenological study, member check is performed by validating the essential structure of the phenomenon (the summative description of the themes) as experienced by all participants with the participants (Colaizzi, 1978). For this study, a limited member check was performed with one participant after data were collected from the first six families. The intention was to perform a member check with each participant. However, it was not possible to do so because it was difficult to maintain contact with the participants either by telephone or written communication due to the long duration of the recruitment phase of this study (five years) secondary to uncontrollable delays in recruitment. For the member check, the participant read a several page summary of a description of the preliminary themes that had been developed from the first six cases. This participant responded that the summary "was really good."

Results

Analysis of the data revealed four themes common to all parents. These are: (a) recognizing problems and responding to the loss, (b) dealing with stressful life events, (c) creating and cherishing memories of their infant, and (d) living with the loss.

Recognizing Problems and Responding to the Loss

Parents experienced intense feelings surrounding their loss and were often unprepared for the baby's death, even in those situations when the mother had experienced serious problems during her pregnancy. To parents, events unfolded very rapidly. Parents clearly recalled the events that occurred prior to their baby's death, especially the day of their infant's death. One father said, "I can remember the entire day vividly." Parents who had experienced other perinatal losses were in disbelief that they were experiencing another loss, and compared their current experience to previous ones.

Misreading symptoms of pregnancy complications.—Most mothers who were diagnosed with preterm labor or prolonged rupture of the membranes were not aware of the symptoms accompanying these conditions, and misread their symptoms as something else, such as urinary incontinence (for ruptured membranes) or gas pains (preterm labor pains). One father said,

Not even with the water being broken, I guess for a week prior, you know, we thought the baby was just sitting on her bladder and that she had no control of that, but, you

know, that's the kind of stuff that comes with being first time parents and really not knowing what happens.

Although some of these mothers had prior labor experience, they could not believe that labor could begin as early as it was occurring. Several parents questioned their own mothers about their symptoms and were reassured that they were normal symptoms of pregnancy. Other mothers knew that something was wrong during the pregnancy and verbalized their concerns, such as abdominal pressure, to health care professionals. However, they felt that the health care professional minimized the mother's physical complaints or offered conflicting advice.

Maintaining hope during medical treatment.—Although parents had experienced prenatal complications, they maintained hope for their infant. Even those parents who were told of their infant's death in utero were hoping that there would be a miracle and they would hear their baby's cries at the time of birth. One mother recalled, "I held her and I kissed her and I was just praying for a miracle. 'Please open your eyes. Please open your eyes.'"

Parents were also hopeful because of their expectations for aggressive treatment for their infant. They were familiar with advances that have been made in perinatal care through their own prior experiences, those of family members, or those publicized in the media. One parent made the comparison to knowing people "who were shot in the head and lived." Only those parents whose infants lived for more than one week and had the opportunity to participate in treatment decisions for their infant realized that their infant might die. One of these mothers said:

They said there was no guarantee that she would live, not unless there was a miracle. I was believing in a miracle because I do believe in God and I had faith.... But then there was really nothing else more they could do for her because she wasn't getting better.... The next day when I went to see her, she was real blowed (sic) up. I couldn't see that was her. So I told them, I guess it's time to take her off the machines because they said that her organs were so damaged, and there was nothing else they could do for her.

Feeling intense emotions after the death.—Parents used words such as "shocked" to describe how they felt when their infants died. One mother who gave birth to a stillborn infant said, "that was the worst moment in my life, the worst moment in my life." Mothers described themselves as being "very emotional" and even feeling like they were "going crazy." A number of mothers recalled suicidal thoughts because of their desire to be with their infant. They cried very often, had insomnia, and initially did not want to talk about the loss. Mothers appreciated nurses who could just sit and hold their hand, especially if the mother was alone. One mother said,

When she found out I lost the baby, she just held me like I was a baby, rubbed my back and everything. She just was real helpful. I don't know if I would have made it. I will never forget her name. I know their job is to be a nurse, but she didn't have to be that nice. She was really nice. She stayed there until my mother came. She sat in the room until my mother came. I felt much better.

Both mothers and fathers recognized the "special attention" that most nurses gave them with regards to their physical, emotional, and spiritual needs, and they found this attention very helpful. One mother shared that she was moved when she found out that one physician came in from home to tell her of her infant's death.

A number of mothers also described themselves as being very irritable and "ready to fight" with others, especially with those health care professionals whom mothers perceived as uncaring. Most mothers were themselves hospitalized at the time of their infant's death, and they wanted to be discharged home as soon as possible. Guilt was prevalent in mothers of

stillborn infants and those mothers who had been treated for pregnancy complications. One mother recalled how she felt when she heard of her infant's in utero death. She said, "Its like my heart stopped and I kept saying, 'I killed my baby. I killed my baby.'" These mothers questioned if they should have sought health care in a timelier manner.

Fathers explained that they hurt badly also and felt a loss of control. One father said, "It has been the worst thing that has happened that I have had no control over." Fathers often kept their emotions under control for fear of further upsetting the mother, but yet were also unsure how to support their partners. Mothers wanted fathers to express their emotions. One mother said, "I wish he would open up to me just once, just once to show me he can cry, too. This was your baby. Don't be embarrassed. You can cry, too. You feel the pain, too."

Dealing with Stressful Life Events

During the pregnancy and surrounding the time of the loss, parents were dealing with stressful life events that were not all related to the loss, such as economic hardship and deaths of other close family members. Many mothers described an overall feeling of being stressed out during their pregnancy. Several of these mothers felt that their stress contributed to their poor pregnancy outcome. Only one mother was receiving mental health counseling during her pregnancy, and an additional few had expressed a need to seek professional care.

Second callout During the pregnancy and surrounding the time of the loss, parents were dealing with stressful life events that were not all related to the loss, such as economic hardship and deaths of other close family members.

Feeling abandoned or unsupported.—Some mothers described feelings of abandonment or not being supported by family, which caused them stress. For several mothers, arguing with their family members was so intense that the mothers relocated to other housing.

Several mothers explained that they were deserted by the father of the baby when they became pregnant. Several other mothers felt abandoned by their own parents who had been separated from them because of the parents' substance abuse or incarceration. As a result, their own parents were not there for them during the pregnancy. One of these mothers had the extra responsibility of raising her three younger siblings because of her mother's substance abuse.

Suffering economic hardship.—Parents described economic factors that caused them stress; for mothers the stress was especially prevalent during their pregnancy. Many mothers expressed concerns about their housing or living arrangements, either because they were unhappily living in another relative's home, or they were concerned about the safety of their neighborhood. Many parents also described the stress of either not having a job or being employed under adverse working conditions, which the mothers felt contributed to their poor pregnancy outcome. For example, one of these mothers explained that she saved enough money to live on her own and in a safer neighborhood, which was especially important for her son and fourteen year old brother whom she was also raising. Then, she discovered that she could barely pay the bills and feed her family, so she secured a job on the night shift at a fast food restaurant. As her pregnancy progressed, she found it more difficult to work, but she described why she had to keep her job. She said,

I was still stressed out. I wasn't getting any rest. It was eight long laboring hours. When I would go to the clinic, they were like you are doing too much. You have to stop. You aren't giving the baby enough rest.... I didn't really say to them what I was really going through, I had to have this job. It was the only way to support the kids and keep eating. ... The money I was getting from public aid was barely enough just to pay the rent... So it would be two or three days I went without eating, just drinking

water and eating crackers. One of them days I felt the baby move. I said, “Oh, you still surviving in there?”

Experiencing others’ serious illnesses or untimely deaths.—Many mothers had close relatives who were seriously ill and/or hospitalized during the mother’s pregnancy. Also, parents had already experienced deaths of other close family members or friends. These included untimely deaths of their: other infants, nieces and nephews, siblings, parents, grandparents, and young male cousins or friends. One father, who was a substitute high school teacher, experienced the deaths of three of his students in one year. Even if the death had not been recent, the current loss triggered painful memories of the other deaths.

Despite their young age, nearly half of the parents had already lost their own parent or parent substitute, such as a grandparent who had raised them. These relatives had experienced an untimely death which occurred while the person was in their 30s, 40s, or 50s. For several, these deaths occurred during the mother’s pregnancy. One of these parents whose own mother and grandmother died during the woman’s pregnancy said, “I can’t take this anymore, another death. I don’t know how I would take it.” A number of parents were survivors of homicidal or suicidal deaths of young male relatives or close friends, which one parent described as “devastating.” One of these fathers explained that his closest friend was murdered two weeks after his baby was buried. This father could barely talk during the interview when he recalled “my best friend, my very first friend was killed.”

Receiving unfair treatment.—Parents perceived that they received unfair treatment mostly related to medical treatment. However, one mother also reported unfair treatment by a funeral director who buried her son sooner than she was told initially and without her presence. According to the mother, the funeral director said that he did not know she wanted to be present for the burial. She said, “I was mad with them because they buried my son without me being there How did he not know that I wanted to go.... My heart just dropped. ... Like I didn’t even get a chance to say goodbye to him.”

Many parents perceived that the mother or the infant received inadequate care, and many attributed this unfair treatment to their insurance status (medical assistance), and therefore perceived a disparity in treatment. One mother whose baby died after being in the neonatal intensive care unit said,

I feel they didn’t do the best that they could. I think it was because of financial. I was on public aid and so much that certain doctors will only do so much for people on a medical card. ... But if I had a better insurance, they would have put their heart and soul into it and I feel they didn’t.

One father said, “I know she (his wife) was not in a private room and was not a VIP patient, but, to me she was a VIP patient.” This father also perceived unfair treatment with regards to paternity establishment. He shared that his name did not appear on his son’s birth certificate even though he provided the requested documentation that he was married to the mother of his son.

Parents recalled the frustration of seeing many different medical providers at their prenatal visits and thus felt they did not have the continuity of care they would have had with a “personal doctor”. Some parents were also very stressed by overhearing health care professionals make negative comments about the mother or argue with each other about the plan of care. One mother who overheard a conversation said, “I could hear the doctor in another room telling him, ‘You should have caught this. Why wasn’t she given a circlage right away?’ I could hear him fussing down the hall.” Another mother was very angry at how she was treated during an evaluation for her pregnancy complications. Because she was worried about child care

arrangements for her young daughter at home, she questioned if hospitalization for herself was necessary. She said, “‘Well, do I have to stay?’ The doctor said, ‘Well, I don’t have a gun to your head.’ So I was like ‘You don’t have to have a gun to my head to make me stay.’”

Parents expected that health care providers would listen to their concerns and that technology would be applied in their situations. All parents whose live born infants were born at the threshold of viability felt that their infants did not receive adequate resuscitative attempts right after birth. One mother said,

When he was born alive, you should have done something... I felt that my son had a chance and they did not give him a chance... When they took him out, they should have rushed him away and said they would see what they could do.

These parents then were dismayed and angry when the care they received was not consistent with the care they expected, especially because they were in a hospital that specializes in high-risk perinatal care and the mothers felt they had done everything prescribed during the pregnancy. Several of these parents even made initial inquiries about a medical malpractice lawsuit.

Creating and Cherishing Memories of their Infant

Parents created memories of their infant by being physically close, noticing family resemblances, and obtaining keepsakes of their infant. Parents cherished these memories. Parents also decided on infant burial, which typically included a memorial or prayer service and provided another memory for the parents.

Being with their infant.—Except for one father of an infant born with gastroschisis, all parents saw their infants. Whereas most mothers readily embraced their infants, most fathers were very reluctant to hold their infants but did with encouragement from nurses or family members. Only two mothers had no physical contact with their infant; one of these mothers explained that she was too heavily medicated at the time to hold her infant. With the exception of one, mothers of the live born infants held their infants at the time of death. Typically, this time was the only time they had an opportunity to hold their infants. During the interviews, these mothers very tearfully recalled this time, and often needed a break from the interview to regain composure and continue. One mother recalled,

And I held him until about four hours. I said I loved him. I don’t want him to go, but God knows best. Maybe I’m struggling so much that it would be just another burden and He knows best. Well I was just holding him and crying. I said, somewhere you get a chance to see your grandma. I tell him when you get there when you get old enough you think about me.

Parents cherished the time they spent with their infants, and were often allowed to keep their infant with them in the mother’s room for many hours. Family members were often present, and also took turns holding the infant. One mother described the scene as a “room full of love”. Most mothers were very averse to having the nurse take the infant away before they were ready, and some even described intense verbal arguments that occurred between them and family members or nurses who were persistent.

Noticing family resemblances.—Many parents identified family resemblances between their infant and themselves. Family resemblances were of physical or behavioral characteristics. Even the mother of the infant who was born at 17 weeks gestation thought that her son resembled his father. Although parents happily described these resemblances during the interview, parents explained that noticing these resemblances initially often made the parent cry.

Gathering keepsakes of and remembering their infant.—Parents received many different types of keepsakes from the nurses, including pictures, identification bracelets, blankets, certificates of life, ointment, gowns, and caps. Parents especially treasured any items that had been worn by or come in contact with their infant. Parents reported that they were grateful for the pictures and especially appreciated nurses who took extra time for carefully arranged photographs. Parents wanted more pictures, especially pictures of their infant after being cleaned up. Parents of live born infants wanted pictures of their baby before the baby's death and without any equipment on the infant, particularly on the face. All infants were named, and one mother chose to remember her infant by having her infant's name tattooed on her body.

Deciding on infant burial.—Parents had to decide on a private or a 'hospital' burial for their infant, and found it very helpful to receive accurate information about these options from health care professionals. A hospital burial was actually carried out by the county and involved burying the infant in a common burial site, which would be 'unmarked' to the parent. Ten families selected hospital burial, even though some of them were advised by family members to arrange for a private burial, and 7 families selected a private burial. Parents explained that they based their decision on what they felt was best for them and their infant. Some parents were concerned about the emotional distress of a private burial, yet others were distressed when they realized what was involved with a hospital burial. One mother was comforted to hear that her infant would be buried with others with a hospital burial, yet another mother thought her infant was "too special" to be buried with others. A few parents were not aware that they could receive financial assistance for a private burial, so economics was a factor in their decision. Even those parents who decided on a hospital burial often selected the burial clothes for their infant and participated in a memorial prayer service which, they very gratefully described, was conducted by the hospital chaplain. Parents found that prayer and reading from scripture were very comforting at the time of the memorial service.

Living with the Loss

In the weeks and early months following the infant's death, parents had to find ways to live with their loss. Parents encountered difficult situations, relied on their spirituality, sought diversions and support, tried to make sense of the loss, and thought about a subsequent pregnancy.

Encountering difficult situations.—Most mothers found it very difficult to be in situations that reminded them of "what could have been" if their infant had not died. Examples of these situations were being around pregnant women or infants, attending baby showers, and celebrating holidays that recognize mothers and involve dressing up children, such as Mother's Day, Christmas, and Easter. For these holidays, mothers explained that they either stayed at home or went out with their partner only late in the evening. Although fathers did not describe any problems with these situations, one father found it difficult to complete job applications which requested information on number of children.

Relying on their spirituality.—Many parents relied on their spirituality to deal with their loss. For some parents, this was in the form of praying, for others it was returning to church. Parents described "turning their lives over to the Lord." One mother said, "If I hadn't started back praying and had the faith and keeping the faith, I don't think I would have been able to do it." Some mothers were comforted to think that their infant was in heaven surrounded by angels. For these parents, it was very helpful to receive religious books after the loss.

Seeking diversions and support.—Parents searched for activities to keep busy and divert themselves from dwelling on the loss, and also sought social support. Fathers, more so than mothers, found it helpful to keep busy and to "move forward" and think positively about the

future. Mothers with other children explained that having other children to care for helped them. Parents sought support from family and friends and described a number of ways that family and friends supported them. The most common support they received was emotional support. The various types of support that parents received is described in detail elsewhere (Kavanaugh, Trier, & Korzec, 2004). Although parents were aware of support groups in their area and expressed an interest in attending, none had participated in them. Some parents would consider attending if their partner attended with them, and others felt it might be something they would need more of in the future.

Making sense of the loss.—Parents tried to make sense of their loss and determine why it occurred. Parents who consented for an autopsy were waiting for the final results; a number of them were very frustrated with the delay in receiving the results. Parents also questioned the death from a philosophical or spiritual perspective, and explained it within a religious context. Many mothers could not understand why the loss occurred, especially because they believed they had done everything to insure a healthy pregnancy and had known others who abused drugs and had healthy babies.

Contemplating future pregnancies.—Parents talked about their plans for a future pregnancy, and several mothers shared that they were possibly pregnant at the time of the second interview. Most parents planned on a subsequent pregnancy in the future, yet those with more than one perinatal loss were very reluctant to plan for another pregnancy. One father explained how a healthy subsequent pregnancy would be part of the healing. He said, “And once it is all right, that will even take away some of the sorrow and pain from the previous times.” However, some mothers did not plan on a subsequent pregnancy because of their concern about their ability to deal with another perinatal loss. Those parents who planned on a subsequent pregnancy described the measures they would take to insure a healthy outcome. These included delaying a pregnancy until there was less stress in their lives and planning on having one consistent primary care provider.

Discussion

Several themes and subthemes described in this study are consistent with findings in perinatal loss research. For example, the subtheme that parents experienced intense emotions after the death and the reported gender differences is well supported in other research (Kavanaugh, 1997a; Lin & Lasker, 1996). The gender differences in this research included the tendency for the father to guard his display of emotions because of his concern for his partner, his inability to talk about his feelings, and his need to move on with his life. Also, the behaviors that parents found helpful, such as physical presence, are similar to the caring behaviors reported in other perinatal loss research (Lemmer, 1991; Sexton & Stephen, 1991) and described in a middle range theory of caring (Swanson, 1991, 1993). However, the subtheme that parents misread symptoms of pregnancy complications has not been well described in the loss research. It is unclear if this finding is unique to this group of parents or, more probably, that this phenomenon has not been examined in other perinatal loss research. The theme of dealing with other stressful life events has also not been reported in the perinatal loss literature. However, it is possible that this finding is related to the parents’ race and/or economic status because of the results of prior research on the prevalence of stress in African Americans (Brodsky, 1999; Schultz et al., 2000).

Parents in this study often were dealing with other stressful events, such as untimely deaths of others close to them, and everyday stress, such as that described by Brodsky (1999). These findings are supported in a study of stress and health status among African American and white women (Schulz et al., 2000). In that study, African American women reported more frequent acute life events and unfair treatment than white women. Mothers in the research reported here

also felt that their stress was cause of their adverse pregnancy outcome. This finding is noteworthy because of the link between perceived stress and preterm birth (Mackey, Williams, & Tiller, 2000; Stancil, Hertz-Picciotto, Schramm, & Watt-Morse, 2000). Also, of particular importance in this study is that nearly 40% of the parents had experienced prior perinatal losses.

Some of these stressful events were clearly related to parents' economic status, such as inadequate housing. Parents also perceived that some stressors, such as perception of unfair medical treatment, were due to their economic status, but it is not clear if other findings are related to race and/or economic status. The reports of treatment disparity are parents' perceptions after a very adverse event; no attempt was made to compare their perceptions with the type of treatment that was provided by health care professionals. It is possible that parents did receive unfair treatment, especially because of the prevalence of racial and ethnic treatment disparity in health care (Smedley, Stith, & Nelson, 2003). However, it is also possible that parents had a different understanding of the events than the health care professionals.

The importance of spirituality in the lives of bereaved parents has been established in prior research (Calhoun, 1994; Sexton & Stephen, 1991) and is also supported in this research. Parents relied on their spirituality to cope with and also to make sense of their loss. These findings are also similar to other research on African American parents who experienced a perinatal loss (Van, 2001; Van & Meleis, 2003). For parents, their spirituality sustained them through this very difficult time.

Recommendations for Practice and Research

These findings demonstrate the importance of recognizing the incidence of other life stressors that parents may be experiencing that could impact on pregnancy outcome or on their response after a loss. Nurses and other health care professionals should assess the presence of stressors in a parent's life, recognize the ways that parents cope, assist them to mobilize coping strategies, and make appropriate referrals as needed.

Third callout Nurses and other health care professionals should assess the presence of stressors in a parent's life, recognize the ways that parents cope, assist them to mobilize coping strategies, and make appropriate referrals as needed.

Specifically, this study illustrates the need for nurses to assess vulnerable pregnant women and their partners for increased instances of life stressors such as the death of a close family member or undesirable employment conditions. The presence of multiple life stressors may have contributed to pregnant women's reported difficulty in identifying or responding to symptoms of pregnancy complications in this study. Nurses may need to implement additional measures of empathy, communication, education, and, when appropriate, social support referrals to optimize the health of these women.

This study also explicates the importance of nursing actions and interventions when parents experience a perinatal loss. Foremost, nurses can increase parents' coping abilities during times of perinatal loss by providing a supportive environment for parents. Examples of specific nursing actions include displaying empathy and providing parents with pictures and mementoes of their deceased baby. These findings coincide with other research examining perinatal loss (Kavanaugh, 1997b; Lundqvist, Nilstun, & Dykes, 2002). This study also accentuates the need for nurses and other health care professionals to articulate clear and explicit information regarding hospital-based policies, such as autopsy and burial practices, to both mothers and fathers. Significant to this study was the finding that fathers were reluctant to hold their infant without additional encouragement from nursing personnel, thereby indicating the special attention that fathers may need.

It is also noteworthy that in this research no parents attended parent support groups. This finding suggests that traditional mechanisms for providing follow-up care for parents after a perinatal loss, such as making a referral to a hospital-based parent support group, may not be appropriate for these parents. The results of the research reported here do support the need to find out parents' understanding of events and desire for involvement in treatment plans.

This study provides a beginning understanding of the experience of low-income, African American parents and is not designed to be generalized to a larger population. Further research needs to be conducted with a larger sample of low-income African American parents. Also, the methodological approach for this study does not allow for inferences to be drawn regarding the influence of race and/or economic status on the study findings. A larger correlational study could be designed to study those variables. Further research is also needed to compare parent and health care professional perceptions of treatment decisions, including parent desire for involvement in treatment decisions.

Conclusion

Limited research has been done on the experience of perinatal loss in low-income, African American parents. The results of this study provide a beginning understanding of the loss experience in this group of parents, and demonstrate some experiences that have not previously been reported in the perinatal loss research literature. Nurses and other health care professionals should be aware of the presence of additional stressful events that parents may be experiencing and intervene appropriately.

Acknowledgements

This study was funded by the National Institute of Nursing Research (1R15NR4224-1A1). The authors gratefully acknowledge the parents who shared their stories and the following individuals: Kristen M. Swanson, PhD, FAAN, Erin Powell Carmany, MS, CGC, Vida Bartasius, MS, CNM, Judy Friedrichs, MS, RN, and Virginia Maikler, PhD, RN for their assistance with this study.

References

- Arias E, MacDorman MF, Strobino DM, Guyer B. Annual summary of vital statistics-2002. *Pediatrics* 2003;112(6):1215–1230. [PubMed: 14654589]
- Beck CT. Reliability and validity issues in phenomenological research. *Western Journal of Nursing Research* 1994;16:254–267. [PubMed: 8036802]
- Best, E. K. (1981). *Grief in response to prenatal loss: An argument for the earliest maternal attachment*. Unpublished doctoral dissertation, University of Florida, Gainesville.
- Bradshaw CT. The grief response to fetal loss among lower SES women: A comparison of brief vs. extended intervention. (Doctoral dissertation, Virginia Commonwealth University, 1985). Dissertation Abstracts International 1986;46:2453B.
- Brodsky A. "Making it": The components and process of resilience among urban, African-American, single mothers". *American Journal of Orthopsychiatry* 1999;69(2):148–160. [PubMed: 10234381]
- Calhoun LK. Parents' perceptions of nursing support following neonatal loss. *Journal of Perinatal and Neonatal Nursing* 1994;8(2):57–66. [PubMed: 7815319]
- Clements PT, Vigil GJ, Manno MS, Henry GC, Wilks J, Das S, Kellywood R, Foster AW. Cultural perspectives of death, grief, and bereavement. *Journal of Psychosocial Nursing* 2003;41(7):18–26.
- Coffman M, Ray M. African American women describe support processes during high-risk pregnancy and postpartum. *Journal of Obstetric, Gynecologic, and Neonatal Nursing* 2002;31:536–544.
- Cohen, M.Z., & Omery, A. (1994). School of phenomenology: Implications for research. In J.M. Morse (Ed.), *Critical issues in qualitative research methods* (pp. 136–156). Thousand Oaks, CA: Sage.
- Colaizzi, P. F. (1978). Psychological research as the phenomenologist views it. In R. S. Valle & M. King (Eds.), *Existential-phenomenological alternatives for psychology* (pp. 48–71). New York: Oxford University Press.

- Crowther ME. Communication following a stillbirth or neonatal death: Room for improvement. *British Journal of Obstetrics and Gynecology* 1995;102:952–956.
- deMontigny F, Beaudet L, Dumas, L. A baby has died: The impact of perinatal loss on family social networks. *Journal of Obstetric, Gynecologic, and Neonatal Nursing* 1999;28:151–156.
- Dimarco MA, Menke EM, McNamara T. Evaluating a support group for perinatal loss. *MCN: The American Journal of Maternal/Child Nursing* 2001;26:135–140.
- Fulmer, R. H. (1987). Special problems of mourning in low-income single-parent families. In J. C. Hansen (Series Ed.) & M. Lindblad-Goldberg (Vol. Ed.), *The family therapy collection: Vol. 23. Clinical issues in single parent households* (19–37). Rockville, Maryland: Aspen.
- Giorgi, A. (1986). Theoretical justifications for the use of description in psychological research. In P.D. Ashworth, A. Giorgi, & A. J. J. de Koning (Eds.), *Qualitative research in psychology* (pp. 3–22). Pittsburgh: Duquesne University Press.
- Kavanaugh K. Gender differences among parents who experience the death of an infant weighing less than 500 grams at birth. *Omega: The Journal of Death and Dying* 1997a;35:281–296.
- Kavanaugh K. Parents' experience surrounding the death of a newborn whose birth is at the margin of viability. *Journal of Obstetric, Gynecologic, and Neonatal Nursing* 1997b;1:43–51.
- Kavanaugh K, Ayres L. "Not as bad as it could have been": Assessing and mitigating harm during research interviews on sensitive topics". *Research in Nursing & Health* 1998;21:91–97. [PubMed: 9472241]
- Kavanaugh K, Trier D, Korzec M. Social support following perinatal loss. *Journal of Family Nursing* 2004;10:70–92. [PubMed: 17426820]
- Lake MF, Johnson TM, Murphy J, Knuppel RA. Evaluation of a perinatal grief support team. *American Journal of Obstetrics and Gynecology* 1987;157:1203–1206. [PubMed: 3688076]
- Lang A, Gottlieb L. Parental grief reactions and marital intimacy following infant death. *Death Studies* 1993;17:233–255.
- Lemmer CM. Parental perceptions of caring following perinatal bereavement. *Western Journal of Nursing Research* 1991;13:475–493. [PubMed: 1897209]
- Lin SX, Lasker JN. Patterns of grief reaction after pregnancy loss. *American Journal of Orthopsychiatry* 1996;66:262–271. [PubMed: 9173804]
- Lincoln, Y. S., & Guba, E. G. (1985). *Naturalistic inquiry* Beverly Hills: Sage Publications.
- Lundqvist A, Nilstun T, Dykes AK. Both empowered and powerless: Mothers' experiences of professional care when their newborn dies. *BIRTH* 2002;29:192–199. [PubMed: 12153650]
- Mackey M, Williams C, Tiller, C. Stress, pre-term labour and birth outcomes. *Journal of Advanced Nursing* 2000;32:666–674. [PubMed: 11012810]
- Madden ME. The variety of emotional reactions to miscarriage. *Women & Health* 1994;21(23):85–104.
- Mekosh-Rosenbaum V, Lasker JN. Effects of pregnancy outcomes on marital satisfaction; A longitudinal study of birth and loss. *Infant Mental Health Journal* 1995;16:127–143.
- Morgan M. Prenatal care of African American women in selected USA urban and rural cultural contexts. *Journal of Transcultural Nursing* 1996;7(2):3–9. [PubMed: 8974415]
- Sexton PR, Stephen SB. Postpartum mothers' perceptions of nursing interventions for perinatal grief. *Neonatal Network* 1991;9(5):47–51. [PubMed: 1944093]
- Schulz A, Israel B, Williams D, Parker E, Becker A, James. Social inequalities, stressors and self reported health status among African American and white women in the Detroit metropolitan area. *Social Science & Medicine* 2000;51:1639–1653. [PubMed: 11072884]
- Smedley, B.D., Stith, A.Y., & Nelson, A.R. (Eds.). (2003). *Unequal treatment: Confronting racial and ethnic disparities in health care*. Washington, DC: The National Academies Press.
- Stancil T, Hertz-Picciotto I, Schramm M, Watt-Morse, M. Stress and pregnancy among African-American women. *Paediatric and Perinatal Epidemiology* 2000;14:127–135. [PubMed: 10791655]
- Stinson KM, Lasker JN, Lohmann J, Toedter LJ. Parents' grief following pregnancy loss: A comparison of mothers and fathers. *Family Relations* 1992;41:218–223.
- Swanson KM. Empirical development of a middle range theory of caring. *Nursing Research* 1991;40:161–166. [PubMed: 2030995]
- Swanson KM. Nursing as informed caring for the well-being of others. *Image: Journal of Nursing Scholarship* 1993;25:352–357.

- Swanson KM. Effects of caring, measurement, and time on miscarriage impact and women's well-being. *Nursing Research* 1999;48:288–298. [PubMed: 10571496]
- Swanson-Kauffman, K. M., & Schonwald, E. (1988). Phenomenology. In B. Sarter (Ed.), *Paths to knowledge: Innovative research methods for nursing* (pp. 97–105). New York: National League for Nursing.
- Theut SK, Pederson FA, Zaslow MJ, Cain RL, Rabinovich BA, Morihisa JM. Perinatal loss and parental bereavement. *American Journal of Psychiatry* 1989;146:635–639. [PubMed: 2535615]
- Thomas V, Striegel P. –5). Stress and grief of a perinatal loss: Integrating qualitative and quantitative methods. *OMEGA* 1994;30:299–311.
- Van P. Breaking the silence of African American women: healing after pregnancy loss. *Health Care for Women International* 2001;22:229–243. [PubMed: 11814070]
- Van P, Meleis A. Coping with grief after involuntary pregnancy loss: Perspectives of African American women. *Journal of Obstetric, Gynecologic, and Neonatal Nursing* 2003;32:28–39.
- Wilson S, Miles M. Spirituality in African-American Mothers coping with a seriously ill infant. *Journal of the Society of Pediatric Nurses* 2001;6(3):116–122. [PubMed: 11529600]

Table 1

Demographic Characteristics of Parents (N = 23)

Characteristic	n
Age (years)	
Mothers: range 19–34 (M = 23.82)	17
Fathers: range 20–34 (M = 27.33)	6
Education range 11–17 years (M = 13.27)	23
Marital status	
Single, living with partner	13
Single, in a relationship	4
Married	3
Single mother, abandoned by partner	3
Yearly income (dollars)	
<5,000	10
5,000–10,000	8
10,001–15,000	1
20,000 or >	4 ^a
Religion	
Baptist	13
Catholic	2
Other	2
No Preference	6

^aThese parents (3 fathers and 1 mother) were kept in the study because the mothers met the sample inclusion criterion of being a recipient of Medicaid.