Comparison of Biomass Dry Weights and Radial Growth Rates of Fungal Colonies on Media Solidified with Different Gelling Compounds

M. REESLEV* AND A. KJØLLER

Department of General Microbiology, University of Copenhagen, DK-1307 Copenhagen, Denmark

Received 21 June 1995/Accepted 19 September 1995

Penicillium commune, *Aureobasidium pullulans*, and *Paecilomyces farinosus* were grown on two different media solidified with agar, Pluronic F-127, Carrageenan X-4910, or Carrageenan X-4910 overlaid with cellophane. Growth on Carrageenan X-4910 was generally the same as that on agar, as was the visual appearance of the colonies, e.g., the pigmentation. The Carrageenan X-4910 gels had a melting point, depending on the medium, of 41 to 46°C, and the dry weights of the colonies were readily determined at 60°C. To determine the dry weights of the colonies grown on agar plates, the gels were boiled for 10 min to melt the agar. Comparison of these two procedures showed that the boiling procedure resulted in a 22% reduction of the biomass dry weight. Cellophane membranes did not affect the radial growth rate profoundly. The biomass density was almost halved for *P. commune* and *P. farinosus* grown with membranes, whereas the presence of the membrane did not affect the biomass density of *A. pullulans*. The biomass densities of the colonies grown on Pluronic F-127 were significantly reduced, while in most cases, the radial growth rates of colonies grown on Pluronic F-127 were significantly higher than those obtained on agar or Carrageenan X-4910. Furthermore, the morphology of the leading hyphae was altered, and the hyphal growth unit length was more than twice that obtained on agar and Carrageenan X-4910. Carrageenan X-4910 is a valuable gelling compound for the study of the growth of fungi, as the biomass dry weight is readily determined and growth is similar to that obtained on agar gels.

The mechanisms involved in the control and regulation of mycelial growth are better studied on solid medium than in submerged cultures, as fungi are adapted to growth on solid substrates (11). Nevertheless, the published data concerning determinations of the biomass dry weights of fungal colonies grown on solid medium are scarce. This probably relates to the ease with which the radial growth rate (K_r), defined as $K_r = (R_1 - R_0)/(t_1 - t_0)$ (16), where R_0 and R_1 are the colony radii at time t_0 and t_1 , respectively, can be determined. Furthermore, the finding that the K_r of a fungal colony was related to the specific growth rate obtained in submerged culture by the equation $K_r = w \times \mu$, where w is the peripheral growth zone and μ is the specific growth rate (16), has led to a widespread use of K_r in assessing the effects of environmental conditions. However, K_r is a valid measure of the specific growth rate only so long as w is constant, which is not the case when the nutrient concentration varies or after treatments with inhibitors or paramorphogens (18, 20). The high melting point of the agar gels which traditionally has been used is another reason for the preference of determining the K_r instead of the biomass dry weight. It is necessary to boil the gel to separate the biomass from the gel, and this harsh treatment results in the degradation of the biomass (10). Also, resolidification of the agar during the filtering process can be a problem. These problems have been circumvented by growing the fungi on agar plates covered with cellophane membranes from which the biomass can be washed or otherwise removed for determination of the dry weight (3, 6, 13). Also, indirect methods for the evaluation of the fungal biomass of colonies grown on agar medium have been used, e.g., determination of the hyphal density by counting the number of hyphae crossing an arc of a known length (1,

15) or determination of the lengths and diameters of hyphae and assuming the density and water content (14). These indirect methods are time-consuming, and the fact that they are used emphasizes the need for a simple and direct method for the determination of biomass dry weight.

Carrageenan is a gelatinous substance extracted from marine algae (e.g., *Eucheuma cottonii*). Carrageenan has been tested as an agar substitute because of its lower cost (8, 19). For this study, we used a Na^+ salt of a κ -carrageenan which has a low melting point suitable for our purpose. Pluronic F-127 is a block copolymer of polypropylene oxide and ethylene oxide (4) and forms a semisolid gel at room temperature but is fluid at low temperature (4°C). The gelling temperature depends on the concentration. The compound has been described as being useful for the isolation of heat-sensitive bacteria (4) and antagonists against oomycetous fungi (9).

In the present investigation, the rates of radial extension and the biomass densities of colonies grown on Carrageenan X-4910, Carrageenan X-4910 overlaid with cellophane, and Pluronic F-127 were compared with those obtained on agar.

MATERIALS AND METHODS

Fungi. *Penicillium commune* (IBT 13713), *Aureobasidium pullulans* (QM 3092), and *Paecilomyces farinosus* (AGM 93001) were obtained from the culture collection of the Institute of Biotechnology, Technical University of Denmark (IBT); the University of Massachusetts, Amherst, Mass. (QM); and the Department of General Microbiology, University of Copenhagen, Copenhagen, Denmark (AGM).

Preparation of the solid media. Two media were used in the experiments. The V8 medium (The Campbell Soup Co., Camden, N.J.) was prepared as described earlier (7), although antibiotics were omitted. A defined medium, b medium, was prepared as described earlier (12), except that the glucose concentration was 10.00 g/liter and the concentration of $(\text{NH}_4)_2\text{SO}_4$ was 0.368 g/liter (100 mg of ammonium per liter). For solid media with Carrageenan X-4910 (Copenhagen Pectin A/S, Lille Skensved, Denmark) and agar (Bacto agar; Difco Laboratories, Detroit, Mich.), a final concentration of 15 g/liter was used. Pluronic F-127 (BASF Corporation, Parsippany, N.J.) was used at a concentration of 300 g/liter. After the medium was autoclaved, 15 ml was pipetted into sterile plastic petri

* Corresponding author. Mailing address: Department of General Microbiology, University of Copenhagen, Sølvgade 83H, DK-1307 Copenhagen K, Denmark. Phone: 45 35 322057. Fax: 45 35 322040.

dishes and allowed to dry. The pH was adjusted to 6.0 prior to autoclaving. The media solidified with carrageenan or agar were dispensed at 65 and 55°C, respectively. After being autoclaved, medium solidified with Pluronic F-127 was placed in the refrigerator at 4°C until it became liquid (the next day). The bottle containing the medium was then placed in an ice-water bath, and 15 ml was dispensed into sterile plastic petri dishes. When Carrageenan X-4910 medium was overlaid with cellophane (PT 300; British Cellophane Ltd.), the cellophane membranes were, prior to the autoclaving of the medium, boiled twice for 15 min in distilled water to remove plasticizers (13).

Inoculation and growth conditions. Conidia were harvested in 2 to 3 ml of 1% aqueous NaCl from cultures grown on V8 agar. One milliliter of this solution containing the conidia was mixed with 1 ml of semisolid agar solution (0.2%), and this suspension was used for one-point inoculation of the plates. The plates were incubated at 25°C in a glass desiccator (inner diameter, 30 cm) to which 500 ml of a glycerol solution had been added to maintain a relative humidity of 98% (5).

Measurements of biomass and K_r . The two growth media gelled with different gelling compounds were inoculated with the fungal test species. Colony diameters were measured with a ruler four to six times at intervals of 24 to 48 h to ensure that the colonies were expanding at a constant rate while their K_r s were being determined. If a colony was not circular, the average of the longest and shortest widths was recorded. The K_r was calculated by linear regression of each colony radius versus time. After the last measurements, typically when the colonies had reached diameters of between 40 and 60 mm, the dry weights of the colonies were determined.

The dry weights of the fungal colonies growing on Carrageenan X-4910 were determined by cutting out a colony and melting the gel in 100 to 200 ml of distilled water at 60°C for 10 min and collecting the colony on a preweighed membrane filter (pore size, 1.2 µm). The filter and filter holder were prewarmed by adding 50 ml of 60°C water to the filtration unit approximately 0.5 min before the filtering, and the same volume of 60°C water was used for washing. The filters were dried at 80°C overnight. Uninoculated plates were used as controls to correct for any precipitations from the medium or resolidification of gel material. There was a slight weight loss by the filters (0.2 to 0.3 mg when treated with 60°C water), for which the data were corrected. Determinations of the dry weights of colonies grown on medium gelled with agar were done like those for the Carrageenan X-4910 cultures, except that boiling water was used instead. The weight losses by the filters were around 1 mg each, for which the data were corrected. Prior to determinations of the dry weights of colonies grown on media solidified with Pluronic F-127, the plates were placed in a refrigerator (4°C) for 15 min, after which the media became liquid. The colony and the liquid medium were poured into 300 ml of ice water and filtered on a precooled (50 ml of ice water) filtration unit, and the dry weight biomass was determined. On media gelled with Carrageenan X-4910 and overlaid with cellophane, the cellophane was lifted from the gel, the colony was gently washed off the cellophane membrane onto a preweighed filter, and the dry weight was determined. The biomass density of a colony was calculated as the dry weight (in milligrams) divided by the area of the colony (in square centimeters). Six replicates were used for determinations of biomass density and K_r . P values were determined by Student's t test.

Measurements of morphology. Measurements of the mycelial morphology of *P. commune* were carried out with an image analysis system (Graphical Image Processing System; Image House Inc., Copenhagen, Denmark) coupled to a Leitz (orthoplan) microscope. The hyphal diameter and the hyphal growth unit length (G_1) were determined. Both parameters were measured on leading surface hyphae in the peripheries of the colonies. The diameter was measured 200 to 300 µm behind the hyphal apex of 32 to 45 leading hyphae at a magnification of $\times 630$. The G_1 values for at least 25 leading hyphae were determined by measuring the total length of a leading hypha and the branches for a distance of approximately 350 µm from the apex of the leading hypha and dividing that by the number of hyphal tips. Further, the visual appearances (e.g., the pigmentation) of the colonies were observed.

RESULTS

Growth on Carrageenan X-4910 and agar. Only small differences between the K_r obtained on medium solidified with agar and that obtained on medium solidified with Carrageenan X-4910 were found, with the exception of that for *A. pullulans* grown on b medium (Table 1). Additionally, the visual appearances (e.g., pigmentation) of the colonies grown on media solidified with agar and Carrageenan X-4910 were similar. No significant difference ($P > 0.05$) between the biomass densities of colonies of *P. farinosus* and *P. commune* grown on b medium gelled with either agar or Carrageenan X-4910 was found (Table 2). Biomass density rather than biomass was used, as it compensates for differences in colony sizes caused by differences in the lag phase and/or in the K_r . In order to determine if the procedure necessary for melting agar gels (boiling for 10

TABLE 1. The K_r s of three fungi grown on media with different gelling substances and on plates covered with cellophane

Medium and substance	K_r ($\mu\text{m h}^{-1}$) ^a		
	<i>P. farinosus</i>	<i>P. commune</i>	<i>A. pullulans</i>
b medium			
Carrageenan	82 ± 8	66 ± 4	74 ± 3
Carrageenan with cellophane	81 ± 4	85 ± 3	65 ± 2
Pluronic F-127	77 ± 2	143 ± 4	86 ± 6
Agar	79 ± 4	67 ± 2	97 ± 6
V8 medium			
Carrageenan	83 ± 3	117 ± 3	106 ± 3
Carrageenan with cellophane	77 ± 5	121 ± 4	97 ± 4
Pluronic F-127	99 ± 3	190 ± 7	138 ± 5
Agar	75 ± 2	119 ± 3	108 ± 4

^a Each result is the mean for six replicate plates ± the standard error.

min) led to a thermic degradation of the biomass, 20 colonies of *P. commune* grown on Carrageenan X-4910 plates were harvested. Ten plates were boiled for 10 min, and 10 plates were harvested by the normal procedure for Carrageenan X-4910 plates (60°C). The biomass dry weight was reduced by 22% when the colonies were boiled. Besides the thermal degradation of biomass, resolidification of agar can be a problem, particularly if filtration is slow. A larger standard error on the determination of biomass density from agar plates was therefore not surprising, and because of the difficulties and uncertainties, no further measurements of the biomass dry weights of colonies grown on agar were made.

Growth on Carrageenan X-4910 overlaid with cellophane. The K_r of *P. farinosus* was not affected by a cellophane membrane, and neither was the K_r of *P. commune* grown on V8 medium (Table 1). But *P. commune* grown on b medium expanded significantly faster when the medium was overlaid with cellophane. *A. pullulans* had a slight but significantly ($P < 0.05$) lower K_r on both media when the plates were covered with cellophane. Cellophane had a more profound effect on the biomass densities of colonies of *P. farinosus* and *P. commune*; biomass density was reduced to less than half on the b medium and to around one-third on V8 medium (Table 2). The response of *A. pullulans* was different, as the cellophane membrane had no effect on the biomass density on the b medium. However, biomass density for *A. pullulans* grown on V8 medium increased by approximately 20% with the cellophane membrane.

TABLE 2. The biomass densities of three fungi grown on media with different gelling substances and on plates covered with cellophane

Medium and substance	Biomass density (mg [dry wt]/cm ²) ^a		
	<i>P. farinosus</i>	<i>P. commune</i>	<i>A. pullulans</i>
b medium			
Carrageenan	2.63 ± 0.24	2.54 ± 0.27	2.47 ± 0.16
Carrageenan with cellophane	1.23 ± 0.18	1.03 ± 0.11	2.50 ± 0.24
Pluronic F-127	0.98 ± 0.20	0.67 ± 0.09	1.41 ± 0.12
Agar	2.55 ± 0.49	2.05 ± 0.50	ND
V8 medium			
Carrageenan	2.18 ± 0.27	1.45 ± 0.16	1.02 ± 0.05
Carrageenan with cellophane	1.54 ± 0.24	0.95 ± 0.22	1.23 ± 0.07
Pluronic F-127	1.55 ± 0.14	0.84 ± 0.06	0.72 ± 0.03
Agar	ND	ND	ND

^a Each result is the mean for six replicate plates ± the standard error. ND, not done.

TABLE 3. The hyphal diameters and G_1 s of *P. commune* grown on b medium solidified with different gelling substances and on plates covered with cellophane

Substance	Hyphal diameter (μm) (mean $\pm$ SE)	G_1 (μm) (mean $\pm$ SE)
Carrageenan	3.5 $\pm$ 0.5	107 $\pm$ 24
Carrageenan with cellophane	4.3 $\pm$ 0.6	134 $\pm$ 32
Pluronic F-127	4.4 $\pm$ 0.8	239 $\pm$ 34 ^a
Agar	3.4 $\pm$ 0.5	107 $\pm$ 26

^a The G_1 was determined by measuring the total hyphal length and the number of branches for a distance of 1,000 μm from the apex. Ten leading hyphae were measured.

Growth on Pluronic F-127 gels. When they were grown on media solidified with Pluronic F-127, all three fungi had K_s significantly higher than those obtained on the agar and Carrageenan X-4910 plates, except for that of *P. farinosus* grown on the b medium (Table 1). The biomass densities of the colonies were reduced considerably compared with those of colonies from the Carrageenan X-4910 plates (Table 2), and the visual appearances of the colonies grown on media solidified with Pluronic F-127 seemed less dense and were lighter pigmented.

Hyphal morphology. The morphology of leading hyphae in the peripheries of colonies of *P. commune* grown on b medium was investigated (Table 3). Measurements of diameters and G_1 s of leading hyphae of colonies grown on agar and Carrageenan X-4910 revealed no difference in the morphology, and this determination supports the conclusion that growth rates of *P. commune* on these gelling compounds are similar. Growth on a cellophane membrane led to a significant increase in both the diameters of the leading hyphae ($P < 0.001$) and the G_1 s ($P < 0.01$). Likewise, the leading hyphae of colonies grown on Pluronic F-127 had larger diameters, and the G_1 s were more than doubled compared with those with agar and Carrageenan X-4910.

DISCUSSION

In the present work, fungal colonies were separated from the agar by being immersed in a boiling water bath for 10 min, even though uninoculated agar plates can be melted by less than 10 min of boiling. However, the presence of a fungal culture with hyphae penetrating the agar slows the melting process, and a boiling period of 10 min, also used by Girvin and Thain (6), was found to be necessary to avoid agar residues. This treatment was shown in the present study to lead to a thermal degradation of the biomass and result in a 22% reduction of biomass compared with the procedure used for isolating biomass from Carrageenan X-4910. This result corresponds to the report of Olson (10), who found that boiling fungal biomass in a microwave oven led to a reduction in dry weight of approximately 30% compared with that found by the no-boiling procedure.

We have shown that fungi have a different response to growth on cellophane membranes. The response also depended on the growth medium. Growth on medium overlaid with cellophane membranes has previously been reported to decrease the radial extension rate of two strains of *Neurospora crassa* (17). Our results suggest that the presence of a cellophane membrane affects the biomass density more than it does the K_r . The observed change in the morphology of the leading hyphae suggests that care should be taken when data obtained with fungi grown on cellophane membrane are extrapolated.

Pluronic F-127 has only seldom been used for the cultivation of fungi. A slight difference in the K_s of *Coprinus congregatus* grown on Emerson's YpSs medium solidified with Pluronic F-127 and the same medium solidified with agar has been observed (2). Our findings suggest that Pluronic F-127 affects growth by generally increasing the rate of expansion but reducing biomass density. The morphology of the leading hyphae differed significantly compared with that of hyphae of colonies grown on agar or Carrageenan X-4910. The increase in the hyphal diameter and a reduced branching frequency leads to a greater volume of cytoplasm per tip. The increase in the hyphal extension rate corresponds well with the theory that the rate of radial extension is closely correlated with the volume of cytoplasm supporting the building blocks for tip growth (16).

Carrageenan X-4910 forms gels that are more transparent than the agar gels, and the costs are the same. One drawback is the gelling temperature, which is higher than that of agar and varies according to the concentrations of certain ions, e.g., K^+ , NH_4^+ , and Ca^{2+} . Carrageenan X-4910 allowed the separation of the biomass from the gel at nondestructive temperatures. This property, together with the fact that growth (except for that of *A. pullulans* on b medium) and hyphal morphology were similar to those obtained on agar, makes Carrageenan X-4910 a useful compound for studies of the relationship between the K_r and the mass growth rates of fungal colonies grown on solid medium.

ACKNOWLEDGMENTS

This work was financed by Danish government funds through the Center for Microbial Ecology.

We thank Kirsten Jensen for providing able technical assistance and Bo Jensen for helpful comments on the manuscript.

REFERENCES

- Allan, E. J., and J. I. Prosser. 1985. Mycelial growth and branching of *Streptomyces coelicolor* A3(2) on solid medium. J. Gen. Microbiol. **129**:2029–2036.
- Choi, H. T., and I. K. Ross. 1988. A low-temperature-liquifying medium for growth and fruiting of *Coprinus congregatus*. Exp. Mycol. **12**:80–83.
- Edelstein, L., Y. Hadar, I. Chet, Y. Henis, and L. A. Segel. 1983. A model for fungal colony growth applied to *Sclerotium rolfsii*. J. Gen. Microbiol. **129**:1873–1881.
- Gardener, S., and G. Jones. 1984. A new solidifying agent for culture media which liquefies on cooling. J. Gen. Microbiol. **130**:731–733.
- Gervais, P., P. Molin, W. Grajek, and M. Bensoussan. 1988. Influence of the water activity of a solid substrate on the growth rate and sporogenesis of filamentous fungi. Biotechnol. Bioeng. **31**:457–463.
- Girvin, D., and J. F. Thain. 1987. Growth of and translocation in mycelium of *Neurospora crassa* on a nutrient deficient medium. Trans. Br. Mycol. Soc. **88**:237–246.
- Kjøller, A., and S. Struwe. 1980. Microfungi of decomposing red alder leaves and their substrate utilisation. Soil Biol. Biochem. **12**:425–431.
- Lines, A. D. 1977. Value of K^+ salt of carrageenan as an agar substitute in routine bacteriological media. Appl. Environ. Microbiol. **34**:637–639.
- Olson, L. W., and L. Lange. 1989. The use of BASF Pluronic polyol F127, low temperature liquifying polyol, for the isolation of microbial antagonists. Opera Bot. **100**:197–199.
- Olson, S. 1994. Uptake of glucose and phosphorus by growing colonies of *Fusarium oxysporum* as quantified by image analysis. Exp. Mycol. **18**:33–47.
- Prosser, J. I., and A. J. Tough. 1991. Growth mechanisms and growth kinetics of filamentous microorganisms. Crit. Rev. Biotechnol. **10**:253–274.
- Reeslev, M., B. B. Jørgensen, and O. B. Jørgensen. 1993. Influence of Zn^{2+} on yeast-mycelium dimorphism and exopolysaccharide production by the fungus *Aureobasidium pullulans* grown in a defined medium in continuous culture. J. Gen. Microbiol. **139**:3065–3070.
- Robson, G. D., S. D. Bell, P. J. Kuhn, and A. P. J. Trinci. 1987. Glucose and penicillin concentrations in agar medium below fungal colonies. J. Gen. Microbiol. **133**:361–367.
- Schnürer, J. 1993. Comparison of methods for estimating the biomass of three food-borne fungi with different growth patterns. Appl. Environ. Microbiol. **59**:552–555.

15. **Trinci, A. P. J.** 1969. A kinetic study of the growth of *Aspergillus nidulans* and other fungi. *J. Gen. Microbiol.* **57**:11–24.
16. **Trinci, A. P. J.** 1971. Influence of the width of the peripheral growth zone on the radial growth rate of fungal colonies on solid media. *J. Gen. Microbiol.* **67**:325–344.
17. **Trinci, A. P. J.** 1973. Growth of wild-type and spreading colonial mutant of *Neurospora crassa* in batch culture and on agar medium. *Arch. Mikrobiol.* **91**:113–126.
18. **Trinci, A. P. J.** 1985. Effect of validamycin A and L-sorbose on the growth and morphology of *Rhizoctonia cerealis* and *Rhizoctonia solani*. *Exp. Mycol.* **9**:20–27.
19. **Watson, N., and D. Apiron.** 1976. Substitute for agar in solid media for common usage in microbiology. *Appl. Environ. Microbiol.* **31**:509–513.
20. **Wiebe, M. G., G. D. Robson, and A. P. J. Trinci.** 1992. Evidence for the independent regulation of hyphal extension and branch initiation in *Fusarium graminearum* A 3/5. *FEMS Microbiol. Lett.* **90**:179–184.