Representative appressorium stage cDNA library of *Magnaporthe grisea**

LU Jian-ping (卢建平), LIU Tong-bao (刘同宝), YU Xiao-yun (于晓云), LIN Fu-cheng (林福呈)[†]

(School of Life Sciences, Zhejiang University, Hangzhou 310029, China)

[†]E-mail: fuchenglin@zju.edu.cn

Received Apr. 29, 2004; revision accepted June 15, 2004

Abstract: A mature appressorium cDNA library of rice blast fungus, *Magnaporthe grisea*, was constructed in a λ TriplEx2 vector by SMARTTM cDNA library containing 2.37×10^6 independent clones about 100% of which harbor foreign cDNA inserts with average size of 660 bp. Of 9 randomly selected clones, 2 expressed sequence tags (ESTs) sequences did not have homologous EST sequences of *M. grisea* in GenBank. The appressorium cDNA library is suitable for gene expression analysis and function analysis of the late stages of appressorium formation and the early stages of penetration of *M. grisea*.

Key words: *Magnaporthe grisea*, Appressorium, cDNA library

doi:10.1631/jzus.2005.B0132

Document code: A

CLC number: Q949.32

INTRODUCTION

The rice blast fungus *Magnaporthe grisea* causes one of the most destructive diseases of rice. Genetic studies of this important pathogen during the past decade made it excellent for investigating fungal-plant interactions. Appressorium is an infection structure of *M. grisea*. Many genes expressed in appressorium, such as *PMK1* (Xu and Hamer, 1996), *CYP1* (Viaud *et al.*, 2002), *MAGB* and *MAC1* (Choi and Dean, 1997; Liu and Dean, 1997), *GAS1* and *GAS2* (Xue *et al.*, 2002), and *MPG1* (Lau and Hamer, 1996; Talbot *et al.*, 1993; 1996), are required for appressorium formation and pathogenesis by the fungus. Many approaches, such as expressed sequence tags (ESTs) sequencing (Kamakura *et al.*, 1999; Kim *et al.*, 2001; Raayaree *et al.*, 2001), SAGE (Serial analysis of gene expression) (Irie *et al.*, 2003), microarray (Takano *et al.*, 2003), and mutants (Balhadere *et al.*, 1999; Shi *et al.*, 1995; Sweigard *et al.*,

1998), have been used to identify genes already expressed in appressorium. By now, more than 10000 ESTs have been sequenced from several cDNA libraries representing different stages of fungal growth, differentiation and plant infection of the pathogen, including appressoria, conidia, and mycelia from complete and nitrogen starvation cultures (Xu and Xue, 2002). However, appressorium cDNA libraries in existence were only constructed from conidia germinated for less than 8 h on an inductive surface (Kamakura *et al.*, 1999; Takano *et al.*, 2003), causing a significant problem for completely evaluating gene expression during the late stages of appressorium formation. We report here the result of producing an appressorium stage cDNA library of *M. grisea* from mature appressoria formed for 23.5–24.5 h.

MATERIALS AND METHODS

M. grisea isolate, appressorium induction and RNA isolation

The appressorium of *M. grisea* GUY11 was used to isolate total RNA for construction of cDNA library.

* Project supported by the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry, and in part by the National Natural Science Foundation of China (No. 30270049)

For appressorium formation, droplets (20 μ l) of conidium suspension (1×10^6 – 1.5×10^6 conidia/ml in distilled water) were inoculated on an inductive surface of projection membrane (Gaoke, China) for 23.5–24.5 h at 25 °C. And then appressoria were harvested and from which total RNA was isolated with Trizol (Molecular Research Center, USA) following the manufacturer's procedure.

Appressorium cDNA synthesis and library construction

The ss cDNA, ds cDNA and library construction were carried out according to Smart cDNA Library Construction Kit User Manual (CLI, 2001). Three μ l (~5.23 μ g) total RNA was used to reverse transcribe to single-strand cDNA with SMART IVTM Oligonucleotide, CDS III/3' PCR Primer and PowerScriptTM Reverse Transcriptase at 42 °C for 1 h. Two μ l of first-strand cDNA was used to carry out ds cDNA synthesis by 23 cycles of PCR (95 °C for 15 s, 68 °C for 6 min) with 5' PCR primer, CDS III/3' PCR primer and AdvantageTM 2 polymerase mix. Half of the ds cDNA were digested with proteinase K and purified with phenol:chloroform: isoamylalcohol. Then all purified ds cDNA was digested with Sfi I enzyme. The reaction product was fractionated by CHROMA SPIN-400 column and the first three fractions containing ds cDNA were collected. The cDNA of all three fractions containing ds cDNA was concentrated by ethanol and resuspended in 7 μ l deionized H₂O. At last, these Sfi I-digested cDNAs were ligated to Sfi I-digested, dephosphorylated λ TriplEx2 vector at 16 °C overnight. The ligated cDNA/ λ TriplEx2 vector was packed using MaxPlaxTM Lambda Packaging Extracts (Epicentre, USA).

Analysis of the appressorium cDNA library

The unamplified cDNA library was titered and the percentage of recombinant clones was determined by blue/white screening according to the instructions in manual of SMARTTM cDNA library construction kit (Clontech, USA).

Twelve randomly selected plaques were used for the conversion of λ TriplEx2 clone to pTriplEx2 plasmid according to the instructions of manual of SMARTTM cDNA library construction kit. The plasmids were purified by CTAB-method DNA Mini Preparation and digested by Sfi I enzyme. The inserts

were analyzed by separation on a 1% agarose/EtBr gel. And the expressed sequence tags (EST) of nine randomly selected clones were sequenced with T7 primer on an ABI 377 DNA sequencer. The sequence data were analyzed using VecScreen program (NCBI) for vector masking, and then the adaptor sequences were removed from these cDNA sequences. These EST sequences were processed using software BioEdit (Hall, 1999) for contig assembly. Process sequences were subjected to similarity searches against GenBank database using Blast 2.2.8 (Altschul et al., 1997) and against Phytopathogenic Fungi and Oomycete EST database (Version 1.4) in COGEME (<http://cogeme.ex.ac.uk/index.html>) and *Magnaporthe grisea* database (<http://www.broad.mit.edu/annotation/fungi/magnaporthe/>).

Appressorium cDNA library amplification and titer of the amplified library

The manual of SMARTTM cDNA library construction kit suggested amplification of the unamplified cDNA library and then titering of the amplified library.

RESULTS

Appressorium induction of *M. grisea*

More than 96% conidia inoculated on the hydrophobic surface of a projection membrane for 23.5–24.5 h at 25 °C germinated and formed appressoria (Fig.1).

Fig.1 Appressoria of *M. grisea* GUY11 on a projection membrane inoculated for 23.5–24.5 h at 25 °C

Total RNA isolation of appressorium

The ratio of OD260/OD280 of total RNA isolated from appressoria incubated for 23.5–24.5 h was 1.63. From the gel, total RNA appeared as two bright bands (28s rRNA and 18s rRNA) and showed intact (Fig.2).

Fig.2 Appressorium total RNA (5 µl) on a 1.2% denaturing formaldehyde/agarose gel

cDNA synthesis and library construction quality analysis

The sizes of the ds cDNA produced from the 24 h appressorium mRNA ranged from 200 bp to 6 kb, mainly between 500 bp and 5 kb on a 1.1% agarose gel (Fig.3). After fractionation, the first three fraction containing cDNA (8, 9 and 10 fractions) were collected and used for ligation to vector.

Fig.3 Ten µl ds cDNA product synthesized from appressorium RNA electrophoresed on a 1.1% agarose/EtBr gel. Lane M, 1-kb DNA size marker (Fermentas, Lithuania) (0.2 µg loaded)

The cDNA was ligated unidirectionally into λTriplEx2 vector by three ligation reactions. From the three ligations combined, a primary library of 2.37×10^6 independent clones was obtained. The result of blue/white screening of the primary cDNA library in *E. coli* XL1-Blue showed that the ratio of white (recombinant) to blue (nonrecombinant) in two LB/MgSO₄ plates containing IPTG and X-gal were 684/0 and 931/0 respectively. So, the recombination efficiency of cDNA ligation reached 100%. After amplification, the amplified cDNA library had a titer of 2.86×10^9 pfu/ml.

Twelve randomly selected lambda plaques of the primary library were converted to plasmids by in vivo

conversion and were analyzed for insert size. The plasmids were purified with CTAB method and digested by Sfi I. All 12 plasmids showed that the insert sizes ranged from 300 bp to 900 bp, average of 660 bp (Fig.4).

Fig.4 Sfi I-digested plasmids of 12 clones on a 1% agarose/EtBr gel

Lane M, 1-kb DNA marker size (Fermentas, Lithuania). 1~12: clones 1~12

Nine randomly selected cDNA clones were partially sequenced using T7 primers. Analysis of the 9 cDNA sequences using BioEdit software (Hall, 1999) generated 8 nonredundant EST sequences. All 8 unique EST sequences were subjected to similarity searches against GenBank (Blastn 2.2.8 and Blastx 2.2.8) (Altschul *et al.*, 1997), COGEME Db and *Magnaporthe grisea* Database. Six of them had homologous sequences of *M. grisea* in dbEST and the other 2 clones' EST sequences were found to be in-homologous sequences (Table 1) until submission. Clone C02 was homologous to putative protein of cDNA of *Neurospora crassa* (GenBank Accn: AW723279, Expect value= $3e-53$). As for clone C04, no homologous putative protein sequence or EST sequence hits with *E*-value less than $e-3$ were found in NCBI blast. These sequences have been deposited in the dbEST of GenBank database (GenBank Accn: CK725218~CK725225).

Table 1 Homologous sequences from *M. grisea* of 8 EST sequences by Blastn in GenBank

Clone noes	Putative product/function
C01	Glutathione peroxidase (BU641649, $e-140$)*
C02	Not found
C04	Not found
C05	Cyclophilin (peptidyl-prolyl cis-trans isomerase) (BI808723, 0.0)
C06	ATP synthase subunit 9, lipid-binding protein, mitochondrial (AA415145, 0.0)
C09	Unknown (CD031047, 0.0)
C11	Unknown (CD036090, $e-119$)
C12	Unknown (BM872180, 0.0)

* GenBank accession numbers and *E*-values for homologues are in parentheses

DISCUSSION

Although there are many *M. grisea* cDNA libraries in the world, mainly in Ebbole's lab at Texas A&M University and Ralph Dean's lab at North Carolina State Biotechnology Center (Ebbole *et al.*, 2002; Xu and Xue, 2002), and there are five reported appressorium cDNA libraries built from germinated conidia on appressorium-inductive surface (http://www.tigr.org/tdb/tgi/mggi/searching/xpress_search.html), no cDNA library was built strictly from mature appressorium. The appressorium cDNA lambda expression library reported here was constructed from mature appressoria inoculated on an inductive surface for 23.5~24.5 h at 25 °C, and had 2.37×10^6 pfu independent clones with 100% recombinants enough to screen genes expressed in appressorium. In addition, all cDNA sequences inserted into MCS of λ TriplEx2, the vector of this library, can be expressed for screening genes with protein antigens. And the vector λ TriplEx2 is easily converted from phage to a plasmid vector via Cre-lox-mediated subcloning.

Two of 8 non-redundant EST sequences have no homologues with *E*-value less than e^{-3} in *M. grisea* ESTs available in GenBank until submission. This cDNA library probably contains many genes missed by previous *M. grisea* cDNA libraries. Clone C05 encoded a reported peptidyl prolyl cis-trans isomerase (cyclophilin). Cyclophilin gene *CYP1* is a virulence factor in *M. grisea* (Viaud *et al.*, 2002). *CYP1* mutants show reduced virulence and are impaired in associated functions, such as penetration peg formation and inducement of appressorium turgor. From this cDNA library made from mature appressoria, a 3.0 kb long cDNA sequence of *MGT1* gene has been successfully cloned (¹Wang, unpublished). The appressorium cDNA library ESTs can be used for identification of genes expressed in appressorium. For example, promoters of 12 genes were isolated from the rice blast fungus based on the sequences of randomly ESTs (appressorium formation stage cDNA library of *Magnaporthe* available from GenBank) and then these promoters can be used to identify the genes expressed during infection stages (Banno *et al.*, 2003).

So, this 24 h appressorium cDNA library is a representative cDNA library that reveals the gene expression state in appressorium and is suitable for gene expression and function analysis of *M. grisea* appressorium.

References

- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., Lipman, D.J., 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.*, **25**:3389-3402.
- Balhadere, P.V., Foster, A.J., Talbot, N.J., 1999. Identification of pathogenicity mutants of the rice blast fungus *Magnaporthe grisea* using insertional mutagenesis. *Mol. Plant-Microbe Interact.*, **12**:129-142.
- Banno, S., Kimura, M., Tokai, T., Kasahara, S., Higa-Nishiyama, A., Takahashi-Ando, N., Hamamoto, H., Fujimura, M., Staskawicz, B.J., Yamaguchi, I., 2003. Cloning and characterization of genes specifically expressed during infection stages in the rice blast fungus. *FEMS Microbiol Lett*, **222**:221-227.
- Choi, W., Dean, R.A., 1997. The adenylate cyclase gene *MAC1* of *Magnaporthe grisea* controls appressorium formation and other aspects of growth and development. *Plant Cell*, **9**:1973-1983.
- Ebbole, D.L., Yuan, J., Li, D., Thomas, T.L., Dean, R.A., 2002. Gene Discovery in the Rice Blast Fungus, *Magnaporthe Grisea*: Analysis of ESTs. In: Abstracts for Plant. Animal & Microbe Genomes X, San Diego, CA, p.18.
- Hall, T.A., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids. Symp. Ser.*, **41**:95-98.
- Irie, T., Matsumura, H., Terauchi, R., Saitoh, H., 2003. Serial Analysis of Gene Expression (SAGE) of *Magnaporthe grisea*: genes involved in appressorium formation. *Mol. Genet. Genomics*, **270**:181-189.
- Kamakura, T., Xiao, J., Choi, W., Kochi, T., Yamaguchi, S., Teraoka, T., Yamaguchi, I., 1999. cDNA subtractive cloning of genes expressed during early stage of appressorium formation by *Magnaporthe grisea*. *Biosci. Biotechnol. Biochem.*, **63**:1407-413.
- Kim, S., Ahn, I.P., Lee, Y.H., 2001. Analysis of genes expressed during rice-*Magnaporthe grisea* interactions. *Mol. Plant-Microbe Interact.*, **14**:1340-346.
- Lau, G.W., Hamer J.E., 1996. Regulatory genes controlling *MPG1* expression and pathogenicity in the rice blast fungus *Magnaporthe grisea*. *Plant Cell*, **8**:771-781.
- Liu, S., Dean, R.A., 1997. G protein α -subunit genes control growth, development and pathogenicity of *Magnaporthe grisea*. *Mol Plant-Microbe Interact*, **10**:1075-1086.
- Rauyaree, R., Choi, W., Fang, E., Blackmon, B., Dean, R.A., 2001. Genes expressed during early stages of rice infection with the rice blast fungus *Magnaporthe grisea*. *Mol. Plant Pathol.*, **2**:347-354.
- Shi, Z., Christian, D., Leung, H., 1995. Enhanced transforma-

¹Wang, J.Y., Liu, X.H., Lin, F.C., 2004. Cloning and characterization of *MGT1* gene in rice blast pathogen *Magnaporthe grisea*. *Fungal Genetics and Biology*.

- tion in *Magnaporthe grisea* by restriction enzyme mediated integration of plasmid DNA. *Phytopathology*, **85**:329-333.
- CLI (CLONTECH Laboratories, Inc.), 2001. Smart cDNA Library Construction Kit User Manual. USA, p.1-51.
- Sweigard, J.A., Carroll, A.M., Farrall, L., Chumley, F.G., Valent, B., 1998. *Magnaporthe grisea* pathogenicity genes obtained through insertional mutagenesis. *Mol. Plant-Microbe Interact.*, **11**:404-412.
- Takano, Y., Choi, W., Mitchell, T.K., Okuno, T., Dean, R.A., 2003. Large scale parallel analysis of gene expression during infection-related morphogenesis of *Magnaporthe grisea*. *Mol. Plant Pathol.*, **4**:337-347.
- Talbot, N.J., Ebbole, D.J., Hamer, J.E., 1993. Identification and characterization of *MPG1*, a gene involved in pathogenicity from the rice blast fungus *Magnaporthe grisea*. *Plant Cell*, **5**:1575-1590.
- Talbot, N.J., Kershaw, M.J., Wakley, G.E., De Vries, O.M.H., Wessels, J.G.H., Hamer, J.E., 1996. *MPG1* encodes a fungal hydrophobin involved in surface interactions during infection-related development of *Magnaporthe grisea*. *Plant Cell*, **8**:985-999.
- Viaud, M.C., Balhadere, P.V., Talbot, N.J., 2002. A *Magnaporthe grisea* cyclophilin acts as a virulence determinant during plant infection. *Plant Cell*, **14**:917-930.
- Xu, J.R., Hamer, J.E., 1996. MAP kinase and cAMP signaling regulate infection structure formation and pathogenic growth in the rice blast fungus *Magnaporthe grisea*. *Genes Dev.*, **10**:2696-2706.
- Xu, J.R., Xue, C.Y., 2002. Time for a blast: genomics of *Magnaporthe grisea*. *Mol. Plant Pathol.*, **3**:173-176.
- Xue, C.Y., Park, G., Choi, W., Zheng, L., Dean, R.A., Xu, J.R., 2002. Two novel fungal virulence genes specifically expressed in appressoria of the rice blast fungus. *Plant Cell*, **14**:2107-2119.

Welcome contributions from all over the world

<http://www.zju.edu.cn/jzus>

- ◆ The Journal aims to present the latest development and achievement in scientific research in China and overseas to the world's scientific community;
- ◆ JZUS is edited by an international board of distinguished foreign and Chinese scientists. And an internationalized standard peer review system is an essential tool for this Journal's development;
- ◆ JZUS has been accepted by CA, Ei Compendex, SA, AJ, ZM, CABI, BIOSIS (ZR), IM/MEDLINE, CSA (ASF/CE/CIS/Corr/EC/EM/ESPM/MD/MTE/O/SSS*/WR) for abstracting and indexing respectively, since started in 2000;
- ◆ JZUS will feature **Sciences in Engineering** subjects in Vol. A, 12 issues/year, and **Life Sciences & Biotechnology** subjects in Vol. B, 12 issues/year;
- ◆ JZUS has launched this new column "**Science Letters**" and warmly welcome scientists all over the world to publish their latest research notes in less than 3-4 pages. And assure them these Letters to be published in about 30 days;
- ◆ JZUS has linked its website (<http://www.zju.edu.cn/jzus>) to **CrossRef**: <http://www.crossref.org> (doi:10.1631/jzus.2005.xxxx); **MEDLINE**: <http://www.ncbi.nlm.nih.gov/PubMed>; **HighWire**: <http://highwire.stanford.edu/top/journals.dtl>; **Princeton University Library**: <http://libweb5.princeton.edu/ejournals/>.