Journal of Zhejiang University SCIENCE
 ISSN 1009-3095
 http://www.zju.edu.cn/jzus
 E-mail: jzus@zju.edu.cn

Science Letters:

Differentiation of xanthomonads causing the bacterial leaf spot of poinsettia in China from the pathotype strain of *Xanthomonas axonopodis* pv. *poinsettiicola**

LI Bin (李斌)¹, XIE Guan-lin (谢关林)^{†1}, SWINGS J.²

(¹Institute of Biotechnology, Zhejiang University, Hangzhou 310029, China)

(²Laboratorium voor Microbiologie, Universiteit Gent, K. Ledeganckstraat 35, 9000, Gent, Belgium)

[†]E-mail: glxie@zju.edu.cn

Received Jan. 18, 2005; revision accepted Apr. 1, 2005

Abstract: In October 2003, a new bacterial disease with symptoms similar to those caused by *Xanthomonas axonopodis* pv. *poinsettiicola* was observed on poinsettia leaves at a flower nursery in Zhejiang Province of China. Three *Xanthomonas* strains were isolated from infected plants and classified as *X. axonopodis*. They were differentiated from the pathotype strain LMG849 of *X. axonopodis* pv. *poinsettiicola* causing bacterial leaf spot of poinsettia by comparison of pathogenicity, substrate utilization and BOX-PCR genomic fingerprints.

Key words: Bacterial leaf spot of poinsettia, BOX-PCR, Hypersensitive reaction, Pathogenicity, *X. axonopodis* pv. *poinsettiicola*
doi:10.1631/jzus.2005.B0451 **Document code:** A **CLC number:** S432.4

INTRODUCTION

In recent years poinsettia cultivation has blossomed from a small garden business into a multimillion-dollar industry in China. Some fungal diseases, such as *Rhizoctonia* stem and root rot, *Pythium* root rot, *Fusarium* wilt, and *Botrytis* blight have been reported to cause considerable damage during propagation (Sun *et al.*, 2003). A new bacterial disease characterized by leaf spot appearance was first observed on poinsettia in October 2003, in the Xiaoshan district of Zhejiang Province, China, one of the major poinsettia-production areas. Three bacterial strains have been isolated and classified as *X. axonopodis*. However, they have not been characterized yet on the pathovar level.

Bacterial leaf spot of poinsettia with symptoms similar to those caused by *X. axonopodis* pv. *poinsettiicola* (formerly *X. campestris* pv. *poinsettiicola*)

was first reported by Patel *et al.* (1951) in India. DNA-DNA hybridization studies indicated that *X. axonopodis* was the largest and most heterogeneous genomic group (Vauterin *et al.*, 1995). Phenotypic characterization including hypersensitive reaction, pathogenicity, and substrate utilization is essential for the characterization of phytopathogens, but insufficient for differentiating strains at or below the pathovar level. The rep-PCR technique has proved to be a rapid, simple and reproducible method to classify and differentiate *Xanthomonas* strains (Louws *et al.*, 1994). The objective of the present study was to elucidate the relationship between the causal agent and the pathotype strain of *X. axonopodis* pv. *poinsettiicola*.

MATERIALS AND METHODS

Three strains, R22578, R22579 and R22580, isolated from symptomatic leaves of poinsettia in China and the pathotype strain LMG849 of *X. axonopodis* pv. *poinsettiicola* originally from India

* Project supported by the National Natural Science Foundation of China (No.30370951) and China-Belgium Flemish Government

were used in this study. Cultures were stored in a freezer at -80°C ; and were checked for purity and grown routinely on Tryptic soy agar (Mew and Misra, 1994) at 28°C . All strains were deposited in the BCCMTM, Belgium and Institute of Biotechnology, Zhejiang University, China. Hypersensitive reactions of the 4 strains on tobacco as described by Schaad *et al.* (2001) were observed and pathogenicity test was done on pot-cultivated poinsettia seedling (cv. Qian-xi, Bi-de and Fu-xing) in greenhouse by pricking the leaves with a needle and placing a $10\ \mu\text{l}$ droplet of bacterial suspension (approximately 10^8 cfu/ml) over the wounded site.

BIOLOG test was performed following manufacturer's instructions (BIOLOG, Inc., Hayward, CA) to compare the substrate utilization of the three strains with those of strain LMG849 (Xie *et al.*, 2003). For genotyping, total DNAs of the 4 strains were extracted by the CTAB method (Ausubel *et al.*, 1992) and repetitive PCR genomic fingerprints were obtained by using the BOX A1R primer described by Louws *et al.* (1994). All PCRs were performed in a total volume of $25\ \mu\text{l}$ using 30–50 ng DNA, 500 $\mu\text{mol/L}$ each dNTP, 2 $\mu\text{mol/L}$ BOX A1R primer and 2.0 U *Taq* polymerase. Cycling conditions were: one cycle of 95°C for 7 min, and 35 cycles of 94°C for 1 min, 53°C for 1 min and 65°C for 8 min with a single final extension cycle at 65°C for 15 min and a final incubation at 4°C . PCR products were examined by electrophoresis in 1% agarose gel and DNA was visualized with a gel imaging system.

RESULTS AND DISCUSSION

Infiltration of tobacco leaves with cell suspensions of the 3 strains from China resulted in typical hypersensitivity reactions within 24 h and induced similar symptoms with field observations on pot-cultivated poinsettia seedling 2 weeks after inoculation. On the under leaf surface, new spots were dull grey to brown and slightly water-soaked. The spots later became chocolate-brown and visible on both surfaces. Lesions varied from circular to angular in shape and coalesced to form large, irregular, necrotic areas with a yellow halo. The lesions were often along the veins and midribs, although not confined to these areas. Bacterial exudates often formed on le-

sions, especially on the lower leaf surface. The ooze dried to form a shiny scale-like crest. Three cultivars (Qian-xi, Bi-de and Fu-xing) of poinsettia were all susceptible to the causal agent. A delayed hypersensitivity reaction on tobacco and weak pathogenicity symptoms on leaves of 3 poinsettia cultivars were observed for the pathotype strain LMG849.

Comparisons of metabolic activity on the carbon substrates, by which strains of *X. axonopodis* can be distinguished from those of other *Xanthomonas* species, revealed that the 3 strains had the similar utilization reactions as the type strain LMG849 (Table 1). Strains R22579 and R22580 showed indistinguishable reaction patterns were different from those of strain R22578 on the utilization of succinamic acid and *p*-hydroxyphenylacetic acid. The major difference between the 3 strains and the pathotype strain LMG849

Table 1 Partial substrate utilization profiles of the poinsettia bacterial strains R22578, R22579, R22580 and the pathotype strain LMG849

BIOLOG carbon sources	R22578	R22579	R22580	LMG849
Dextrin	+	+	+	+
Cellobiose	+	+	+	+
Gentiobiose	+	+	+	+
Maltose	+	+	+	+
D-psicose	+	+	+	+
D-trehalose	+	+	+	+
Monomethylsuccinate	+	+	+	+
Succinic acid	+	+	+	+
Bromosuccinic acid	+	+	+	+
Succinamic acid	–	+	+	+
D-alanine	+	+	+	+
L-alanine	+	+	+	+
L-alanylglycine	+	+	+	+
L-glutamic acid	+	+	+	+
Glycyl-L-glutamic acid	+	+	+	+
β -methyl-D-glucoside	±	±	±	–
L-rhamnose	–	–	–	–
Formic acid	+	+	+	–
D-galactonic acid lactone	–	–	–	–
D-galacturonic acid	–	–	–	–
D-glucuronic acid	–	–	–	–
<i>p</i> -hydroxyphenylacetic acid	–	±	±	–
α -ketovaleric acid	–	–	–	–
Quinic acid	–	–	–	–
Glucuronamide	–	–	–	–
L-phenyl-alanine	–	–	–	–
Thymidine	–	–	–	–

BIOLOG data are presented only for the reactions, in which the strains of *X. axonopodis* can be distinguished from other *Xanthomonas* species. +: Positive; ±: Weakly positive; –: Negative

was the utilization of formic acid and β -methyl-D-glucoside. The representative fingerprint patterns of the 4 strains were obtained using the BOX A1R primer (Fig.1), in which the bands, ranging in size from approximately 200 bp to over 2500 bp, were clearly differentiated by agarose gel electrophoresis. The three strains isolated from China showed a similar profile, which showed that a single species might be responsible for bacterial leaf spot of poinsettia in China. The genetic profiles of the 3 strains were clearly different from that of LMG849, which is consistent with the result of phenotypic characterization. The present study revealed the causal agent is different from that of the pathotype strain and confirmed that repetitive sequence-based polymerase chain reaction could be used as rapid, highly discriminatory screening techniques to determine the genomic diversity of *X. axonopodis*.

Fig.1 Genomic fingerprinting of bacterial strains amplified using primer BOX

Lane 1: R22578; 2: R22579; 3: R22580; 4: LMG849; M: 100 bp DNA molecular marker (Bioasia)

References

- Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A., Struhl, K., 1992. Current Protocols in Molecular Biology. Greene Publishing Associates and Wiley Interscience, New York.
- Louws, F.J., Fulbright, D.W., Stephens, C.T., de Bruijn, F.J., 1994. Specific genomic fingerprints of phytopathogenic *Xanthomonas* and *Pseudomonas* pathovars and strains generated with repetitive sequences and PCR. *Applied and Environmental Microbiology*, **60**:2286-2295.
- Mew, T.W., Misra, J.K., 1994. A Manual of Rice Seed Health Testing. International Rice Research Institute, Manila, Philippines, p.109-110.
- Patel, M.K., Bhatt, V.V., Kulkarni, Y.S., 1951. Three new bacterial diseases of plants from Bombay. *Indian Phytopathology*, **4**:144-151.
- Schaad, N.W., Jones, J.B., Chun, W., 2001. Laboratory Guide for Identification of Plant Pathogenic Bacteria. American Phytopathological Society, St. Paul, Minn.
- Sun, J.X., Wang, J.Q., Wang, X., Liu, X.Q., Lu, X.L., 2003. Study on the roots rot diseases of *Euphorbia Pulcherrima* Willd and their control. *Journal of Laiyang Agricultural College*, **20**(3):168-171 (in Chinese).
- Vauterin, L., Hoste, B., Kersters, K., Swings, J., 1995. Re-classification of *Xanthomonas*. *International Journal of Systematic Bacteriology*, **45**:472-489.
- Xie, G.L., Soad, A., Swings, J., Mew, T.W., 2003. Diversity of gram negative bacteria antagonistic against major pathogens of rice from rice seed in the tropic environment. *Journal of Zhejiang University Science*, **4**(4):463-468.

Welcome visiting our journal website: <http://www.zju.edu.cn/jzus>

Welcome contributions & subscription from all over the world

The editor would welcome your view or comments on any item in the journal, or related matters

Please write to: Helen Zhang, Managing Editor of JZUS

E-mail: jzus@zju.edu.cn Tel/Fax: 86-571-87952276/87952331