Verapamil pharmacokinetics

In the April issue of the Journal Schwartz *et al.* (1985) provided interesting information concerning the apparent accumulation of verapamil during chronic therapy. We suggest, however, that their principal conclusion that the observed pharmacokinetic changes cannot be attributed to changes in liver blood flow is based on inappropriate interpretation of their data.

The pharmacokinetic data presented in their paper are very similar to our findings in a group of normotensive volunteers (Meredith *et al.*, 1985). We have shown that there are significant differences in elimination half-life and AUC when comparing continued administration with first dosing and that this occurs in the absence of a significant change in the AUC verapamil/AUC norverapamil ratio. Schwartz *et al.* (1985) have additionally demonstrated that there were no significant changes in the urinary excretion pattern of four other *N* and *O*-dealkylated metabolites of verapamil and we would certainly concur with their interpretation that the prolongation of verapamil half-life and decrease in apparent clearance is not due to impairment of the hepatic biotransformation of verapamil.

However, we differ fundamentally in our interpretation of the liver blood flow findings. We have demonstrated that the acute administration of verapamil is associated with a significant ($P < 0.005$) increase in apparent liver blood flow (Meredith *et al.*, 1985). This was measured 1 h after drug administration in order to coincide with peak plasma drug levels. Apparent liver

blood flow increased to a mean value ($\pm$ s.d.) of $1401 (\pm 363)$ ml min⁻¹ compared to $954 (\pm 219)$ ml min⁻¹ for placebo but with continued administration there was then a significant ($P < 0.01$) reduction towards placebo values at $1167 (\pm 375)$ ml min⁻¹. Because the increase in apparent liver blood flow is transient and returns to basal levels during continued administration, the failure of Schwartz *et al.* (1985) to detect the acute increase in ICG clearance was almost certainly due to timing their baseline administration of ICG *prior* to drug dosing.

Thus it is a transient but profound increase in liver blood flow which is associated with a shorter half-life and more rapid clearance on first dosing with the drug. When liver blood flow returns to baseline values during continued administration the 'true' verapamil kinetics are restored. Our hypothesis, therefore, is that the accumulation of verapamil during chronic therapy is an *apparent* accumulation and that the change in kinetic parameters is more appropriately described as a shortening of elimination half-life and an increase in clearance following acute dosing.

P. A. MEREDITH, H. L. ELLIOTT &
J. L. REID

*University Department of Materia Medica,
Stobhill General Hospital, Glasgow G21 3UW*

Received 12 August 1985,
accepted 10 December 1985

References

- Meredith, P. A., Elliott, H. L., Pasanisi, F., Kelman, A. W., Sumner, D. J. & Reid, J. L. (1985). Verapamil pharmacokinetics and apparent hepatic and renal blood flow. *Br. J. clin. Pharmac.*, **20**, 101-106.
- Meredith, P. A., Pasanisi, F., Elliott, H. L. & Reid, J. L. (1985). The effect of nisoldipine on apparent liver blood flow and effective renal plasma flow. *Br. J. clin. Pharmac.*, **20**, 235-237.
- Schwartz, J. B., Abernethy, D. R., Taylor, A. A. & Mitchell, J. R. (1985). An investigation of the accumulation of verapamil during regular dosing in patients. *Br. J. clin. Pharmac.*, **19**, 512-516.