THE USE OF SHORT- AND LONG-ACTING HYPNOTICS IN CLINICAL MEDICINE

A.N. NICHOLSON

Royal Air Force Institute of Aviation Medicine
Farnborough, Hampshire, England

1 Activity of short- and long-acting benzodiazepines is reviewed with reference to pharmacokinetics and residual sequelae, and to efficacy and adverse effects.

2 Some benzodiazepines may not lead to obvious effects on performance, such as nordiazepam and clobazam, and the persistence of residual sequelae may not relate obviously to elimination half-lives (as with diazepam and possibly flunitrazepam). However, benzodiazepines with mean half-lives less than 8 h may have residual sequelae, whereas hypnotics with mean half-lives greater than 16 h are likely to lead to impaired performance and/or anxiolytic effects the next day.

3 Potassium chlorazepate 15 mg, with its long-acting metabolite nordiazepam, would seem to be the drug of choice for insomnia secondary to anxiety. For the insomniac without significant psychopathology, temazepam 10–20 mg, triazolam 0.125–0.25 mg and for occasional use, diazepam 5–10 mg, provide the initial approach. Flurazepam hydrochloride 15–30 mg, nitrazepam 5–10 mg and flunitrazepam 1 mg and above, have persistent residual effects and should be reserved for refractory patients, and for those in whom some impairment of performance the next day would be acceptable.

4 There is little or no evidence to suggest that the proper use of the short-acting hypnotics, triazolam and temazepam, leads to a worsening of sleep on withdrawal. However, some benzodiazepines may lead to disturbances of sleep and/or rebound insomnia, and nitrazepam and flunitrazepam may be implicated.

Introduction

Drugs used in the management of insomnia (Figure 1) present a continuing process of development, although, without doubt, the most important event has been the introduction of the benzodiazepines. Nitrazepam and flurazepam hydrochloride have proved to be relatively safe if an overdose is taken, and the activity of flurazepam is without serious adverse effects on sleep. However, nitrazepam has an elimination half-life of 18–36 h (Breimer, Bracht & DeBoer, 1977), and the activity of flurazepam is probably related to a relatively long-acting metabolite (Kaplan, deSilva, Jack, Alexander, Strojny, Weinfeld, Puglist & Weissman, 1973). Persistent plasma levels are associated with residual effects, and it is now widely accepted that overnight ingestion of nitrazepam 10 mg and flurazepam 30 mg impairs performance well into the next working day (Bond & Lader, 1972, 1973; Borland & Nicholson, 1975a). The lower dose of flurazepam 15 mg is likely to have less persistent effects on performance, but its efficacy as an hypnotic at this dose has not been established.

A partial answer to the problem of impaired performance the next day is the use of diazepam. Ingestion of diazepam 5–10 mg overnight is uncomplicated

by residual effects (Clarke & Nicholson, 1978), even though it has a long elimination half-life of 14–90 h (Mandelli, Tognoni & Garattini, 1978). The phenomenon is probably related to the equilibration phase as initial recovery from impaired performance is rapid. However, daily ingestion will lead to accumulation of both the parent drug and its long-acting metabolite, nordiazepam, and so is unlikely to be without residual effects, and should be used only occasionally if sequelae are to be avoided.

Hypnotics in which individual elimination half-lives do not exceed 24 h are much less likely to lead to impaired performance. The most important compounds in this group are, nevertheless, structurally similar to diazepam, although they are without active, long-acting metabolites. The hydroxylated metabolites of diazepam, temazepam (3-hydroxydiazepam) and oxazepam (3-hydroxy-N-desmethyldiazepam) have elimination half-lives of 4–10 h and 6–24 h respectively (Fucella, Bolcioni, Tamassia, Ferrario & Tognoni, 1977; Sjoqvist & Sundwall, 1977) and their chlorinated derivatives, lorazepam and lorazepam, have elimination half-lives of 9–22 h (Greenblatt, Joyce, Comar, Knowles, Schader,

Figure 1 Structural formulae for diazepam, nitrazepam, flurazepam, flunitrazepam, fosazepam and triazolam.

Kyriakopoulos, MacLaughlin & Ruelius, 1977) and 11–15 h (Schering AG, internal report) respectively, whereas triazolam, a triazolo-1,4-benzodiazepine with an orthochlorophenyl group, has an elimination half-life of 3–5 h (Eberts, Ko & Thomas, 1979). Temazepam and triazolam have particularly short elimination half-lives, and, at present, it is around these compounds that interest concerning the place of short-acting hypnotics in therapeutics is centered.

Duration of activity and the possibility of residual effects play an important part in deciding the place of various hypnotics in the management of sleep. These properties, together with effectiveness and adverse effects on sleep, need careful assessment to select a drug either as an hypnotic or as an anxiolytic. Elimination half-lives vary considerably (Figure 2). However, although longer values increase the possibility of residual effects and of accumulation of the

parent drug or of an active metabolite, they are not necessarily disadvantageous. Insomnia with a significant anxiety component is best treated by a drug which provides an hypnotic effect overnight and an anxiolytic effect the next day. It is the relative advantages and disadvantages related to clinical need which require consideration, and in this paper attention will be given to the interplay between pharmacokinetics and residual effects on one hand, and efficacy and adverse effects on the other.

Residual effects

Residual impairment of performance with overnight ingestion of hypnotics is well established and with many hypnotics such effects extend well into the next day even with doses which are within the currently

Figure 2 Effective equilibration half-life of some benzodiazepines used as hypnotics. The data for flurazepam refers to its desalkyl metabolite. From Breimer (1979).

accepted therapeutic range. A variety of tasks have been used by several workers to investigate residual effects. Our work has related severity and persistence with dose. We have used visuomotor coordination (Borland & Nicholson, 1974), and in early studies with the barbiturate, heptabarbitalone, we have shown that decrements in performance persisted for 10 h after 200 mg, 13 h after 300 mg and 19 h after 400 mg (Figure 3). Sequelae were related to dose both in the decrement at any given time interval and in the persistence of the impairment; and in this way the studies supported previous observations (Von Felsinger, Lasagna & Beecher, 1953; Malpas, Rowan, Joyce & Scott, 1970; Bond & Lader, 1972) and showed, as did Kornetsky, Vates & Kessler (1959), that impaired performance persists far longer with higher doses which are nevertheless often within the accepted therapeutic range. These studies established the sensitivity of visuomotor coordination to the effects of drugs, and so we continued to use the technique, although we would emphasize the importance of the many other performance tests which have been used by other workers in the solution of the overall problem.

With the benzodiazepines we found that although performance was impaired for 16 h after flurazepam hydrochloride 30 mg and to at least 19 h after nitrazepam 10 mg (Borland & Nicholson, 1975a), residual effects of diazepam were much more limited. Indeed, impaired performance with diazepam 10 mg was restricted to a few hours after ingestion, and there was no evidence of residual impairment with overnight ingestion as long as the dose did not exceed 10 mg (Borland & Nicholson, 1977). It was with this finding that we undertook a detailed analysis of diazepam and its metabolites (Figure 4), 3-hydroxydiazepam

Figure 3 Residual effects of heptabarbitalone 200 mg (●), 300 mg (○) and 400 mg (▲) ingested overnight, on visuomotor coordination (arbitrary units). Severity and persistence of effects are related to dose. From Borland & Nicholson (1974). Significance levels: * $P > 0.05$; ** $P > 0.01$; *** $P > 0.001$.

(temazepam) and 3-hydroxy-N-desmethyldiazepam (oxazepam). With morning ingestion, coordination was impaired from 2.5–4.5 h after diazepam 10 mg, and from 0.5–2.5 h with temazepam 20 mg, but from 4.5–6.5 h after oxazepam 30 mg (Figure 2). There were no residual effects after diazepam 10 mg, temazepam 10, 20 and 30 mg, or after oxazepam 15 and 30 mg. However, over the dose range 10–30 mg temazepam there was a trend toward impaired performance, and with oxazepam 45 mg performance was impaired the next day (Clarke & Nicholson, 1978).

These observations were in agreement with other studies. Similar results have been reported with diazepam by Seppala, Kortilla, Hakkinens & Linnoila (1976) using coordination skills and visual functions related to driving. The recovery from impaired performance within a few hours after ingestion of diazepam has been observed by Hart, Hill, Bye, Wilkinson & Peck (1976) with a variety of tasks including auditory vigilance, reaction time, short-term memory and digit symbol substitution. With temazepam our results are comparable with those of Hindmarch (1975) and Roth, Piccione, Salis, Kramer & Kaffeman (1979); thus, a residual effect may exist with the 30 mg dose. Other workers have observed the slow onset of impaired performance with oxazepam, and Molander & Duvhok (1976) have recorded depres-

Figure 4 Structural formulae of diazepam, temazepam, nordiazepam and oxazepam. Temazepam and nordiazepam are primary metabolites of diazepam. The main metabolic pathway for diazepam is demethylation to nordiazepam (N-desmethyldiazepam). Hydroxylation to temazepam (3-hydroxydiazepam), is a minor pathway, as is demethylation of temazepam to oxazepam.

Figure 5 Immediate effect of diazepam 10 mg (●), temazepam 20 mg (▲) and oxazepam 30 mg (○) ingested in the morning, on visuomotor coordination (arbitrary units). From Clarke & Nicholson (1978).

sion of critical flicker fusion frequency 3.0 h after ingestion of oxazepam 20 and 40 mg and impaired coordination with oxazepam 40 mg.

Diazepam also has a long-acting metabolite, nordiazepam, with a half-life between 30 and 60 h (Mandelli, Tognoni & Garattini, 1978; Post, Lindgren, Bertler & Malmgren, 1977). Nordiazepam, which is also the active metabolite of fosazepam (Nicholson, Stone & Clarke, 1976) and potassium clorazepate, has anxiolytic activity, but it is difficult to establish effects on performance with these benzodiazepines. With the morning ingestion of nordiazepam 5–10 mg or potassium clorazepate 15 mg, there is a gradual decline in performance over the day (Borland & Nicholson, 1977; Clarke & Nicholson, 1978). It would seem that nordiazepam may impair the ability of subjects to maintain a high level of performance, and this may be due to its anxiolytic effect, which can be overcome to some extent by increased effort. Anxiolytic activity without impairment of performance is also seen with the 1,5-benzodiazepine, clobazam (Borland & Nicholson, 1975b) but, although this drug has a relatively short elimination half-life, it is unlikely to be suitable as hypnotic (Nicholson, Stone & Clarke, 1977).

The long-acting metabolite, nordiazepam, with its anxiolytic activity, may be a disadvantage in the daily use of diazepam as an hypnotic; thus, benzodiazepines the activity of which is uncomplicated by such metabolites, have been developed. Triazolam, which has a heterocyclic moiety across the methyl position (Figure 1), is an example of a triazolo-1,4-benzodiazepine (Rudzik, Hester, Tang, Straw & Friis (1973), and has a half-life of 3–5 h (Eberts *et al.*, 1979). In doses around 1 mg it has residual effects on performance (Veldkamp, Straw, Metzler & Demissianos, 1974), but with smaller doses the residual effect is much less marked or even absent. With triazolam 0.25 mg, the highest dose currently recommended in the United Kingdom, visuomotor coordination is impaired for only 5.0 h after ingestion, and is without residual effects when ingested overnight. With triazolam 0.50 mg there are minimal residual effects, but these disappear within 11.5 h of ingestion (Nicholson & Stone, 1980).

A further development in the modification of the benzodiazepine molecule has been that of flunitrazepam (Figure 1). Flunitrazepam has a nitro group like nitrazepam and an orthofluorophenyl moiety like flurazepam hydrochloride, and would seem to be a potent hypnotic. Flunitrazepam 1 mg overnight leads to impaired performance the next day (Bond & Lader, 1975), although it has been suggested from studies with daytime ingestion that this dose would be free of residual effects (Amrein, Cano, Hartmann, Ziegler & Dubois, 1979). However, the recovery from impaired performance may be much quicker when the task is being practised throughout the day, a

synergistic effect may exist with the relatively low level of circadian performance in the morning, and metabolism night and day may vary (Nicholson, Stone & Pascoe, 1980).

Nevertheless, there is the possibility that flunitrazepam may be a useful hypnotic without residual sequelae. Flunitrazepam 0.5 mg increases total sleep time, and decreases awake activity and drowsy sleep, and is free of residual effects (Nicholson & Stone, 1980). It has been suggested that the profile of the equilibration phase may lead to a rapid recovery from impaired performance (Amrein *et al.*, 1979), and in this way it would be similar to diazepam. However, it must be assumed, at least for the moment, that studies with overnight ingestion are more likely to reflect accurately residual effects than extrapolation from daytime studies, and that repeated ingestion of flunitrazepam 1 mg may lead to sequelae either as impaired performance or as an anxiolytic effect.

Sleep in the healthy adult

Although the drug itself may not lead to clear-cut impairments of performance (as with nordiazepam and clobazam), and persistence of impaired performance may not relate obviously to elimination half-lives (as with diazepam and possibly flunitrazepam), it is considered that hypnotics with mean elimination half-lives of less than 8 h are less likely and drugs with mean elimination half-lives greater than 16 h are more likely to have residual sequelae. In this way daily ingestion of nordiazepam 10 mg, and flunitrazepam 1 mg and above, will probably lead to residual effects either as impaired performance or as a daytime anxiolytic effect. On the other hand oxazepam 30 mg, triazolam 0.25 mg, and temazepam 20 mg can be expected to be without such effects. It is in the context of dose ranges free of residual sequelae that we have studied hypnotic activity, and have chosen diazepam as well as the benzodiazepines with short elimination half-lives.

In young adults diazepam 5–10 mg and temazepam 10–20 mg reduced sleep onset latencies and awake activity and increased total sleep time, whereas temazepam also reduced drowsy sleep (Nicholson & Stone, 1976; Nicholson, Stone & Clarke, 1976). The effect of oxazepam 15–30 mg was similar to that of temazepam, although it had no effect on sleep onset latencies, and this is consistent with performance studies and evidence from pharmacokinetics that the drug is poorly absorbed, at least, in its present formulation (Nicholson & Stone, 1978). In middle-aged patients the effects of diazepam and temazepam were less obvious. There was no increase in the total sleep time of middle-aged patients even though both drugs increased the much longer sleep of young adults, and although sleep onset latencies were similar between

the two groups, it was difficult to establish an effect in middle age except with diazepam. Essentially, the effect of diazepam and temazepam in middle age was to reduce awake activity (Nicholson & Stone, 1979). However, all three drugs, in the doses cited, did not alter slow wave sleep or REM sleep with single night ingestion, except that the first REM period was delayed with the higher dose of temazepam in young adults, though this effect was not seen in the middle aged.

With triazolam 0.25 and 0.50 mg, total sleep time was increased and awake activity and drowsy sleep decreased. There were no changes in slow wave sleep, although the first period of REM sleep was delayed with the higher dose. However, although triazolam reduced the duration and percentage of REM sleep during the early part of the night, REM sleep over the whole night was not changed. There was only marginal evidence of a greater hypnotic effect in the healthy individual with the higher dose of triazolam (Nicholson & Stone, 1980).

Effectiveness in insomnia

Currently, it is evident that the short-acting hypnotics, triazolam and temazepam, are the most likely drugs to have both a useful hypnotic profile and be free of residual effects on daily ingestion. It is therefore important to assess the effectiveness of these drugs in the insomniac. In a long-term study with insomniacs aged 46–62 yr, Mitler, Carskadon, Phillips, Sterling, Zarcone, Spiegel, Guilleminault & Dement (1979) have shown that temazepam 15–30 mg prolongs total sleep time and reduces night-time awakenings, and have demonstrated a trend towards reduced sleep latency. There was no evidence that REM sleep was modified, but with 30 mg there was some depression of slow wave sleep. There was also no evidence of tolerance. The effect of temazepam was similar to that of flurazepam in its efficacy and its effect on slow wave sleep (Dement, Carskadon, Mitler, Phillips & Zarcone, 1978), but flurazepam also suppressed REM sleep (Kales, Kales, Bixler & Slye, 1971; Kales, Kales, Bixler & Scharf, 1975). In outpatients with insomnia, Heffron & Roth (1979) and Fillingim (1979) have found that temazepam 30 mg increased total sleep time, reduced difficulty in falling asleep and improved sleep with less awakenings, particularly in the early morning.

Studies with triazolam in insomniacs have also established its efficacy. Roth, Kramer & Lutz (1976) have evaluated the effect of triazolam 0.5 mg on the sleep of insomniacs aged 35–55 years. Triazolam 0.5 mg was effective in inducing and maintaining sleep, but with a delay to the first REM period which has also been observed in healthy subjects of a similar age. In other studies, Vogel, Thurmond, Gibbons,

Edwards, Sloan & Sexton (1975) and Vogel, Barker, Gibbons & Thurmond (1976) have established the effectiveness of triazolam in doses up to 0.50 mg, whereas Sundaresan, Wardell, Weintraub & Lasagna (1979) have found it equally effective to flurazepam 30 mg.

It is with these observations in healthy individuals and in insomniacs, together with studies on performance, that a scheme for the use of hypnotics in general practice may be proposed (Nicholson, 1979), although it is accepted that clinical experience may usefully modify indications based on experimental studies. Potassium chlorazepate 15 mg, with its long-acting metabolite, nordiazepam, would seem to be

the drug of choice for the patient with insomnia secondary to anxiety because its anxiolytic activity has minimal effects on performance. For the patient with insomnia without significant psychopathology, temazepam 10–20 mg, triazolam 0.125–0.25 mg and occasionally diazepam 5–10 mg should provide the initial approach. Flurazepam hydrochloride 15–30 mg, nitrazepam 5–10 mg and flunitrazepam 1 mg and above, have persistent sequelae, and until the activity of flurazepam at the lower dose and of flunitrazepam at doses less than 1 mg are clarified, they should be reserved for refractory patients, and for those in whom some impairment of performance the next day would be acceptable (Figure 6).

Figure 6 Immediate and residual effects of two short-acting hypnotics, triazolam 0.25 mg (Δ) and temazepam 20 mg (▼), and two long-acting hypnotics, flurazepam 30 mg (○) and nitrazepam 10 mg (●), on visuomotor coordination (arbitrary units). The doses used are the maximum of the dose range presently recommended for each hypnotic in the United Kingdom. Composite data from Borland & Nicholson (1975a), Clarke & Nicholson (1978), Nicholson & Stone (1980). Significance levels: * $P > 0.05$, ** $P > 0.01$, *** $P > 0.001$.

Rebound insomnia

However, recently it has been suggested that short-acting hypnotics *per se* may lead to a considerable worsening of insomnia on withdrawal (Kales, Scharf & Kales, 1978; Kales, Scharf, Kales & Soldatos, 1979). Patients apparently suffer from worse insomnia than experienced before treatment, with prolonged sleep latencies and increased wakefulness after sleep onset. It is said to have been observed after flunitrazepam, nitrazepam and triazolam, although in each study there was little evidence of subjective complaints of insomnia. It has not been observed with diazepam or flurazepam hydrochloride, and on this evidence the proposed syndrome has been related to short elimination half-lives, although those of flunitrazepam and nitrazepam can hardly be considered short as they are between 15–30 h (Amrein, Cano & Hugin, 1976; Cano, Soliva, Hartmann, Ziegler & Amrein, 1977) and 18–36 h (Breimer, Bracht & DeBoer, 1977), respectively.

The possibility that rebound insomnia may occur with short-acting hypnotics is based on studies involving five benzodiazepines (Kales *et al.*, 1979), although when the data are reviewed, several uncertainties arise. Four studies are cited in which triazolam is believed to have led to rebound insomnia. It is accepted that the study by Kales, Kales, Bixler, Scharf & Russek (1976) shows an obvious deterioration of sleep with increased sleep latency and wakefulness on withdrawal after 14 d with triazolam 0.50 mg, but there were unusual features of these patients. They had a very high incidence of awakenings, even for insomniacs, and they became tolerant to triazolam within a relatively short period of time. This has not been observed by other workers. The patients used by Kales *et al.* (1976) may not represent the usual insomniac population, and worsening of sleep after medication may be a problem specific to them.

The three other papers which are quoted include the study by Roth *et al.* (1976), in which rebound insomnia was not observed in insomniacs treated for 14 d with triazolam 0.5 mg. The study by Vogel *et al.* (1975) has also failed to show any deterioration of sleep after triazolam 0.25 or 0.50 mg after 6 days. In a further paper by Vogel *et al.* (1976) there would seem to be a worsening of sleep on daily use with triazolam 0.50 mg, but careful examination of the data shows a considerable variability of baseline measures, and so it is not possible to deduce that a worsening of sleep

occurred after withdrawal from the drug.

The question of rebound insomnia with short-acting hypnotics is discussed in greater detail elsewhere (Nicholson, 1980), but it is considered that there is no experimental evidence to suggest that the proper use of a short-acting hypnotic such as triazolam leads to a deterioration of sleep on withdrawal, and this has been supported by studies with the other short-acting hypnotic, temazepam (Mitler *et al.*, 1979). However, it is accepted that some benzodiazepines may lead to disturbances of sleep and/or rebound insomnia, and nitrazepam and flunitrazepam may be implicated. The studies of Oswald & Priest (1965) and Adam, Adamson, Brezinova, Hunter & Oswald (1976) have certainly shown a worsening of insomnia on withdrawal from nitrazepam, and it is evident that flunitrazepam in doses of 1 mg and above has adverse effects on sleep (Monti & Allier, 1973; Javanovic, 1977; Gaillard & Phellipeau, 1977) as well as adverse effects on withdrawal (Bixler, Kales, Soldatos & Kales, 1974). Longer elimination half-lives may be associated with persistent effects both on performance and on sleep, but short elimination half-lives cannot be causally related to adverse effects on withdrawal.

Conclusions

It is evident that currently available benzodiazepines provide a wide spectrum of activity relevant to selected use in clinical medicine. The short-acting hypnotics, temazepam 10–20 mg and triazolam 0.125–0.25 mg, are free of residual sequelae in the doses cited, and there is no evidence that in these doses they have adverse effects on sleep or lead to a worsening of insomnia on withdrawal. Diazepam 5–10 mg is also likely to be free of residual sequelae when used occasionally. On the other hand nitrazepam 5–10 mg and flunitrazepam 1 mg and above, may adversely affect sleep, and these two drugs, together with flurazepam hydrochloride 15–30 mg, are likely to have persistent daytime effects. Additional information is needed to clarify the activity of low doses of flurazepam hydrochloride and of doses of flunitrazepam less than 1 mg. Short-acting hypnotics, such as temazepam and triazolam, should be used in the initial management of insomnia except in patients with significant anxiety when potassium clorazepate 15 mg is likely to be the drug of choice.

References

- ADAM, K., ADAMSON, L., BREZINOVA, V., HUNTER, W.M. & OSWALD, I. (1976). Nitrazepam: Lastingly effective but trouble on withdrawal. *Br. med. J.*, 1, 1558–1560.
- AMREIN, R., CANO, J.P., HARTMANN, D., ZIEGLER, W.H. & DUBOIS, R. (1979). Clinical and psychometric effects of flunitrazepam observed during the day in relation to pharmacokinetic data. In *Sleep Research*. Ed. Priest,

- R.G., Pletscher, A. & Ward, J. Pp. 83-98. Lancaster, England: MTP Press.
- AMREIN, R., CANO, J.P. & HUGIN, W. (1976). Pharmacokinetic and pharmacodynamic findings after a single intravenous, intramuscular and oral application of "Rohypnol". In *Bisherige Erfahrungen mit "Rohypnol" (Flunitrazepam) in der Anesthesiologie und Intensivtherapie*. Ed. Hugin, W., Hossli, C. & Gemperle, M. Pp. 39-56. Basel: Roche Editions.
- BIXLER, E.O., KALES, A., SOLDATOS, C.R. & KALES, J.D. (1974). Flunitrazepam, an investigational hypnotic drug: Sleep Laboratory evaluations. *J. clin. Pharmacol.*, **17**, 569-578.
- BOND, A.J. & LADER, M.H. (1972). Residual effects of hypnotics. *Psychopharmacologia Berlin*, **25**, 117-132.
- BOND, A.J. & LADER, M.H. (1973). The residual effects of flurazepam. *Psychopharmacologia Berlin*, **32**, 223-235.
- BOND, A.J. & LADER, M. (1975). Residual effects of flunitrazepam. *Br. J. clin. Pharmacol.*, **2**, 143-150.
- BORLAND, R.G. & NICHOLSON, A.N. (1974). Human performance after a barbiturate (heptabarbitalone). *Br. J. clin. Pharmacol.*, **1**, 209-215.
- BORLAND, R.G. & NICHOLSON, A.N. (1975a). Comparison of the residual effects of two diazepam (nitrazepam and flurazepam hydrochloride) and pentobarbitalone sodium. *Br. J. clin. Pharmacol.*, **2**, 9-17.
- BORLAND, R.G. & NICHOLSON, A.N. (1975b). Immediate effects on human performance of a 1,5-benzodiazepine (clobazam) compared with the 1,4-benzodiazepines, chlordiazepoxide hydrochloride and diazepam. *Br. J. clin. Pharmacol.*, **2**, 215-221.
- BORLAND, R.G. & NICHOLSON, A.N. (1977). Residual effects of potassium clorazepate, a precursor of nordiazepam. *Br. J. clin. Pharmacol.*, **4**, 86-89.
- BREIMER, D.D. (1979). Pharmacokinetics and metabolism of various benzodiazepines used as hypnotics. In: Temazepam and related 1,4-benzodiazepines. Eds. Lader, M.H., Makin, E. & Nicholson, A.N. *Br. J. clin. Pharmacol.*, **8**, 7S-13S.
- BREIMER, D.D., BRACHT, H. & DeBOER, A.G. (1977). Plasma level profile of nitrazepam following oral administration. *Br. J. clin. Pharmacol.*, **4**, 709-711.
- CANO, J.P., SOLIVA, M., HARTMANN, D., ZIERLER, W.H. & AMREIN, R. (1977). Bioavailability of various galenic formulations of flunitrazepam. *Arzneimittel-Forschung*, **27**, 2383-2395.
- CLARKE, C.H. & NICHOLSON, A.N. (1978). Immediate and residual effects in man of the metabolites of diazepam. *Br. J. clin. Pharmacol.*, **6**, 325-333.
- DEMENT, W.C., CARSKADON, M.A., MITLER, M.M., PHILLIPS, R.L. & ZARCONI, V.P. (1978). Prolonged use of flurazepam: A sleep laboratory study. *Behav. Med.*, **5**, 25-31.
- EBERTS, F.S., PHILOPOULOS, Y. & VLIEK, R.W. (1979). Disposition of 14C-Triazolam; a short-acting hypnotic, in man. *The Pharmacologist*, **21** (3), 131.
- FILLINGHIM, J.M. (1979). Double-blind evaluation of the efficacy and safety of temazepam in out-patients with insomnia. In: Temazepam and related 1,4-benzodiazepines. Effects on sleep and on performance. Ed. Lader, M.H., Makin, E. & Nicholson, A.N. *Br. J. clin. Pharmacol.*, **8**, 73S-77S.
- FUCELLA, L.M., BOLCIONI, G., TAMASSIA, V., FERRARIO, L. & TOGNONI, G. (1977). Human pharmacokinetics and bioavailability of temazepam administered in soft gelatin capsules. *Eur. J. clin. Pharmacol.*, **12**, 383-386.
- GAILLARD, J.M. & PHELLIPEAU, M. (1977). Modification des contenus mentaux en stade paradoxal et en stade 2 sous l'effet d'une benzodiazepine le flunitrazepam. *L'Encephale*, **3**, 35-47.
- GREENBLATT, D.J., JOYCE, T.H., COMER, W.H., KNOWLES, J.A., SHADER, R.I., KYRIAKOPOULOS, A.A., MACLAUGHLIN, D.S. & RUELIUS, H.W. (1977). Clinical pharmacokinetics of lorazepam. II. Intramuscular injection. *Clin. Pharmacol. Ther.*, **21**, 222-230.
- HART, J., HILL, H.M., BYE, C.E., WILKINSON, R.T. & PECK, A.W. (1976). The effects of low doses of amylobarbitone sodium and diazepam on human performance. *Br. J. clin. Pharmacol.*, **3**, 289-298.
- HEFFRON, W.A. & ROTH, P. (1979). Double-blind evaluation of the safety and hypnotic efficacy of temazepam in insomniac patients. In: temazepam and related 1,4-benzodiazepines. Effects on sleep and on performance. Ed. Lader, M.H., Makin, E. & Nicholson, A.N. *Br. J. clin. Pharmacol.*, **8**, 69S-72S.
- HINDMARCH, I. (1975). A 1,4-benzodiazepine: Temazepam (k3917): Its effect on some psychological parameters of sleep and behaviour. *Arzneimittel-Forschung*, **25**, 1836-1839.
- JAVANOVIC, U.J. (1977). Polygraphic sleep recordings before and after the administration of flunitrazepam. *J. int. Med. Res.*, **5**, 77-84.
- KALES, A., SCHARF, M.B. & KALES, J.D. (1978). Rebound insomnia: A new clinical syndrome. *Science*, **201**, 1039-1041.
- KALES, J., KALES, A., BIXLER, E.O. & SLYE, E.S. (1971). Effects of placebo and flurazepam on sleep patterns in insomniac subjects. *Clin. Pharmacol. Ther.*, **12**, 691-697.
- KALES, A., KALES, J., BIXLER, E. & SCHARF, M. (1975). Effectiveness of hypnotic drugs with prolonged use. *Clin. Pharmacol. Ther.*, **13**, 356-363.
- KALES, A., SCHARF, M.B., KALES, J.D. & SOLDATOS, C.R. (1979). Rebound insomnia. A potential hazard following withdrawal of certain benzodiazepines. *J. Am. med. Assoc.*, **241**, 1692-1695.
- KALES, A., KALES, J.D., BIXLER, E.O., SCHARF, M.B. & RUSSEK, E. (1976). Hypnotic efficacy of triazolam: Sleep laboratory evaluation of intermediate-term effectiveness. *J. clin. Pharmacol.*, **16**, 399-406.
- KAPLAN, S.A., deSILVA, J.A.F., JACK, M.L., ALEXANDER, K., STROJNY, N., WEINFELD, R.E., PUGLIST, C.V. & WEISSMAN, L. (1973). Blood level profile in man following chronic oral administration of flurazepam hydrochloride. *J. Pharmacol. Sci.*, **62**, 1932-1935.
- KORNETSKY, C., VATES, T.W. & KESSLER, E.K. (1959). A comparison of hypnotic and residual psychological effects of single doses of chlorpromazine and secobarbital in man. *J. Pharmacol. exp. Ther.*, **127**, 51-54.
- MALPAS, A., ROWAN, A.J., JOYCE, C.R.B. & SCOTT, D.F. (1970). Persistent behavioural and electroencephalographic changes after single doses of nitrazepam and amylobarbitone sodium. *Br. med. J.*, **2**, 762-764.
- MANDELLI, M., TOGNONI, G. & GARATTINI, S. (1978). Clinical pharmacokinetics of diazepam. *Clin. Pharmacol.*, **3**, 72-91.
- MITLER, M.M., CARSKADON, M.A., PHILLIPS, R.L., STERLING, W.R., ZARCONI, V.P., SPIEGEL, R., GUILLEMINAULT, C. & DEMENT, W.C. (1979). Hyp-

- notic efficacy of temazepam: A long term sleep laboratory evaluation. In: Temazepam and related 1,4-benzodiazepines: Effects on sleep and on performance. Ed. Lader, M.H., Makin, E.J.B., & Nicholson, A.N. *Br. J. clin. Pharmac.*, **8**, 63S-68S.
- MOLANDER, L. & DUVHOK, C. (1976). Acute effects of oxazepam, diazepam and methylperone, alone and in combination with alcohol in sedation, coordination and mood. *Acta. Pharmac. Toxic.*, **38**, 145-160.
- MONTI, M. & ALLIER, H. (1973). Flunitrazepam (Ro 5-4200) and sleep cycle in normal subjects. *Psychopharmacologia Berlin*, **32**, 343-349.
- NICHOLSON, A.N. (1979). Hypnotics today. *The Practitioner*, **223**, 479-484.
- NICHOLSON, A.N. (1980). Hypnotics: Rebound insomnia and residual sequelae. *Br. J. clin. Pharmac.*, **9**, 223-225.
- NICHOLSON, A.N. & STONE, B.M. (1976). Effect of a metabolite of diazepam, 3-hydroxydiazepam (temazepam) on sleep in man. *Br. J. clin. Pharmac.*, **3**, 543-550.
- NICHOLSON, A.N. & STONE, B.M. (1978). Hypnotic activity of 3-hydroxy, N-desmethyldiazepam. *Br. J. clin. Pharmac.*, **5**, 469-472.
- NICHOLSON, A.N. & STONE, B.M. (1979). Diazepam and 3-hydroxydiazepam (temazepam) on sleep of middle age. *Br. J. clin. Pharmac.*, **7**, 463-468.
- NICHOLSON, A.N. & STONE, B.M. (1980). Activity of the hypnotics, flunitrazepam and triazolam in man. *Br. J. clin. Pharmac.*, **9**, 187-194.
- NICHOLSON, A.N., STONE, B.M. & CLARKE, C.H. (1976). Effect of diazepam and fosazepam (a soluble derivative of diazepam) on sleep in man. *Br. J. clin. Pharmac.*, **3**, 533-541.
- NICHOLSON, A.N., STONE, B.M. & CLARKE, C.H. (1977). Studies on the effect of the 1,5-benzodiazepines, clobazam and triflubazam, on sleep in man. *Br. J. clin. Pharmac.*, **4**, 567-572.
- NICHOLSON, A.N., STONE, B.M. & PASCOE, P.A. (1980). Studies on sleep and performance with a triazolo-1,4-thienodiazepine (brotizolam). *Br. J. clin. Pharmac.*, **10**, 75-81.
- OSWALD, I. & PRIEST, R.G. (1965). Five weeks to escape the sleeping pill habit. *Br. med. J.*, **2**, 1093-1099.
- POST, C., LINDGREN, S., BERTLER, A. & MALMGREN, H. (1977). Pharmacokinetics of N-desmethyldiazepam in healthy volunteers after single doses of dipotassium clorazepate. *Psychopharmacology*, **53**, 105-109.
- ROTH, T., KRAMER, M. & LUTZ, T. (1976). Intermediate use of triazolam: A sleep laboratory study. *J. int. Med. Res.*, **4**, 59-63.
- ROTH, T., PICCIONI, P., SALIS, P., KRAMER, M. & KAFFEMANN, M. (1979). Effects of temazepam, flurazepam and quinalbarbitone on sleep: Psychomotor and cognitive function. In: Temazepam and related 1,4-benzodiazepines: Effects on sleep and on performance. Eds. Lader, M.H., Makin, E.J.B. & Nicholson, A.N. *Br. J. clin. Pharmac.*, **8**, 47S-54S.
- RUDZIK, A.D., HESTER, J.B., TANG, A.H., STRAW, R.N. & FRIIS, W. (1973). Triazolo-benzodiazepines, a new class of central nervous system-depressant compounds. In *The Benzodiazepines*. Ed. Garattini, S., Mussini, E. & Randall, L.O. New York: Raven Press.
- SEPPALA, T., KORTILLA, K., HAKKINENS, S. & LINNOILA, M. (1976). Residual effects and skills related to driving after a single oral administration of diazepam, medazepam or lorazepam. *Br. J. clin. Pharmac.*, **3**, 831-841.
- SJOQVIST, F. & SUNDWALL, A. (Eds) (1977). The pharmacokinetic profile of oxazepam. *Acta Pharmac. Toxic.*, **40**, suppl. I.
- SUNDARESAN, P.R., WARDELL, W.M., WEINTRAUB, M. & LASAGNA, L. (1979). Methodology for demonstrating sustained efficacy of hypnotics: A comparative study of triazolam and flurazepam. *Clin. Pharmac. Ther.*, **25**, 391-398.
- VELDKAMP, W., STRAW, R.N., METZLER, C.M. & DEMISIANOS, H.V. (1974). Efficacy and residual effect evaluation of a new hypnotic, triazolam. *J. clin. Pharmac.*, **14**, 102-111.
- VOGEL, G., THURMOND, A., GIBBONS, P., EDWARDS, K., SLOAN, K.B. & SEXTON, K. (1975). The effect of triazolam on the sleep of insomniacs. *Psychopharmacologia Berlin*, **41**, 65-69.
- VOGEL, G., BARKER, K., GIBBONS, P. & THURMOND, A. (1976). A comparison of the effects of flurazepam 30 mg and triazolam 0.5 mg on the sleep of insomniacs. *Psychopharmacology*, **47**, 81-86.
- VON FELSINGER, J.M., LASAGNA, L. & BEECHER, H.K. (1953). The persistence of mental impairment following a hypnotic dose of a barbiturate. *J. Pharmac. exp. Ther.*, **109**, 284-291.