LABETALOL: THE NINETEEN-EIGHTIES

J.I.S. ROBERTSON

Medical Research Council, Blood Pressure Unit, Western Infirmary, Glasgow G11 6NT, UK

Introduction

The present symposium on the role of labetalol, a drug with both α - and β -adrenoceptor-blocking actions, in clinical medicine, is timely. Two previous supplements to this journal (Richards & Turner, 1976; Richards & Prichard, 1979) reviewed the earlier studies with this so far unique agent (see also Brogden *et al.*, 1978). The past few years have seen increasingly more widespread use of the drug, and hence greater familiarity with its clinical value and enhanced assurance in its applications. This has coincided with an acceptance of β -adrenoceptor blockade not only as first-choice therapy in hypertension (WHO, 1978) but also in the secondary prevention of myocardial infarction (Wilhelmssen *et al.*, 1974; Andersen *et al.*, 1979; Norwegian Study, 1981; Sleight, 1981; Hampton, 1981). The evidence for a primary preventive effect of β -blockade on ischaemic heart disease, when used as treatment for hypertension, though suggestive, is still inconclusive (Robertson, 1978, 1979). Concurrently, there has been increasing appreciation of the value of post-synaptic α -adrenoceptor blockade in the treatment of hypertension (Gross, 1982) and also of congestive cardiac failure (Symposium on Congestive Heart Failure, 1981). Labetalol, a drug which combines actions applicable in all these areas, could therefore scarcely escape the attentions of the general physician, the clinical pharmacologist and the cardiologist alike.

Clinical pharmacology

The place of β -adrenoceptor blockade and of post-synaptic α -blockade in the treatment of hypertension was reviewed at the present symposium by Gross (1982).

The decrease in pressure with β -blockade is of modest extent and independent of orthostasis; it is seen with all β -blockers regardless of cardioselectivity and of intrinsic sympathomimetic activity. The mechanism of this valuable hypotensive effect remains in the view of Gross (1982) without satisfactory explanation: reduction of cardiac output and suppression of renin secretion seem inadequate. A possible central site of action has also been discussed, but since entry to the central nervous system

varies widely with different β -blockers, although all reduce blood pressure similarly, it seems to Gross (1982) highly improbable that accumulation in the brain is responsible for the anti-hypertensive effect. However, Rand *et al.* (1981) have provided evidence that low doses of adrenaline stimulate presynaptic β -receptors, and so facilitate release of sympathetic neurotransmitter. The effect is prevented by β -adrenoceptor antagonists and this might therefore be a major anti-hypertensive property of β -blockers.

Blockade of postsynaptic α_1 -adrenoceptors has also been shown to cause consistent and sustained reduction in arterial pressure. Earlier so-called "classical" α -adrenolytic agents such as phentolamine, were disappointing in their hypotensive effect. One likely explanation for this is that phentolamine blocks both α_1 - and α_2 -presynaptic adrenoceptors (Drew, 1979). The neurotransmitter noradrenaline stimulates the presynaptic α_2 -adrenoceptor; this action inhibits further noradrenaline release and consequently limits stimulation of α_1 -postsynaptic adrenoceptors on the effector site. Thus blockade of both pre- and postsynaptic α -adrenoceptors is likely to yield a poor or even insignificant hypotensive effect because the presynaptic α_2 -blockade will permit continued noradrenaline release which because of the competitive nature of the blockade, will overcome the α_1 -blockade. However, it is now known that drugs like labetalol or prazosin compete with noradrenaline at postsynaptic α_1 -adrenoceptors, while leaving the presynaptic α_2 -adrenoceptors unoccupied. Thus physiological inhibition of noradrenaline release is unimpaired. This selectivity of postsynaptic α_1 -adrenoceptor blockade seems to be an important aspect of the effectiveness of labetalol and of prazosin in reducing blood pressure.

Labetalol combines in the same molecule non-selective β -blockade with postsynaptic α_1 -blockade, the effect on β -receptors predominating by some 4:1. In pharmacological studies in man, this association results in an enhanced blood pressure lowering effect compared with plain β -blockers and a reduced tendency to orthostatic reactions compared with plain α_1 -blockers.

McNeill *et al.* (1982) have found that during the

chronic treatment of hypertension, mean steady-state plasma labetalol concentrations were consistently higher than would have been predicted from initial pharmacokinetic studies. Thus, bioavailability of labetalol increases with chronic dosage. Patients with lower steady-state plasma levels on fixed labetalol dose usually require higher maintenance doses for blood pressure control.

Labetalol has a large apparent volume of distribution and a relatively high plasma clearance; these aspects seem to be unchanged in severe renal failure (Wood *et al.*, 1982).

Labetalol 100 mg intravenously was found by Agabiti-Rosei *et al.* (1982) to cause a reduction in blood pressure and heart rate within 5 min. These effects were greater in patients with the higher pretreatment plasma noradrenaline values, an observation which seems to accord with the predictions of Rand *et al.* (1981). Cardiac output was not significantly altered in these acute studies. Cohn *et al.* (1982) similarly found that blood pressure reduction with oral labetalol over 7 d was associated with an unchanged cardiac output. However, blood pressure reduction with oral labetalol given for 2.5 weeks was found by Fagard *et al.* (1982) to be accompanied by a reduction in heart rate and cardiac output both at rest and during maximal exercise. Total peripheral resistance was not affected at recumbent rest, but was reduced sitting and at exercise. Pulmonary arterial and capillary wedge pressures were reduced only in the sitting position; pulmonary vascular resistance remained unchanged.

These various studies generally confirm the consensus of earlier reports showing that blood pressure reduction is mainly due to reduced peripheral resistance, with little accompanying change in resting heart rate or cardiac output (Prichard & Richards, 1982). Significant lowering of plasma renin and aldosterone was also confirmed (Fagard *et al.*, 1982).

Long-term labetalol did not cause significant reduction in glomerular filtration rate in hypertensive patients with or without renal impairment (Malini *et al.*, 1982; Wood *et al.*, 1982). Renal plasma flow increased and renal vascular resistance decreased both at rest and during exercise after labetalol treatment (Malini *et al.*, 1982). No modification of dosage is advised in renal impairment (Wood *et al.*, 1982).

Labetalol is as effective in reducing systolic and diastolic pressures over 24 h whether given twice or three times daily (Mancia *et al.*, 1982). Indeed, Breckenridge *et al.* (1982) found that a single daily dose was effective, but advised that a twice daily schedule is generally more suitable because a large (>1 g) single dose might sometimes cause nausea and postural hypotension.

Essential hypertension

The now extensive experience of the use of labetalol in treating essential hypertension has been reviewed by Prichard and Richards (1982) and by Breckenridge *et al.* (1982). The decrease in blood pressure with oral labetalol is rapid, being evident in 2 h and maximal at about 3 h. Tolerance to the hypotensive effect is not a feature with prolonged therapy.

In a variety of studies, using both fixed and titrated doses, labetalol has been found to be at least as effective in lowering arterial pressure as a wide range of plain β -blockers given alone, as β -blockers plus diuretics, β -blockers plus vasodilators, β -blockers plus phentolamine, adrenergic neurone blockers, and methyldopa. Administered alone, or with a diuretic, labetalol is often effective when other regimes, including β -blockers, have failed (Waal-Manning *et al.*, 1982; Kane, 1982).

Labetalol has now been extensively used in both hospital and general practice in Europe, Australia, New Zealand and Japan (Prichard & Richards, 1982; Breckenridge *et al.*, 1982; Waal-Manning & Simpson, 1982; Takeda *et al.*, 1982).

Side-effects of oral labetalol include postural hypotension, nausea, epigastric discomfort, vomiting, tingling of the scalp, lassitude, diarrhoea, constipation, a variety of skin rashes, stranguity (in male patients), impaired sexual drive and performance, and vivid dreams. Quantitatively, the burden of side-effects with labetalol seems roughly equivalent to that of simple β -blocking drugs, although certain features of β -blockers, such as vivid dreams, Raynaud's phenomenon, asthma and cardiac failure, are less frequent (but not absent) with labetalol. The adverse effects of labetalol seem most pronounced in the early weeks of treatment and may often subside with continued therapy. The two most prominent side-effects, nausea and postural hypotension, are clearly dose-related; they are infrequent with single doses of less than 400 mg, and with total daily doses less than 1 g. The reported withdrawal rate from labetalol therapy because of side-effects ranged from 25% in the New Zealand study (Waal-Manning & Simpson, 1982), to 6-13% in general practice (Kane, 1982), and to only 4% in the Japanese multicentre study (Takeda *et al.*, 1982). Side-effects seemed to be distinctly minimized when the starting dose was low – only 150 mg daily was used initially in the Japanese trial (Takeda *et al.*, 1982).

The final daily requirement in essential hypertension might be as low as 200 mg and as high as 2 g, given in two doses. There is a suggestion that while Londoners of West Indian or West African origin are rather resistant to labetalol (Jennings &

Parsons, 1976), Japanese may be particularly sensitive (Takeda *et al.*, 1982).

Phaeochromocytoma

The value of labetalol in controlling the hypertension and tachycardia of phaeochromocytoma, reported in earlier studies (Agabiti-Rosei *et al.*, 1976; Kaufman, 1979) has been extensively confirmed. Takeda *et al.* (1982) found that good blood pressure control was obtained in 21 of 30 patients with this disease. Labetalol was more effective in patients with predominantly adrenaline-secreting tumours, and in patients with sustained rather than paroxysmal hypertension. The former observation is of interest in view of the pressor effect of infused adrenaline reported by Rand *et al.* (1981).

Labetalol is also effective in the analagous situation of severe hypertension following clonidine withdrawal (Cumming *et al.*, 1982).

Intravenous labetalol

For the rare circumstances requiring parenteral anti-hypertensive therapy, labetalol offers several advantages over alternative drugs, as it should permit controlled steady lowering of arterial pressure without increasing heart rate or cardiac output. Thus the two particular dangers of intravenous therapy, cerebral and myocardial infarction, are minimized. Graded incremental infusion of labetalol has been found to be the most effective method of parenteral administration and that least prone to unwanted effects (Cumming *et al.*, 1982; Dal Palu *et al.*, 1982). This approach has successfully been used in hypertensive encephalopathy, and in hypertensive crises associated with phaeochromocytoma, aortic coarctation and pregnancy (Cumming *et al.*, 1982).

Ischaemic heart disease

The theoretical advantages of using labetalol in managing acute myocardial infarction have been reviewed by Timmis *et al.* (1982). β -blockade alone should overall be beneficial, in that myocardial oxygen supply would be enhanced and demand diminished by a reduction in heart rate and contractility and in blood pressure, while these advantages are only partly offset by a lowering of coronary perfusion pressure and increased coronary arterial tone. However, labetalol should have more favourable effects than plain β -blockers. Blockade of both α - and β -receptors should minimize the

tendency to coronary artery spasm induced by β -blockade alone, while any possible dysrhythmic effect of α -stimulation (Sheridan *et al.*, 1980) would also be offset. However, too marked a decrease in blood pressure could jeopardize myocardial perfusion. Pilot studies (Timmis *et al.*, 1982) showed that infusing labetalol intravenously to hypertensive patients with acute myocardial infarction, starting at 500 μ g/min and increasing by 500 μ g every 30 min as necessary, reduced blood pressure and heart rate, leaving cardiac output little altered.

Condorelli *et al.* (1982), in a study of ten normotensive patients with angiographic evidence of coronary artery disease, found that labetalol reduced myocardial oxygen consumption and increased exercise tolerance, without impairing left ventricular function.

Opie *et al.* (1982) also considered that labetalol should be effective in alleviating angina in hypertensive patients, by reducing arterial pressure and minimizing coronary spasm.

Labetalol, unlike plain β -blockers, is unlikely to provoke cardiac failure, presumably because the indirect positive inotropic effect of afterload reduction counters the direct negative inotropic effect of β -blockade (Opie *et al.*, 1982; Timmis *et al.*, 1982; Breckenridge *et al.*, 1982). However, labetalol should not be given to normotensive patients actually in cardiac failure unless they are already receiving digitalis and diuretics.

Overall, these initial exploratory steps in the use of labetalol in ischaemic heart disease are encouraging; more extensive, and more definitive, reports are awaited.

Pregnancy hypertension

Oral labetalol, in doses up to 1200 mg daily, has now been extensively used in pregnant women with severe hypertension. None of 91 infants born to women treated with labetalol through pregnancy showed congenital malformations. Maternal side-effects were few and minor, comprising postural hypotension, lethargy, scalp tingling and skin rash (Michael, 1982).

Intravenous labetalol by infusion has been safely and effectively used in treating severe pregnancy-induced hypertension (Cumming *et al.*, 1982).

Anaesthesia

Labetalol has been valuable in anaesthetic practice to correct uncontrolled hypertension before anaesthesia and surgery, when it can be given either orally or intravenously; to counter acute hyper-

tensive responses during laryngoscopy and various surgical procedures; to prevent hypertension and arrhythmias during removal of phaeochromocytoma; and, combined with halothane, to induce controlled hypotension (Scott, 1982).

Conclusions

The extensive clinical experience surveyed at this symposium left no doubt that labetalol is now well-established as being free from serious toxic effects. Moreover, no adverse maternal or foetal features followed its use in pregnancy.

There was proper concern nevertheless with the frequency and severity of side-effects. The consensus of a now substantial number of trials was that quantitatively the overall side-effect burden was no greater than with plain β -blockers, while some specific disadvantages, notably Raynaud's

phenomenon, and asthma, were less prominent with labetalol. Not surprisingly, side-effects were dose-related, and several speakers emphasized the value of introducing labetalol at low doses (150-200 mg daily) in circumventing these. Postural hypotension seems unlikely to be encountered at total daily doses under 1 g. Twice daily dosage is adequate to maintain control of blood pressure through 24 h.

Was labetalol introduced before its time? There is little doubt that the full advantages of postsynaptic α -adrenergic blockade were not appreciated at the time of the Torquay Symposium (Richards & Turner, 1976), and indeed were less clear than they have now become at the London meeting (Richards and Prichard, 1979). Thus, the full potential of labetalol, especially in ischaemic heart disease, remains to be exploited. Perhaps, as Gross (1982) suggested in his introductory review, the best days of labetalol are yet to come.

References

- AGABITI-ROSEI, E., ALICANDRI, C.L., BESCHI, M., CASTELLANO, M., FARIELLO, R., MONTINI, E., MUIESAN, M.L., ROMANELLI, G., MUIESAN, G. (1982). The acute and chronic hypotensive effect of labetalol and the relationship with pre-treatment plasma noradrenaline levels. *Br. J. clin. Pharmac.*, **13**, suppl. 1 87S-92S.
- AGABITI-ROSEI, E., BROWN, J.J., LEVER, A.F., ROBERTSON, A.S., ROBERTSON, J.I.S. & TRUST, P.M. (1976). Treatment of phaeochromocytoma and of clonidine withdrawal hypertension with labetalol. *Br. J. clin. Pharmac.*, **3**, suppl. 3, 809-815.
- ANDERSEN, M.P., FREDERIKSEN, J. & JURGENSEN, H.J. (1979). Effect of alprenolol on mortality among patients with definite or suspected myocardial infarction. *Lancet*, **2**, 865-868.
- BRECKENRIDGE, A., ORME, M., SERLIN, M.J. & MACIVER, M. (1982). Labetalol in essential hypertension. *Br. J. clin. Pharmac.*, **13**, suppl. 1 37S-39S.
- BROGDEN, R.N., HEEL, R.C., SPEIGHT, T.M. & AVERY, G. (1978). Labetalol: a review of its pharmacology and therapeutic use in hypertension. *Drugs*, **15**, 251-270.
- COHN, J., MEHTA, J. & FRANCIS, G.S. (1982). A review of the unique haemodynamic effects of labetalol in man. *Br. J. clin. Pharmac.*, **13**, suppl. 1 19S-26S.
- CONDORELLI, M., BREVETTI, G., CHIARIELLO, M., MICELI, D., LAVECCHIA, G., PAUDICE, G., VERRIENTI, S. (1982). Effects of combined alpha and beta blockade by labetalol in patients with coronary artery disease. *Br. J. clin. Pharmac.*, **13**, suppl. 1 101S-110S.
- CUMMING, A.M.M., BROWN, J.J., LEVER, A.F. & ROBERTSON, J.I.S. (1982). Intravenous labetalol in the treatment of severe hypertension. *Br. J. clin. Pharmac.*, **13**, suppl. 1 93S-96S.
- DAL-PALU, C., PESSINA, A.C., SEMPLICINI, A., HLEDE, M., MORANDIN, F., PALATINI, P., SPERTI, G. & ROSSI, G.P. (1982). Intravenous labetalol in severe hypertension. *Br. J. clin. Pharmac.*, **13**, suppl. 1 97S-99S.
- DREW, G.M. (1979). Pre-synaptic alpha adrenoceptors and their pharmacological characterisation and functional significance, *Advances in Bioscience*, Vol. 18. Ed. Langer, Starke & Dubocovic. Oxford: Pergamon Press.
- FAGARD, R., LIJNEN, P. & ANTOON, A. (1982). Response of the systemic and pulmonary circulation to labetalol at rest and during exercise. *Br. J. clin. Pharmac.*, **13**, suppl. 1 13S-17S.
- GROSS, F. (1982). The place of alpha adrenoceptor and beta adrenoceptor blockade in the treatment of hypertension. *Br. J. clin. Pharmac.*, **13**, suppl. 1 5S-11S.
- HAMPTON, J.R. (1981). The use of beta-blockers for the reduction of mortality after myocardial infarction. *Eur. Heart J.*, **2**, 259-268.
- JENNINGS, K. & PARSONS, V. (1976). A study of labetalol in patients of European, West Indian and West African origin. *Br. J. clin. Pharmac.*, **3**, suppl. 3, 773-775.
- KANE, J.A. (1982). Labetalol in general practice. *Br. J. clin. Pharmac.*, **13**, suppl. 1 59S-63S.
- KAUFMAN, L. (1979). Use of labetalol during hypotensive anaesthesia and in the management of phaeochromocytoma. *Br. J. clin. Pharmac.*, **8**, suppl. 2, 229-232.
- MALINI, P.L., STROCCHI, E., NEGRONI, S., AMBORSIONI, E. & MAGNANI, B. (1982). Renal haemodynamics after chronic treatment with labetalol and propranolol. *Br. J. clin. Pharmac.*, **13**, suppl. 1 123S-126S.
- MANCIA, G., POMIDOSSI, G., PARATI, G., BERTINEIRI, G., GRASSI, G., NAVONE, F., FERRARI, A.,

- GREGORINI, L. & ZANCHETTI, A. (1982). 24 Hour blood pressure response to labetalol in twice and thrice daily administration. *Br. J. clin. Pharmacol.*, **13**, suppl. 1 27S-35S.
- MCNEIL, J.J., ANDERSON, A.E. & LOUIS, W.J. (1982). Labetalol steady state pharmacokinetics in hypertensive patients. *Br. J. clin. Pharmacol.*, **13**, suppl. 1 75S-80S.
- MICHAEL, C.A. (1982). The evaluation of labetalol in the treatment of hypertension complicating pregnancy. *Br. J. clin. Pharmacol.*, **13**, suppl. 1 127S-131S.
- NORWEGIAN MULTICENTRE STUDY GROUP. (1981). Timolol-induced reduction in mortality and reinfarction in patients surviving acute myocardial infarction. *New Eng. J. Med.*, **304**, 801-807.
- OPIE, L.H., WHITE, D., LEE, J. & LUBBE, W.F. (1982). Alternatives to beta blockade in therapy of hypertension with angina pectoris: role of nifedipine or of labetalol. *Br. J. clin. Pharmacol.*, **13**, suppl. 1 115S-122S.
- PRICHARD, B.N.C. & RICHARDS, D.A. (1982). Comparison of labetalol with other antihypertensive drugs. *Br. J. clin. Pharmacol.*, **13**, suppl. 1 41S-47S.
- RAND, M.J., MAJEWSKI, H. & TUNG, L.-H. (1981). Adrenaline-induced hypertension in rats. Abstract No. 360. *Eighth Scientific Meeting of the International Society of Hypertension*. Milan, May 31-June 3, 1981.
- RICHARDS, D.A. & PRICHARD, B.N.C. (Eds.) (1976). Proceedings of a Symposium on Labetalol, Torquay, April, 1976. *Br. J. clin. Pharmacol.*, **3**, suppl. 3, 677-824.
- RICHARDS, D.A. & TURNER, P. (Eds.) (1979). Proceedings of the Second Symposium on Labetalol, London, March, 1979. *Br. J. clin. Pharmacol.*, **8**, suppl. 2, 85-244.
- ROBERTSON, J.I.S. (1978). Complications of mild hypertension: Discussion. *Ann. N.Y. Acad. Sci.*, **304**, 64-65.
- ROBERTSON, J.I.S. (1979). Hypertension, ischaemic heart disease and left ventricular hypertrophy. *Roy. Soc. Med. Int. Cong. Symp. Series*, No. 9, Pp. 1-4.
- SCOTT, D.B. (1982). The use of labetalol in anaesthesia. *Br. J. clin. Pharmacol.*, **13**, suppl. 1 133S-155S.
- SHERIDAN, D.J., PENKOSKE, P.A., SOBEL, B.E. & CORR, P.B. (1980). Alpha-adrenergic contribution to dysrhythmia during myocardial infarction and reperfusion in cats. *J. clin. Invest.*, **65**, 161-171.
- SLEIGHT, P. (1981). Beta-blockade after myocardial infarction. *New Eng. J. Med.*, **304**, 837-838.
- SYMPOSIUM ON CONGESTIVE HEART FAILURE. (1981). *Am. J. Med.*, **71**, 131-160.
- TAKEDA, T., KANEKO, Y., OMEA, T., YOSHINAGA, K., MUSUYAMA, Y., NUKADA, T. & SHIGIYA, R. (1982). The use of labetalol in Japan. Results of multicentre clinical trials. *Br. J. clin. Pharmacol.*, **13**, suppl. 1 48S-57S.
- TIMMIS, A.D., FOWLER, M.B., JAGGARAO, N.S.V., VINCENT, R. & CHAMBERLAIN, D.A. (1982). The role of labetalol in acute myocardial infarction. *Br. J. clin. Pharmacol.*, **13**, suppl. 1 111S-114S.
- WAAL-MANNING, H.J. & SIMPSON, F.O. (1982). Review of long term treatment with labetalol. *Br. J. clin. Pharmacol.*, **13**, suppl. 1 65S-73S.
- WHO: ARTERIAL HYPERTENSION. (1978). Proceedings of a WHO Expert Committee. WHO Technical Report Series No. 628. Pp. 1-58.
- WILHELMSSON, C., VEDIN, J.A., WILHELMSEN, L., TIBBLIN, G. & WERKÖ, L. (1974). Reduction of sudden deaths after myocardial infarction by treatment with alprenolol. *Lancet*, **2**, 1157-1163.
- WOOD, A.J., FERRY, D.G. & BAILEY, R.R. (1982). Elimination kinetics of labetalol in severe renal failure. *Br. J. clin. Pharmacol.*, **13**, suppl. 1 81S-86S.