BIAS

G. ROSE

Department of Medical Statistics and Epidemiology,
London School of Hygiene and Tropical Medicine,
Keppel Street (Gower Street), London WC1E 7HT

Controlled trials are rigid, cumbersome and frequently insensitive. On the credit side they have one key virtue, and that is freedom from bias. Their findings are—or should be—beyond the possibility of distortion by the desires or expectations of either investigators or subjects. This vital asset can easily be lost unless certain basic rules and procedures are observed.

Armitage has drawn attention to what seems to be the original randomised controlled trial, carried out in Holland in 1662:

'Let us take out of the Hospitals, out of the Camps, or from elsewhere, 200, or 500 poor People, that have Fevers, Pleurisies, etc.

'Let us divide them into halves, let us cast lots, that one halfe of them may fall to my share, and the other to yours; we shall see how many Funerals both of us shall have.

'Let the reward of the wager be 300 Florens, deposited on both sides.'

In addition to such admirable features as definitions of cases and their source, and estimation (rather approximately) of sample size, this design also includes the basic means for avoiding bias, namely, randomisation, analysis of all randomised cases, and complete ascertainment of simple and objective critical events. By contrast with most trial reports there is also a frank recognition of the investigator's vested interest in the outcome. Though usually covert (and not usually financial) this must always be assumed to be present; for most investigators hope (privately) for a particular conclusion to their trial. The aim of the design and analysis is to assure the reader that such hopes cannot have influenced the result. The less that he needs to trust the authors, the better.

This article will review some of the ways in which, at each stage of a trial, unwanted biases may creep in.

Randomisation

The purpose of randomisation is to ensure that at the outset the trial groups differ only by chance. Randomisation is basically an offence to reason, because one can argue that judgement and care will

produce a better matching than if the constitution of the groups is left to chance. Thus most doctors basically mistrust randomisation, and when it leads to evident mis-matching they will be quick to point out that 'like is not being compared with like'; and hence, it is implied, the trial is invalidated. This reveals a misunderstanding of what randomisation does and does not achieve. It does not maximise equality of the groups. If, for example, the investigator were to notice that one group were accumulating an excess of high-risk patients, he might in some way interfere with allocations and thereby improve the balance. But the price of that improvement would be that he gained an opportunity to manipulate the groups in such a way as to influence the results; and that possibility cannot be tolerated.

Randomisation is not the best way of producing equality. It is simply the only way of precluding bias in the constitution of the trial groups—the only guarantee of an honestly set up comparison. It is also the only basis for calculating the significance of differences in outcome, and fortunately the *P* value attached to the results takes into account the potentially misleading effects of chance, including initial chance imbalance. The latter may produce error, but not bias; and the risk and limits of such error can be precisely calculated.

There is another reason why randomisation is nearly always preferable to matching, and it arises from our limited ability to predict outcome. If prognosis could be known with certainty, then there would be no need for trials: any departure from the predicted outcome could confidently be attributed to treatment. This happened when streptomycin therapy for tuberculous meningitis was found to be followed by recovery in a disease that previously was predictably fatal. Usually, however, outcome is less certain, and even when known prognostic factors are taken into account it is still impossible to make definite predictions. This implies that even when groups are apparently well matched they may still have unlike prognosis. Matching cannot be guaranteed to achieve its aim.

The problem may be illustrated by the difficulty in assessing the life-saving effect of coronary care units. Attempts have been made to compare mortality in

the same hospital before and after the service was introduced, after matching for known prognostic factors; but this approach would only be acceptable if it had been shown that groups of coronary cases when matched for known prognostic factors always have the same outcome, and this has not been possible. The method of matched comparisons cannot take account of unidentified prognostic factors, whereas the virtue of randomisation is that it copes with all factors, both known and unrecognised. It cannot assure equality but it does exclude bias, in a way that the investigator himself could never achieve.

Stratification

By randomising within sub-groups (strata) it is possible to constrain the limits of imbalance. For example, for each class of age and severity a separate set of allocation envelopes may be prepared, such that within each sequence of (say) eight envelopes there will be an equal number of allocations to each treatment. For practical reasons it is usually impossible to stratify for more than two or at most three variables; and the larger the trial the less the benefit, since there is less risk anyway of chance producing serious imbalance. In a multicentre trial it is usually wise to stratify by centre, thus regulating a major source of heterogeneity.

The purpose of stratification is not to reduce bias but simply to limit the play of chance. However, bias can enter a trial if the investigator has advance knowledge of the group to which the next patient will be allocated, since this may influence his decision on whether or not a particular patient should enter the trial. This happens in trials based on alternating allocation, or when allocation envelopes are translucent. It can also happen in stratified random allocation if the investigator knows that allocation is balanced within each sequence of n envelopes: he can then work out the group to which each n^{th} patient will be allocated. ' n ' must therefore be a closely guarded secret until the end of the trial if it is not double blind.

Unbalanced randomisation

The report on any controlled trial is likely to contain a table setting out the initial comparison of the trial groups. Any notable imbalance will pose two kinds of question.

(1) *Was the randomisation carried out properly? Were any files removed or suppressed after randomisation?* For example, in a trial that is not 'double-blind' the doctor in charge of a particular patient may decide that the allocated treatment is unsuitable, or else perhaps that the patient is too sick for control status to be ethically acceptable; either way he may take the patient out of the trial. He thereby offends a most

fundamental rule: after randomisation, no one can be withdrawn from the trial. Their treatment may sometimes have to be changed ('withdrawal from randomised treatment') or they may be lost to follow-up, but the phrase 'withdrawn from the trial' is inappropriate and should never be used. Allocation envelopes must be numbered, and the final analysis should account for every envelope that has been issued.

If it appears that randomisation somehow went wrong, so that bias crept into the initial constitution of the groups, then there is no remedy and the whole results must be discounted. This happened in a trial by Cochrane in Cardiff, comparing the inpatient treatment of varicose veins by ligature with outpatient treatment by injection and compression. At the conclusion of the trial it was found that the inpatient group seemed to have too few patients, and on enquiry it emerged that a surgical registrar had diverted some of the admissions into another study of his own. Cochrane destroyed all the results and started again.

(2) *Has a chance imbalance influenced the conclusions?* If two randomised groups are contrasted with respect to, say, ten independent characteristics, then clearly it may readily occur that in one of the characteristics they may differ at the 5% level of significance. When this happens, the readers tend to get agitated. What they, and the investigators, should do is to estimate quantitatively the effect of the imbalance on the outcome and conclusions. Suppose, for example, that there are fewer smokers in the control group, then the first step would be to examine the relation of smoking to outcome in the total results. From this can be estimated the contribution of the imbalance to the apparent effect of treatment. Usually it will prove to be slight. If it is not, then a correction procedure, such as Cox's multiple regression model (Cox, 1970) will probably take care of the problem.

Theoretically the precision of a trial can always be improved by correcting the results statistically to allow for initial imbalance, whether small or large; some statisticians will therefore practice such adjustments as a routine. However, a theoretical gain has to be set against a practical loss. The results of a trial are useless unless they influence medical practice. If the readers of the report are baffled by the statistics then they will be less likely to trust the conclusion.

Bias in management

The interpretation of a randomised trial rests on the assumption that any differences in outcome are the result either of chance (whose effects can be estimated) or of the test treatment(s). This assumption

will be invalidated if the groups are not handled equally in respect of every other component of treatment which may influence outcome. In the U.S. National Diet Heart Study subjects in the intervention group were given cholesterol-lowering dietary advice; but when the results were analysed it transpired that a quarter of them had also stopped smoking. Obviously this distorted the trial's primary purpose, which was to test the diet-fat hypothesis of coronary heart disease; the observed result could not be interpreted.

The problem of bias arises (or at least it cannot be excluded) in interpreting any trial that is not double-blind. For example, a trial of coronary artery by-pass surgery necessarily confounds the effects of surgery *per se* with the effects of different drug usage by the two groups. A comparison of the frequency of sudden death in home v hospital care of myocardial infarction confounds the intrinsic merits of the two types of care with the potential arrhythmogenic effects of digoxin and potassium-depleting diuretics, which are used more in hospital. Even in a double-blind trial the effects of the test treatment may influence the indications for using other drugs. β -adrenergic receptor blockers, for example, will tend to decrease the use of other anti-anginal drugs (whose effects on survival are unknown) whilst increasing the use of drugs for controlling cardiac failure.

The purposes of a trial have been usefully classified into two types (Schwartz *et al.*, 1980). The first or 'pragmatic' purpose is to evaluate the effects of a therapeutic decision. For this purpose it is fully appropriate to analyse the patients according to their randomisation ('intention to treat'); and any consequent imbalance in management is not a problem, but simply one component of the package which is being evaluated. Cholesterol-lowering dietary advice might be beneficial, even if the reason were that those so advised also tended to stop smoking; and admission to a coronary care unit might sometimes be worse than staying at home, if only through overuse of arrhythmogenic drugs.

The second purpose of a trial is 'explanatory'—the attempt to explain the actions of a specific therapy. This purpose is present to some degree in all trials, since one always wants to interpret a pragmatic result. Interpretation is hazardous if there were many withdrawals from randomised treatment, or if there was bias in the management of the groups. If differences are recognised then some allowance may be possible. Unrecognised treatment bias is more sinister, for one can never exclude the explanation that has not been considered. One advantage of the double-blind method is that it much increases the chance that groups will be comparably managed. Nevertheless even in a double-blind trial it is good practice to review the comparability of use of the main supplementary or alternative treatments.

Assessing progress and outcome

The benefits of randomisation are vitiated if there is bias in assessing the outcome, and there are many ways in which this can happen.

Dangers of 'before/after' comparisons

Hypertensives tend to improve during the early weeks of treatment, even if the treatment is only tablets of lactose. This was seen in the Medical Research Council Hypertension Trial, where during the first three months in the trial the blood pressure fell in the controls by almost half as much as in those receiving active treatment. This trend presumably reflected habituation to measurement, and possibly also some delayed 'regression to the mean'. Without controls the benefit of treatment would have been greatly over-estimated.

In another trial, designed to study the effect of an autogenous bacterial vaccine on the frequency of colds, volunteers receiving the vaccine reported an average of 1.8 colds in the following winter, compared with 5.6 in the winter before; the experience of the control group was almost identical. Without the control results, everyone concerned might well have concluded that the vaccine was very effective. The explanation of this misleading improvement is a common phenomenon: patients with a chronic or relapsing condition naturally tend to seek help during exacerbations, and these naturally tend to be followed by improvement. Misinterpretation can be avoided either by establishing a stable pre-treatment baseline or by a concurrent control group.

Defining the outcome criteria

The Anturane Re-infarction Trial (1980) was set up to study the possibility of reducing the incidence, in myocardial infarction survivors, of re-infarction and also (so the report claimed) total mortality. The results failed to achieve conventional significance, but the investigators noted with surprise an apparent reduction in sudden deaths, confined to the first 7 months of follow-up. In the light of this finding they defined this as a new category of critical events, for which they claimed statistical significance. They thereby offended two criteria for an honest trial:

- (1) the precise definition of outcome criteria should be written down before any results are known; and
- (2) probability cannot be calculated for hypotheses generated by the results, since these are chosen from among an unknown number of partly interdependent options.

The Edinburgh/Newcastle trials of clofibrate in the treatment of coronary heart disease provided an extreme example of a common fault. The investigators were particularly thorough in their search for

sub-groups of patients in whom the treatment might have been beneficial. A total of eight analyses were 'significant at the 5% level', but it appeared that in at least 183 other instances a *t*-test was not significant. Clearly the number of "successes" could be interpreted only if one also took into account the number of chances; but in practice this is not possible. The thoughtful investigator cannot say in how many different ways he has inspected the results, formally or informally. Moreover many such inspections may be partly interdependent (for example, older patients may also tend to be those with more severe disease), and again no precise allowance is possible.

By changing the definitions after seeing the results the investigators in the Anturane Trial dug another trap for themselves, which they then fell into. When an independent committee reviewed the trial it appeared that the criteria for distinguishing sudden death and fatal re-infarction had been wrongly applied, and the errors were biased in favour of the treatment. When these events were re-assigned correctly, then the *P* value became quite unexciting. The lesson is that the decision as to which patients meet the criteria for a critical event must be made without any knowledge of the treatment group and in advance of analysing the results.

Double-blindness, and its possible failures

Double-blindness is desirable (though not necessarily possible) whenever the knowledge of a patient's treatment might bias the assessment of outcome. Experience with trials of β -adrenoceptor blockers in coronary heart disease illustrates some ways in which it may sometimes fail to provide the expected protection.

(1) In a trial of oxprenolol a patient bit into one of his tablets and found that it tasted sweet. He asked the other patients attending the same clinic to taste their tablets; some proved sweet, others bitter, and the blindness was destroyed. A similar disaster has overtaken trials where tablets were labelled simply A or B, and the identification of one single patient's treatment has unmasked the whole trial. Tablets should be indistinguishable in taste and consistency as well as in appearance, and each patient's supply must be uniquely identified.

(2) β -adrenoceptor blockers often produce characteristic side-effects, such as cold hands and a fall in heart rate, so that double-blindness cannot be relied upon. Sometimes the investigator can be denied access to unmasking information. But in general it is best to assume that a trial of a drug with obvious effects will not be fully double-blind, and potentially biased end-points must be treated with suspicion. Often it is possible to reduce the scope for bias by the clinician; e.g., symptoms may be recorded on a questionnaire completed by the patient without prompting, blood

pressures may be measured by an automatic or blind-reading sphygmomanometer, and laboratory technicians or radiologists may receive no clue as to the treatment.

(3) Where double-blindness is not achieved the investigator may decide to send progress data (biochemistry, electrocardiograms, X-rays, etc.) to be classified by some independent and fully 'blinded' referee. This certainly guards against false-positive verdicts, since the referee makes an unbiased decision on which cases to accept. But it does not protect against false-negative errors, which may arise because some genuine cases were not referred to him; and such non-referral could be biased.

(4) A drug may alter the manifestations by which disease is recognised without necessarily altering the disease itself. For example, β -adrenoceptor blockers modify anginal pain, and this may mean that in a treated patient the doctor is less likely to be alerted to an ischaemic episode and hence less likely to take an electrocardiogram. Such a bias will again not be detected by an external review of electrocardiograms. Wherever such a danger exists, the assessment of outcome should rest primarily on standardised fixed-interval follow-up examinations rather than on investigations initiated at the investigator's discretion.

This problem has been further illustrated by experience in trials of cimetidine in duodenal ulcer, where healing was assessed by endoscopy. In earlier trials endoscopy was undertaken only if a patient continued to experience pain. Naturally there were many fewer endoscopies in cimetidine-treated patients, and many fewer active ulcers were found. In later trials endoscopy has been performed systematically in all patients after a fixed interval, and the incidence of active ulcer is now seen to be reduced much less. The treatment influences symptoms more than disease.

Inappropriate measures of outcome

In every therapeutic assessment there is a danger of accepting a treatment only because it improves the results of the tests. This arises because it is so much easier to use investigations rather than clinical events as a measure of outcome. Patients, however, are concerned only with symptoms and survival, not with the results of tests. It was reported that β -adrenoceptor blockers might alter the dimensions of electrocardiographic Q-waves. Fortunately it seems that this has not actually introduced serious bias into trials of β -adrenoceptor blockade and myocardial infarction; but it points the need, in any trial depending on test results, to question the assumption that treatment does not bias their validity. Further, it emphasises that therapeutic benefit is best related to symptoms and survival rather than to clinical signs and test results.

The problem of interim results

Many trials last a long time, and interim results become available whilst recruitment and/or follow-up still continue. It is best that such information should be withheld from the doctors responsible for care of trial patients. Any apparent trend will not only cause anxiety on behalf of the group faring worse, it may also influence their management, thereby confusing the interpretation of the trial or (if it is not double-blind) biasing subsequent results.

Interim results may also influence the decision on when to stop recruitment or follow-up. By chance alone the groups may at some point in time differ impressively, even though the treatment is in fact useless. By biding his time and stopping when it suits him, the investigator enhances his chance of apparent success. The methods of sequential analysis make allowance for multiple stopping options, but they are not always applicable. Of course, the investigator may decide in advance that he will not inspect the results until the end of the trial, thereby avoiding this problem; but some kind of interim monitoring is essential if the outcome events are serious. So long as the number of inspections is fixed in advance (for example, after every n critical events) then allowance can be made for the effect on significance estimates (Minawa *et al.*, 1981).

In practice this simple remedy cannot always be followed. In large and complex trials there may be unforeseen and worrisome patterns of results: something not specified as a critical event occurs with undue frequency in one group, or else sub-groups of the trial show heterogeneity of effects. The decision to stop may have to be taken in the light of circumstances not specified in advance, implying that the number of potential stopping options has been underestimated. The situation is statistically unsatisfactory but medically it may be unavoidable. Two guidelines are suggested. (1) The trial protocol should state the stopping rules precisely, insofar as they can be foreseen, together with the expected number and timing of occasions on which the results will be so assessed. (2) If in the event the decision to stop has some other basis, then the trial report should make clear what happened, and significance estimates will be correspondingly a little less credible.

Generalisation

Those who conduct a clinical trial tend to forget that their readers are interested, not in what happened to the patients in the trial, but only in whether the findings are a fair guide to what may happen in their own patients. If the report does not permit generalisation, then it is of no value as a guide to clinical practice. To decide whether 'his results apply to my patient' is a matter of judgement more than statistics, but the authors of the report can do much to facilitate the judgement.

(1) Selection of study sample

The study group forms a non-random sample of all cases of the disease. Often the sample is biased in the direction of below-average severity, partly because entry criteria exclude bad and complicated cases, and partly because of reluctance to expose sicker patients to the uncertainties of a trial. (This is why those who conduct trials commonly find—to their dismay—that the event rate is below expectation). Contrariwise it can happen that trials are undertaken in specialist centres to which cases of above-average severity are referred. The large effects of recruitment procedures on prognosis are illustrated by the extraordinarily divergent event rates among control subjects in three recent hypertension trials, even among groups of comparable age, sex and level of blood pressure (Table 1).

Thus it is not sufficient just to report the entry criteria. The reader will also want to know something of the selection process which led to the inclusion of those particular patients, and (if possible) the number of eligible but non-included patients during the trial's period of recruitment.

(2) Proportionate and absolute benefit

An estimate of proportionate reduction in event rate may be more generalisable than a measure of absolute benefit; that is to say, it is less inclined to vary when the treatment is applied to groups of differing severity. This was apparent in the results of the Veterans Administration hypertension trial (Table 2). Nevertheless it is absolute rather than relative

Table 1 Event rates among control subjects in three hypertension trials.

	<i>Deaths per 100 per year</i>		
	<i>Total</i>	<i>CHD</i>	<i>Strokes</i>
Australian National Study	0.3	0.1	0.05
U.S. Hypertension Detection Follow-up Programme	1.5	0.5	0.2
Veterans Administration Trial	3.3	1.7	1.1

Table 2 The results of the Veterans Administration hypertension trial.

<i>Age (years)</i>	<i>Cardiovascular renal complications</i>	<i>% mortality reduction</i>	<i>Lives saved per 100 treated</i>
<50	No	59	6
	Yes	62	14
50+	No	50	15
	Yes	60	29

benefit that is the ultimate guide to a treatment's utility. In this instance, clearly there was far more benefit from blood pressure control to the high-risk than to the low-risk groups. This fact was quite obscured in the original report, which presented only the proportionate estimates of benefit. There is thus something of a paradox here: proportionate estimates are more generalisable (since they vary less between groups of differing risk), but absolute estimates are the proper guide to decisions and efficient application of results. Both should be presented.

(3) Risk/benefit ratios

Although benefits (in absolute terms) tend to be less in low-risk patients, this is unlikely to be true of the risks; and thus the risk/benefit ratio cannot be generalised. Clofibrate, so it appears, carries a lethal risk of the order of one death per 1000 patient-years. The World Health Organization trial evaluated the drug in primary prevention where the chance of absolute benefit was necessarily small, and the outcome was a large excess of deaths in the treatment group; but this could be a misleading estimate of the drug's performance in severe hyperlipidaemia, where the balance of risks and benefits could be different.

Cimetidine provides an example of the opposite error. Trials have established its role in hospital cases of duodenal ulcer, most of whom suffered severe or

prolonged disability. But the long-term effects remain unknown, as to both natural history and risks, and it is not justifiable therefore to use the hospital trials as evidence for cimetidine to be used as a first-line drug in general practice.

(4) Unwarranted extrapolation

The evidence from a trial applies directly only to similar patients, treated with that particular product and dosage for that particular length of time. In chronic diseases it may tell nothing of the effects of really long-term treatment; and it usually gives no guide to the appropriate duration of treatment, nor to the effects of stopping. Answers to these key questions remain speculative.

(5) False-negative outcome

The power of controlled trials is less than is generally supposed, and it is still regrettably common to equate a failure to demonstrate an effect with the positive verdict that the treatment is useless. This danger would be avoided if authors and editors made it a practice in any trial with a seemingly negative outcome to give a confidence interval for the estimated effects. If this embraces what would be regarded as a clinically important result, then the only valid conclusion is that the question remains open and a more powerful trial is needed.

References

- ANTURANE RE-INFARCTION TRIAL RESEARCH GROUP (1980). Sufinpyrazone in the prevention of sudden death after myocardial infarction. *New Engl. J. Med.*, **302**, 250-256.
- COX, D.R. (1970). *Analysis of binary data*. London: Methuen & Co.
- MINAWA, A., FIELDING, J.W.L., JOHNS, B.J. & WATERHOUSE, J.A.H. (1981). Monitoring serial clinical data in groups of patients. *J. inst. Statist.* (in press).
- NATIONAL DIET-HEART STUDY (1968). Final report. *Circulation*, **37**, Suppl 1.
- SCHWARTZ, D., FLAMANT, R. & LELLOUCH, J. (1980). *Clinical trials*. London: Academic Press.
- VETERANS ADMINISTRATION COOPERATIVE STUDY GROUP. (1972). Effects of treatment on morbidity in hypertension: III Influence of age, diastolic pressure and prior cardiovascular disease; further analysis of side-effects. *Circulation*, **45**, 991-1004.