Table 1 Percentage binding of chloroquine to plasma protein and 4% bovine serum albumin (BSA) in the presence and absence of aspirin. Values are given as mean $\pm$ s.e. mean of 4 determinations.

Concentration of chloroquine ($\mu\text{g/ml}$)	Concentration of aspirin ($\mu\text{g/ml}$)	Bound drug (%)	
		Plasma	4% BSA
2	—	46.8 $\pm$ 0.8	30.5 $\pm$ 0.7
2	2	46.2 $\pm$ 0.3	29.0 $\pm$ 0.5
4	—	44.5 $\pm$ 0.6	25.1 $\pm$ 0.4
4	4	42.6 $\pm$ 0.8	25.6 $\pm$ 0.7
6	—	43.1 $\pm$ 0.3	24.2 $\pm$ 0.4
6	6	42.3 $\pm$ 1.0	24.3 $\pm$ 0.3
8	—	43.1 $\pm$ 0.4	24.3 $\pm$ 0.4
8	8	42.7 $\pm$ 0.5	24.0 $\pm$ 0.5

fraction of plasma. Other observations (Adelusi & Salako, unpublished observations) have also shown that the aspirin did not affect the rate of absorption or the plasma level of chloroquine in children, and with the present result it is clear that adverse interaction between aspirin and chloroquine is unlikely to affect the total or free plasma concentration of chloroquine in man.

This investigation received support from the UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases.

S. A. ADELUSI & L. A. SALAKO
*Department of Pharmacology & Therapeutics,
University of Ibadan, Ibadan, Nigeria*

Received August 21, 1981

References

- AARONS, L., CLIFTON, P., FLEMING, G. & ROWLAND, M. (1980). Aspirin binding and the effect of albumin on spontaneous and enzyme-catalysed hydrolysis. *J. Pharm. Pharmac.*, **32**, 537–543.
- ADELUSI, S.A. & SALAKO, L.A. (1980). Improved fluorimetric assay of chloroquine in biological samples. *J. Pharm. Pharmac.*, **32**, 711–712.
- BUCHANAN, N. & VAN DER WALT, L.A. (1977). The binding of chloroquine to normal and kwashiorkor serum. *Am. J. trop. Med. Hyg.*, **26**, 1025–1027.
- KLOTZ, I.M., WALTZ, O., WALKER, F.M. & PIVAN, R.B. (1946). The binding of organic ions by proteins. *J. Am. chem. Soc.*, **68**, 1486–1490.
- McCHESNEY, E.W., BANKS, W.F., Jr. & McAULIFF, J.P. (1962). Laboratory studies of the 4-aminoquinoline antimalarials; II. Plasma levels of chloroquine and hydroxychloroquine in man after various oral dosage regimens. *Antibiot. Chemother.*, **12**, 583–594.
- McCHESNEY, E.W., FASCO, M.J. & BANKS, W.F., Jr. (1967). The metabolism of chloroquine in man during and after repeated oral dosage. *J. Pharmac. exp. Ther.*, **158**, 323–331.
- McCHESNEY, E.W. & McAULIFF, J.P. (1961). Laboratory studies of the 4-aminoquinoline antimalarials: I. Some biochemical characteristics of chloroquine, hydroxychloroquine and SN-7718. *Antibiot. chemother.*, **11**, 121–129.
- PARKER, F. & IRVIN, J. (1952). The interaction of chloroquine with the albumin of the bovine plasma. *J. biol. Chem.*, **199**, 897–901.
- RUBIN, M., ZVAIFLER, N., BERNSTEIN, H. & MANSOUR, A. (1965). Chloroquine toxicity. *Proceedings of the Second International Congress of Pharmacology*, pp 467–486. Oxford: Pergamon Press.

FREE SODIUM VALPROATE MONITORING

Plasma anticonvulsant monitoring has contributed significantly to the management of epileptic patients. In most instances, only total drug concentration is measured, both for technical reasons and because it is assumed that the total value reflects the free (unbound—pharmacologically active) drug concentration. Considerable effort has been expended in attempting to define the therapeutic range of serum sodium valproate. Results to date have been dis-

appointing, for although a range of 300–700 $\mu\text{mol/l}$ has been proposed (Schobben *et al.*, 1975), there is little relationship between serum drug concentration and seizure control. Evidence exists that the free serum sodium valproate fraction shows some considerable variation during a single dosing interval at steady state (Bowdle *et al.*, 1980) and that there is marked intersubject variation in the free:total concentration ratio (Patel *et al.*, 1980). Due to the poor

prediction value of serum sodium valproate measurements, it has been proposed that monitoring of free, rather than total, values may be of use (Levy, 1980). The purpose of the present study was to examine the relationship between free and total sodium valproate levels.

Fifty-nine random serum samples obtained from 30 healthy epileptic patients (aged between 3 and 56 years, four of whom were children), referred for routine sodium valproate assay were studied for free and total serum sodium valproate concentrations. All the patients had normal serum albumin concentrations. Seven patients were on monotherapy and the remaining 23 received at least one other anticonvulsant.

Four persons were studied in greater detail, with hourly blood samples collected from an indwelling heparinised venous catheter over a 24 h period. In this group, patient 3 received carbamazepine and phenytoin and patients 1 and 4, carbamazepine as additional therapy.

Total serum sodium valproate was measured by gas liquid chromatography (Wood *et al.*, 1977) as was free sodium valproate following three hours of equilibrium dialysis at room temperature (Buchanan & Ponniah, 1981). Serum free fatty acid was not measured.

All the 59 samples studied had total serum sodium valproate concentrations within the proposed therapeutic range. The variable relationship between free and total valproate concentrations for these samples ($r = 0.68$) is shown in Figure 1. When 95% tolerance limits were calculated, free drug levels varied between 3.4 and 60 $\mu\text{mol/l}$ at 300 $\mu\text{mol/l}$ total drug (17.6 fold), and between 39.7 and 98.7 at 600 $\mu\text{mol/l}$ (2.5 fold). Both free and total drug levels of the four children included in the study fell within one standard deviation of the mean for the whole group. In Figure 2, total and free valproate concentrations are shown for a patient in steady state during two dosing intervals. Considerable fluctuations occur with the peak concentration being 3.2 times higher than the trough concentration for total drug but 6.4 times higher for free drug. With increasing total valproate concentrations the free fraction increases in a non-linear fashion from 9.2% to 18.3%. Four patients (including patient 4 depicted in Figure 2) were studied in greater detail, with regard to the relationship between free and total valproate at different concentrations of the latter. Two of the patients exhibited a linear relationship between free and total drug concentrations (Figure 3) whilst in the other two patients, a non-linear (saturable) relationship was observed.

Sodium valproate is an anticonvulsant for which effective clinical use is not enhanced by the measurement of serum drug levels. Previous studies have suggested that the measurement of free valproate may be beneficial in establishing a therapeutic range

Figure 1 Free v total valproate concentrations in 30 epileptic patients (59 plasma samples). Correlation coefficient $r = 0.68$ (—) and 95% tolerance limits (---)

for this compound (Levy, 1980; Wood *et al.*, 1977). The present study set out to ascertain, in further detail, the relationship between total and free sodium valproate concentrations. The results have shown that whilst the free sodium valproate concentration follows the total drug concentration, there is considerable inter-individual variation in the degree of protein binding of this drug. This has been observed previously and may be partly related to competitive binding for free fatty acids (Patel & Levy, 1979) which probably accounts for about 25% of the variability in the free fraction (Bowdle *et al.*, 1980). Although sodium valproate may displace other anticonvulsants such as phenytoin, from their binding sites, the reverse does not seem to apply and does not account for the observed variability in the free fraction. The present study has shown the general pharmacological assumption that total drug levels reflect free levels, to be invalid for sodium valproate. However, one encouraging observation is that there is a linear relationship between cerebrospinal fluid and free plasma

Figure 2 Plasma level v time profile of total (●) and free (○) valproate in an epileptic patient between two dosing intervals.

Figure 3 Free v total valproate concentrations in four epileptic patients. Patient 1 (○), 2 (▲), 3 (●), 4 (□).

valproate levels (Levy *et al.*, 1980). This would suggest that despite the potential difficulties in interpreting free valproate values, a prospective study relating free valproate concentrations to clinical control is necessary. At present the indication for valproate therapeutic drug monitoring using total drug values is unproven. Until a clinical efficacy versus free drug level study is performed and depending upon the results, the same conclusion must apply to free valproate monitoring.

E. JAGER ROMAN, P. PONNIAH, J. BAIRD LAMBERT & N. BUCHANAN
Paediatric Pharmacology Unit, Department of Paediatrics, Westmead Centre, Westmead, N.S.W. 2145, Australia.

Received September 7, 1981

References

- BOWDLE, T.A., PATEL, I.H., LEVY, R.H. & WILENSKY, A.J. (1980). Valproic acid dosage and plasma protein binding and clearance. *Clin. Pharmac. Ther.*, **28**, 486-492.
- BUCHANAN, N. & PONNIAH, P. (1981). Value of measuring salivary and free sodium valproate concentrations. *Med. J. Aust.*, **1**, 539-540.
- DAHLQVIST, R., BORGA, O., RANE, A., WALSH, Z. & SJOOVIST, F. (1979). Decreased plasma protein binding of phenytoin in patients on valproic acid. *Br. J. clin. Pharmac.*, **8**, 547-552.
- LEVY, R.H. (1980). Monitoring of free valproic acid levels. *Ther. Drug Monitoring*, **2**, 199-201.

- LEVY, R.H., LOCKARD, J.S. & LUDWICK, B.T. (1980). Plasma protein binding and C.S.F. concentration of valproic acid in the monkey. Abstracts, *Apha Acad. Pharm. Sci.*, **24**, 10, 79.
- PATEL, I.H. & LEVY, R.H. (1979). Valproic acid binding to human serum albumin and determination of free fraction in the presence of anticonvulsants and free fatty acids. *Epilepsia*, **20**, 85–90.
- PATEL, I.H., LEVY, R.H., VENKATARAMANAN, R., VISWANATHAN, C.T. & MORETTI-OJEMANN, L. (1980). Diurnal variation in protein binding of valproic acid and phenytoin and the role of free fatty acids. *Clin. Pharmac. Ther.*, **27**, 277 (abstr.).
- SCHOBEN, E., VAN DER KLEIJN, E. & GABREELS, F.M.M. (1975). Pharmacokinetics of di-N-propylacetate in epileptic patients. *Eur. J. clin. Pharmac.*, **8**, 97–105.
- WOOD, M.H., SAMPSON, D.C. & HENSLEY, W.J. (1977). The estimation of plasma valproate by gas liquid chromatography. *Clin. Chim. Acta*, **77**, 343–347.

A COMPARISON OF THE BIOAVAILABILITY OF TWO BRANDS OF PHENYTOIN

Although the bioavailability of different brands of phenytoin available in the United Kingdom has not been widely studied, there have been some suggestions (Richens, 1976; Stewart *et al.*, 1975) that different preparations could produce different steady-state plasma concentrations. In this study tablets obtained from Boots and capsules from Parke-Davis have been compared. Cox, Kerfoot and Macarthy also manufacture phenytoin tablets, available in the U.K., but these were not tested in this study.

Patients whose epilepsy was well controlled on a given dose of phenytoin, with or without another anticonvulsant were entered into the trial. Twenty-two patients, 13 men and 9 women, ages ranging between 17 and 69 years entered the cross-over trial. Patients were randomly allocated initially to either Boots tablets or Parke-Davis capsules of phenytoin (Epanutin) and then crossed over to the other. The aims of the trial were explained to the patients and they were asked to take their tablets regularly. To check patient compliance a return tablet count was performed. At the end of a 2 week period blood samples for phenytoin concentrations and other anticonvulsant concentrations were taken. The time of day at which blood samples were collected was constant for each patient. The number of fits sustained and any side-effects occurring during that fortnight were recorded. The patients then crossed over and received the other brand of phenytoin for 2 weeks and a similar procedure was followed. At the end of the 4

weeks patients were asked to state which preparation they preferred.

The analysis for phenytoin developed by one of us (D.B.) was performed by adding 50 μ l of an internal standard (50 mg methyl primidone/l of methanol) to 0.5 ml of buffered plasma (pH 5.4). The drugs were extracted into dichloroethane which was separated and evaporated to dryness. After reconstitution with methanol the residue was examined by gas-liquid chromatography using a nitrogen specific detector. Each batch of analyses was quality controlled internally, and long term performance assessed by participation in an external quality control scheme, organised at St Bartholomew's Hospital, London.

The results are shown in Table 1. Using a paired *t*-test there was no significant difference between the plasma phenytoin concentrations obtained with either the Boots or Parke-Davis preparations ($P = 0.3$). The tablet count showed excellent compliance. In the majority, the tablet count was exact and the worst patient was only 4 tablets out. This good result is not surprising as the patients chosen were already known to be complying and the trial was a short one.

This comparison was repeated for the 10 patients taking phenytoin alone, and for the 12 taking phenytoin with another anticonvulsant. In each of these two groups there was no significant difference between phenytoin levels for either preparation ($t = 1.5$ with 9 degrees of freedom and $P = 0.17$; $t = 16$ with 11 degrees of freedom and $P = 0.88$ respectively).

Table 1 Comparison of phenytoin levels while on Parke-Davis capsules or Boots tablets.

	Total number of patients	Mean serum phenytoin level (mg/l)	s.d.
Parke-Davis	22	7.52	4.40
Boots	22	6.82	4.59
Paired difference		0.68	2.98 $P = 0.30$