Commentary: Patient's experience

Mrs Patel with the support of Roxy Senior

Department of
Cardiovascular
Medicine,
Northwick Park
Hospital, Harrow,
Middlesex HA1 3UJ
Roxy Senior
consultant
cardiologist
roxy.senior@
virgin.net

Initially my husband and I thought that the symptoms I was experiencing (pain in my chest and right hand) were down to muscular pain and due to me carrying our baby around.¹ This pain gradually increased and got unbearable, causing me to sit down often when out and about. I took paracetamol on our doctor's advice, but this did not seem to help. After I had been to accident and emergency and referred to a cardiologist, it became apparent that I had heart disease.

At the time of the discovery we were shattered to hear the news, and more so because of my age. However, we were relieved that the problem was discovered before

I had a heart attack. We were also glad that the doctors were straight with us and told us the seriousness of the situation as soon as they found out. The cardiologist was very helpful and explained things to us at every stage, which was also good. Since the angioplasty, I seem to be recovering well and am leading a normal life thanks to the prompt treatment I received.

Competing interests: None declared.

1 Dwivedi G, Hayat SA, Violaris AG, Senior R. A 28 year old postpartum woman with right sided chest discomfort: case outcome. *BMJ* 2006;332:643.

Commentary: Cardiological management

Peter A Henriksen, Nicholas A Boon

Department of
Cardiology, New
Royal Infirmary of
Edinburgh,
Edinburgh
EH16 4SA
Peter Henriksen
registrar in cardiology
Nicholas Boon
consultant
cardiologist

Correspondence to:
P Henriksen
phenriksen@
blueyonder.co.uk

Mrs Patel's case shows the challenge of assessing young patients with possible cardiac chest pain.¹ The location of her chest discomfort may be considered atypical, but Marin Marinovic correctly notes that the close relation with exertion raises the possibility that it is cardiac in origin.² Overlap in the location of symptoms between patients presenting with cardiac and non-cardiac chest pain makes discrimination difficult using this factor alone.³

Several respondents on *bmj.com* flag the possibilities of pulmonary embolism or pneumothorax as causes of chest pain in the postnatal period.² However, the absence of dyspnoea and the non-pleuritic nature of the pain points away from these diagnoses. Acute coronary syndromes do occur infrequently during pregnancy when a prothrombotic state and changes in haemodynamic forces can lead to occlusive coronary thrombosis or arterial dissection respectively. Mrs Patel's presentation seven months after giving birth was not so dramatic, perhaps indicating a subacute process.

Investigation and further management

Postmortem studies show that atherosclerotic coronary disease is common in young adults. Factors responsible for driving plaque growth and the development of an obstructive coronary stenosis leading to angina are incompletely understood. Risk assessment using variables such as smoking can predict events in populations but not individuals. Mrs Patel does have risk factors.

Stephen Head notes correctly that her maintenance dose of thyroxine is low.² We learn subsequently that she is biochemically hypothyroid and has raised low density lipoprotein cholesterol concentrations. She does not have features of severe hypothyroidism, and delaying the increase in her thyroxine replacement until after revascularisation would arguably have avoided the small risk of precipitating further ischaemia.

A troponin measurement might also have been helpful in view of her particularly severe episode of pain. Although normal troponin concentrations do not exclude serious disease in patients with exertional symptoms, it is a marker of intracoronary thrombus (noted on Mrs Patel's angiogram), the likely cause of an acute deterioration in a patient with hitherto stable symptoms.⁴

Her resting electrocardiogram shows abnormalities. Although the changes may reflect her weight, they should in the context of exertional chest pain prompt further investigation for an underlying cardiac problem. She had echocardiography to look for evidence of infarction or an alternative explanation such as myocarditis or hypertrophic cardiomyopathy for the symptoms. The predictive power of treadmill testing depends on disease prevalence in the population tested, according to bayesian principles. This is of particular relevance in that predictive accuracy is reduced for young women.⁵ Abnormal results on resting electrocardiography further complicate interpretation of the test. The development of symptoms limits performance on the treadmill,

Stress echocardiography isn't widely available in the UK