Primary care trusts to continue direct provision of services

The UK government has insisted that reconfigured primary care trusts (PCTs) will be free to continue to provide as well as to commission a range of services under its latest reforms. The announcement is part of its response, published last week, to a scathing health select committee report that attacked the reforms as "ill judged" and the policy of contracting out provider services as "incoherent" (*BMJ* 2006;332:72, 14 Jan).

Ministers originally said that the PCTs would have to shed all provided services, such as district nursing and family planning, and concentrate solely on commissioning. They later softened this line, saying trusts could continue to run services in certain circumstances. But it was widely believed that trusts would have to disengage sooner or later.

The government has now said, however, that it will support PCTs "whether or not they chose to divest, provided that the approach they are taking is genuinely best for local patient care."

Andrew Cole *London*

The government's response to the health committee's report on changes to primary care trusts is available at www.dh.gov.uk.

Russian woman may lose grandson conceived from dead son's frozen sperm

A Russian woman who had a grandson two years after her only son died by using his frozen sperm and a surrogate mother faces losing the child just three months after he was born. Headmistress Ekaterina Zakarova, aged 55, lost her law student son Andrei two years ago but managed to persuade doctors, after his death, to use his frozen sperm to fertilise a donor egg that was then implanted into a surrogate mother. Andrei had

had his sperm frozen before having treatment for the cancer that eventually killed him.

But now Russian officials say that because Andrei died two years ago he cannot be registered as the father, and, because the egg donor was anonymous, the baby also does not have a mother. Consequently the baby has no legal parents, does not officially exist, and cannot have a birth certificate.

The Civil Registry Office says that Mrs Zakarova has no claim on the boy, and as she is too old to adopt him it wants to take him away from her and place him in an orphanage. The office has now taken her to court.

Michael Leidig *Vienna*

GPs' leader sets conditions for electronic care records

A long standing disagreement between GPs and leaders of the NHS's information technology programme may be moving towards rapprochement. The dispute is over whether patients should have to opt in to the new systems of electronic health records being created in England, or have the option to keep their medical records off the system if they do not wish them to be shared across the NHS.

Officials at Connecting for Health, the agency building the new electronic records service, argue for "opt out," but the council of the Royal College of General Practitioners last month recommended "opt in" as essential to creating explicit consent.

The college indicated that it would change its position, however, if the NHS tackled areas of concern. Mayur Lakhani, the college's chairman, said that opt in posed "huge logistical difficulties."

To accept opt out, however, Dr Lakhani has asked for action on three points – clarification about which data will go on to the electronic record, action to clean up data going into the system, and "clearly understood access rules" covering the system's personal demographics service. (See p 656.)

Michael Cross *London*

Poorest children under 10 are 40% more likely to die than the richest

Roger Dobson *Abergavenny*

Children in the United Kingdom born into the lowest socioeconomic group have a 40% increased chance of dying in the first 10 years of life compared with children born into the highest group, a new study has shown.

The study was published in the same week that the UK government had to admit that it had narrowly failed to meet its target of lifting 400 000 children out of poverty by 2004-5. It also showed that hospital admission rates for 14 groups of diseases were also related to social class (*Paediatric and Perinatal Epidemiology* 2006;20:14-23).

"We found a clear association between low social class at the time of birth and increased mortality and morbidity during the first 10 years," said the lead author, Stavros Petrou, and colleagues. "Children born into a social class V household [in which the main breadwinner is unskilled] were found to have a substantially increased risk of mortality during the 10-year time horizon of the study, compared with children born into a social class I household."

The authors looked at the association between social class of the head of household, and mortality and morbidity in the first 10 years of life, using data from a cohort of 117 212 children born to women in Oxfordshire or West Berkshire between 1979 and 1988. The study is thought to

be one of the largest studies into social class inequalities in childhood mortality and morbidity. In total, 93 657 children were included in the analyses.

In the first 10 years, mortality was lowest in children born into a socioeconomic group II household (with the main breadwinner in a managerial or technical role), at 7.03 deaths per 1000 births, and greatest in children born into a social class V household, at 13.07 per 1000 births.

The adjusted odds of mortality for each drop in social class category was estimated at 1.08 (95% confidence interval 1.03 to 1.14): "This translates into a 40% increased odds of mortality, on average, for a child born into a social class V household compared with a child born into a social class I household," the authors, from the University of Oxford, said.

The results also show that for each drop in social class in the first 10 years of life, the mean number of hospital admissions increased by 5%.

The researchers say that the results support the existence of a social class gradient. "Because the gradient is seen across such a wide range of outcomes, the effects of low social class are unlikely to be mediated simply through a few recognised biological, behavioural, and environmental factors, such as low birth weight, parental smoking, or exposure to infection."