matriculation qualifications and/or experience"; in 1985, "No considerations other than academic attainment and aptitude are set as entrance requirements."

When quotation marks are used by your correspondent to suggest a verifiable source for some emotional remarks one doubts if the real concern lies with factual data and truth.

H P WASSERMANN

Faculty of Medicine,
University of Stellenbosch,
Stellenbosch,
RSA

Severe extravasation injury

SIR,—We agree with Dr D A R Bard and colleagues (25 May, p 1579) that prevention rather than treatment of extravasation injuries is of paramount importance, and to this end our special care baby unit has a policy of ensuring that infusion sites are visible and are inspected each hour. Despite this policy we see extravasation injuries, and in neonatal practice, with the presently available infusion pumps, this will probably continue to be the case.¹

We have developed a simple technique which seems to limit scarring and reduce the need for

to free the growing epithelium from the central coagulum. This process is continued until epithelialisation is complete. The infant's mother can be shown how to care to the wound in this way and can continue with it at home if necessary. Very little discomfort is caused, provided the epithelial edge is kept separated from the coagulum.

Using this regimen we have seen fewer severe extravasation injuries and we have found that healing is rapid and a good cosmetic and functional result is obtained even in large lesions, which might otherwise be considered for grafting (figure). Rapid re-epithelialisation is probably encouraged because the regular freeing of the growing edge removes the slowing effect of producing collagenoses to burrow beneath the overlying coagulum. Scarring is probably reduced because the wound contracting effect of myofibroblasts is disrupted.

As Dr Bard points out, problems of finding matched controls makes clinical trials in this area very difficult. We recommend this technique in neonatal practice because it is simple, reduces scarring, and is not prone to further complications.

D C DAVIDSON
J GILBERT

Special Care Baby Unit,
Fazakerley Hospital,
Liverpool L9 7AL

¹ Wales JKH, Hall MA, Walton P. Neonatal infusion site complications. *British Journal of Parenteral Therapy* 1985;6:8-5.

Potential hazard of clotting during blood transfusion using a blood warming pack

SIR,—Drs J Blagdon and T Gibson gave an account of an incident of coagulation of whole blood when mixed with compound sodium lactate intravenous infusion (Hartmann's solution) when a blood warmer was in use (18 May, p 1475). We want to report a similar phenomenon which occurred when fresh frozen plasma was preceded and followed by Hartmann's solution through a blood warming coil.

The patient, who was undergoing surgery for an aortic aneurysm, had received normal saline followed by four units of whole blood (CPDA) using an Avon A80 blood warming coil at 37°C. A Pall filter was in use, which was changed after the first two units of blood. About 200 ml of Hartmann's solution was then given, followed by a unit of fresh frozen plasma, after which the Hartmann's solution was restarted. At this point large fibrin clots were seen in the tubing of the giving set. The infusion was stopped immediately and the cannula withdrawn from the vein. No additional drugs had been given to the patient via the giving set. Some investigations were performed to examine the nature of the problem.

The table shows the capacity of Hartmann's solution, at various dilutions, to induce clotting of stored blood and fresh frozen plasma at 37°C and at room temperature. The fact that the clotting times are shorter in the presence of the higher proportion of Hartmann's solution can be accounted for by the greater amount of calcium present. At a ratio of 1:10 the clotting time of the fresh frozen plasma approaches that of a sample of time expired stored blood under the same conditions.

In view of our experience and the incident

Clotting times (minutes) of various dilutions of blood and fresh frozen plasma (FFP) with Hartmann's solution

FFP or blood: Hartmann's:	37°C				Room temperature			
	1:1	1:2	1:4	1:10	1:1	1:2	1:4	1:10
Fresh frozen plasma	>30	>30	14	5	>30	>30	30	18
Blood (time expired, 28 days)	6	4	3	3	30	30	17	16

Prothrombin time of blood used for investigations 22 s; prothrombin time of fresh frozen plasma 12 s.

reported by Drs Blagdon and Gibson it is clear that neither blood nor fresh frozen plasma should be preceded or followed by any solution containing calcium through the same giving set, especially when using a blood warming coil.

L PARAPIA
F E EDGAR
T PIGHILLS
J COX

Bradford Royal Infirmary,
Bradford BD9 6RJ

Thiocyanate and nicotine half lives in plasma

SIR,—Dr Burckhard Junge's paper (6 July, p 22) was a revelation to me and no doubt to many others also. Thiocyanate concentrations in body fluids have been widely used as a quantitative measure of the intake of tobacco smoke in active and passive smokers and for the validation of self reported abstinence from smoking, although they are probably less useful than other markers such as nicotine, cotinine, and carbon monoxide concentrations.¹ Dr Junge has pointed out that the generally accepted value of two weeks for the half life of thiocyanate in plasma or serum is based on findings in a single subject published 13 years ago² and that the true value is probably about six days. However, he also states, without citing a reference, that the half life of nicotine is 30 minutes. He may have obtained this value from two sources which failed to differentiate the initial (distributional) phase from the terminal (elimination) phase.^{3,4} The half life of nicotine in plasma is biexponential, with an initial phase of about 9 minutes⁵ and a terminal half life of 2 to 2.2 hours.⁶ I write lest Dr Junge's paper in destroying one myth may inadvertently help to prolong another.

M A H RUSSELL

Addiction Research Unit,
Institute of Psychiatry,
London SE5 8AF

- 1 Jarvis MJ, Tunstall-Pedoe H, Feyerabend C, Vesey C, Saloojee Y. Biochemical markers of smoke absorption and self-reported exposure to passive smoking. *J Epidemiol Community Health* 1984;38:335-9.
- 2 Pettigrew AR, Fell GS. Simplified colorimetric determination of thiocyanate in biological fluids, and its application to investigation of the toxic amblyopias. *Clin Chem* 1972;18:996-1000.
- 3 Isaac PF, Rand MJ. Cigarette smoking and plasma levels of nicotine. *Nature* 1972;236:308-10.
- 4 Armitage AK, Dollery CT, George CF, Houseman TH, Lewis PJ, Turner DM. Absorption and metabolism of nicotine from cigarettes. *Br Med J* 1975;iv:313-6.
- 5 Feyerabend C, Ings RMJ, Russell MAH. Nicotine pharmacokinetics and its application to intake from smoking. *Br J Clin Pharmacol* 1985;19:239-47.
- 6 Benowitz NL, Jacob P, Jones RT, Rosenberg J. Interindividual variability in the metabolism and cardiovascular effects of nicotine in man. *J Pharmacol Exp Ther* 1982;221:368-72.

Benign skin tumours

SIR,—I would agree with Dr Stephen K Jones and his colleagues (1 June, p 1747) that the treatment of giant pigmented naevi is controversial and apologise for adding to the confusion.

The exact definition of a giant naevus is imprecise and varies from a lesion covering at least 30% of the body surface area to one which can be completely excised and closed primarily in a single operation. Histologically these lesions can also vary. In some the naevus cells are superficial but in

Top: extravasation injury in a preterm neonate; bottom: the same injury 12 weeks later.

grafting. If the tissues are tense the limb is gently massaged until they are soft and a good capillary flush is seen in all potentially viable tissue. This process is repeated every 15 minutes for two to three hours and subsequently every two to three hours until the tissues remain soft. Ensuring that the tissues remain soft encourages good perfusion of any potentially ischaemic tissue. No dressings are used. The necrotic skin is allowed to dry and demarcate. Once (sometimes twice) each day the skin around the wound is stretched sufficiently just

others the lesion is far more hamartomatous and the naevus cells lie deeply in the dermis, even possibly in the underlying musculature.¹ There is little disagreement that these giant naevi can undergo malignant change, but the true incidence of giant naevi is not known in the United Kingdom, though it is less than 1 case per 1000 live births.² Malignant melanoma in the United Kingdom under the age of 15 years is extremely rare, being about two to three cases per year over the last 18 years, but this may be an under-estimation (A Qabu, personal communication). Unfortunately the prognosis in childhood of this cancer is extremely grave.

Controversies surrounding treatment depend on two aspects. The first is over prophylaxis against malignant change. Dr Jones and his colleagues think that prophylactic excision prevents the future development of malignant melanoma. There are, however, reports of cases of melanoma occurring despite the fact that the lesion had been previously excised and skin grafted. This may reflect the inadequacy of the initial excision, which if undertaken should be aided by frozen section reports to make sure that all the naevus cells are being removed. There are several cases in published reports of non-epidermal origin of melanoma associated with giant naevi.¹ One must also take into account that there is a morbidity, even mortality, from repeated anaesthesia and excision of large areas of skin with skin grafting. Against this background some surgeons, including myself, take a very conservative view on the excision of giant naevi as a prophylaxis against malignant change, and I do not think anyone with the available facts can be dogmatic.

The second consideration is purely cosmetic. I agree that shaving or dermabrasion of these lesions is unlikely to be a prophylaxis against malignant change, but as a compromise treatment it may well have something to offer purely on cosmetic grounds, as the pigmentation is usually reduced and the hair growth can also be lessened. In most cases, excision and skin grafting leaves such a bad cosmetic result that it should not be used as an argument for treatment.

D M DAVIES

Royal Postgraduate Medical School,
London W12 0HS

1 Rhodes AR, Wood WC, Sober AJ, Mihim MC. Non epidermal origin of malignant melanoma associated with a giant congenital neurocutaneous naevus. *Plastic and Reconstructive Surgery* 1981;69:782-90.

2 Walton RG, Jacobs AH, Cox AJ. Pigmented lesions in newborn infants. *Br J Dermatol* 1976;95:389.

Fine needle aspiration cytology in isolated thyroid swellings

SIR,—Mr Hilal M Al-Sayer and others (18 May, p 1490) correctly point out that the predictive accuracy of cytology can be reliably established only by "blind" comparison with the histological diagnosis, and they present the results of decision analysis performed on data from such a study. They then, however, defy their own argument by presenting results from a second study for which not all patients had a histological diagnosis. The assumption made in the analysis of this second study—that the 18 patients who had no histological diagnosis had benign disease—not only makes the results of less weight, as the authors state, but also invalidates the analysis. The only assumption which could correctly be made is that the false negative fraction was the same as for the previous study and thus that 14% of the 18 patients had malignant disease. The claim that the false negative fraction fell from 14% in the first year to

4.6% in the second is not substantiated because it is incorrect to discuss a false negative fraction when there are no data on final diagnosis for 18 out of 62 patients.

If, as the authors correctly suggest, clinical judgment is always an overriding arbiter of the indications for operation, the first year study should be repeated with the criteria for positive classification being changed from "suspicious or malignant aspiration cytology" to "suspicious or malignant cytology or positive clinical indications." A study such as this might well show a reduced false negative fraction.

The false negative fraction so calculated, if still greater than zero, would then be the correct figure to use in an assessment on whether the monetary and human cost savings (due to reduced bed occupancy and fewer operations) which could accrue from the suggested method of use of aspiration cytology would be sufficiently great to justify the consequent risk of mortality to the patients with malignant disease who would not be detected.

SUSAN V ELLAM
M N MAISEY

Department of Radiological Sciences,
Guy's Hospital Medical and Dental Schools,
London SE1 9RT

Treatment of primary breast cancer in Great Britain

SIR,—The paper by Mr Jean-Claude Gazet and others (15 June, p 1793) failed to address the question which the authors set—that is, whether current clinical trials of treatment had in fact influenced surgical practice in the management of primary breast cancer. It is hardly surprising that there is a lack of consensus among surgeons about the best method of managing primary breast carcinoma, particularly in view of the authors' own admission that, at least as regards survival, there seems to be no difference between conservative surgery and mastectomy.

If the questionnaire had related to, say, the most appropriate management of chronic duodenal ulcer or rectal carcinoma, or even the best method of inguinal hernia repair, it would have been equally impossible to achieve surgical agreement. It is surely excessively naive to assume that there is necessarily a single best treatment for all patients with primary breast cancer. It is clear that a number of treatments, varying from lumpectomy and radiotherapy to some form of mastectomy, offer equal control of local disease and have similar survival rates provided the patient is appropriately selected.

The problem, of course, is how should the surgeon select the most appropriate treatment for a particular patient. With respect to local control, there are a number of trials currently in progress which should define the precise indications for lumpectomy and radiotherapy and for mastectomy. Current trials of adjuvant treatment—that is, hormone treatment and chemotherapy—are also answering the question of appropriate systemic therapy. Combination chemotherapy has now been shown to have beneficial effects on survival in premenopausal patients with metastases in the lymph nodes, and tamoxifen has shown a survival benefit in premenopausal patients with affected nodes and all postmenopausal patients. The paper merely reflects the current state of play in the UK and tells us no more than we know already.

E A BENSON

General Infirmary,
Leeds LS1 3EX

A unit management team's second year of business

SIR,—I sympathise with Dr Roger Morgan (15 June, p 1828) on the amount of paper circulated to his hospital management team which he has to digest, but I was surprised that one point in his original paper was not challenged: he appeared to make a virtue of members submitting no written material at all. Items presented verbally, however, often waste the time of teams, which would have been saved by a brief, clear report outlining the issue, allowing members to consider it in advance. It is also a good discipline for the presenter of an item to set out his case clearly on paper.

In my experience, which has included membership of a hospital management committee and of two district management teams, excessive circulation of paper occurs in the health service for three reasons. Firstly, a lack of trust between and within disciplines leads to individuals demanding to see the full version of every document and being unwilling to accept a précis; secondly, little attention is given to good writing in some of the professions concerned; and, thirdly, as Mr Kenneth Clarke recently said, there are too many second line managers in first line managerial posts.

GEOFFREY PATEY

Colchester, Essex CO6 2AR

Junior hospital doctors' hours of work campaign

SIR,—No doubt ASTMS/MPU will be defending its campaign and informing people about whom it represents, but Mr Keith Porter's letter (29 June, p 1989) raises some important points.

He states that he and his colleagues (How many? Representing whom?) "feel" that rotas less strenuous than 1 in 3 will not provide adequate clinical experience. Does he mean 1 in 3 for a busy house officer gaining experience as an all night clerking machine; 1 in 3 for a registrar in a less strenuous specialty, on call from home; or does he mean with prospective cover (in effect 1 in 2 for much of the year)? Who has decided on this arbitrary 1 in 3? How busy must it be to provide adequate training? He mentions continuity in management and post-operative and critical care patients, but surely this is one of the functions of the consultant.

Sir, the issues in the hours of work debate are complex and not to be reduced to simplistic statements. The Hospital Junior Staff Committee, representing and "open to all hospital junior staff," has spent much time and effort negotiating to improve hours of work and to eliminate excessive rotas. This work can and must continue. Perhaps if Mr Porter and his colleagues were to attend these meetings and join the debate they could more usefully contribute. A simple statement of, "1 in 3 no less" is no more valid than the cry, "When I was a boy I worked seven days a week," and adds as little.

PETER HAWKER

West Midlands Junior Staff Committee,
BMA, Birmingham 15

Medical manpower

SIR,—The junior and senior staff conferences welcomed the government's action in removing the exemption to immigration controls for overseas doctors (15 June, p 1843). Both conferences also supported the policy for a reduction in medical student intake.

In the same issue of your journal there are advertisements for 65 senior house officer posts in