CORRESPONDENCE

More corneal grafts A B Tullo, FRCS, and others 735	Chronic bronchial sepsis and progressive lung disease P R Goddard, FRCP, and others 739	Abuse of anabolic steroids S T Garner, BSC, and N A Miles, MB 741
Serum aluminium concentration and aluminium deposits in bone in patients receiving haemodialysis S A Charhon, MD, and P J Meunier, MD; R J Winney, FRCP, and J F Cowie 735	CSM Update: adverse drug reactions and the liver G Lake-Bakaar, MD, and H A Cameron, MRCP; Sir Abraham Goldberg, FRCP 739	IgM and IgG antibodies to HTLV-III in the lymphadenopathy syndrome and subjects at risk for AIDS P Pouletty, MD, and J Kadouche, PHD; F Aiuti, MD 741
Economics of coronary artery bypass grafting A Williams, BCOM; R M Acheson, FFCM 736	Idiopathic hypopituitarism in the elderly W J Jeffcoate, MRCP, and R E Cotton, FRCPATH 739	A fair share of consultations D P B Pound, MRCP 741
General practitioners' advice on smoking to patients referred for barium meals P G Preston, MRCP 737	Screening for Down's syndrome using serum α fetoprotein P R Wyatt, MD 740	Points Rh(D) haemolytic disease of the newborn (H B M Lewis); Death and injury to children in cars (S E Lessels); Respiratory sequelae of whooping cough (J E Cotes); Gastric microflora (D J Seddon); Limited list: limited effects? (F H Leonard); Acute allergic reactions to depot neuroleptic drugs (J M Simister); Quinine for night cramps (H W Daniell); Surgery and the pill (M Waldram); Unit 731 (F G Neild); Body-matters (Caroline van den Brul and M J Hughes-Games) 742
Coma and brain stem areflexia in brain stem encephalitis C Pallis, FRCP; D Wainwright Evans, FRCP, and others 737	Increase of oleic acid in erythrocytes associated with malignancies Su Metcalfe, PHD, and others 740	Correction: Withdrawal of funds for animal experiments (Moore) 742
Metabolic control of insulin dependent diabetes after pancreas transplantation A A P Connolly, MSC 737	Short, Black, Baird, Himsworth, and social class differences in fetal and neonatal mortality R J Lilford, MRCP 740	
Malnutrition, ignorance, and poverty G T Nurse 738	Do locum duties help or hinder acquisition of clinical knowledge by final year medical students? N Banatvala 740	
The low dose aspirin controversy solved at last? R D Eastham, FRCPATH 738	Nurses' education Bernice J M West, MA 741	

Because we receive many more letters than we have room to publish we may shorten those that we do publish to allow readers as wide a selection as possible. In particular, when we receive several letters on the same topic we reserve the right to abridge individual letters. Our usual policy is to reserve our correspondence columns for letters commenting on issues discussed recently (within six weeks) in the BMJ.

Letters critical of a paper may be sent to the authors of the paper so that their reply may appear in the same issue. We may also forward letters that we decide not to publish to the authors of the paper on which they comment.

Letters should not exceed 400 words and should be typed double spaced and signed by all authors, who should include their main degree.

More corneal grafts

SIR,—Over the past decade the number of patients awaiting a corneal graft at this hospital has risen steadily. This reflects two trends. Firstly, as surgical technique has improved more such operations have become feasible. Secondly, fewer, but more specialised, surgeons now undertake keratoplasty. This has resulted in a higher referral rate to regional centres. However, we have until recently been unable to cope with the increased workload because of the lack of donor material. At the beginning of 1984 the United Kingdom Transplant Service (UKTS) first offered the means of providing tissue matched donor material which would be expected to benefit about one third of patients awaiting keratoplasty. UKTS was also able to provide non-matched corneas which would otherwise not have been used in the centres where they were obtained (often the second eye of a pair). Outside the London conurbation Manchester has been the main beneficiary of this service. At the beginning of 1984 our waiting list stood at 120 patients. During the year 76 corneal grafts were performed, an increase of over 100% on the previous year, and our waiting list was reduced by half. Twenty seven donor eyes (35%) were "imported" via UKTS; in the same year we "exported" 13 donor eyes. During the first seven months of 1985 a further 44 grafts have been performed, and the waiting list now stands at 70 patients.

Central to the success of the scheme has been the indexing of information about patients at a local level. We use the computing facilities of the renal transplant service, which issues a monthly printout of all tissue typed patients and many of those awaiting a non-matched corneal transplant.

The removal of eyes should always be considered when permission is being sought for the donation of any other organ. Indeed the eyes can often be removed from a patient who is not suitable to donate another organ. Further details can be obtained from the nearest eye unit, which should be expected to remove the eyes.

UKTS now enables us to accept regional referrals and offer a realistic chance of being able to perform keratoplasty within months rather than years. We also believe it will provide a means of

collating information on a national basis, which is long overdue.

A B TULLO
M K TUTTON
A E RIDGWAY

Manchester Royal Eye Hospital,
Manchester M13 9WH

P DYER

Regional Tissue Typing Laboratory,
St Mary's Hospital,
Manchester M13

Serum aluminium concentration and aluminium deposits in bone in patients receiving haemodialysis

SIR,—We would like to reply to Dr Peter Garrett's comments (29 June, p 1986) on our article (1 June, p 1613).

We have indeed shown that a serum aluminium concentration of over 3.7 $\mu\text{mol/l}$ (100 $\mu\text{g/l}$) was indicative of a high probability (95%) of stainable aluminium in bone. However, the presence of stainable aluminium in bone was not associated with histological evidence of a bone disease related to aluminium accumulation, such as osteomalacia or aplastic bone, in all patients. Thus, the presence of aluminium in bone does not lead unavoidably to aluminium bone disease. A serum aluminium value of over 3.7 $\mu\text{mol/l}$ does not predict whether aluminium deposits in bone have induced decreased bone formation and therefore we believe that a bone biopsy should be recommended in patients with high serum aluminium values to determine the possible part played by aluminium accumulation in the bone disease.

A major problem for the clinician is to diagnose aluminium induced osteomalacia before the patients develop bone fractures. Patients with histological evidence of aluminium induced osteomalacia can be asymptomatic for a long time. There are few serum biochemical indices, other than serum aluminium measurement, including mild hypercalcaemia and relatively low parathyroid hormone levels, which can alert the clinician. But none are specific for aluminium related bone disease. Although bone biopsy is minimally invasive, we can hardly recommend systematic bone biopsies in patients with low serum aluminium values unless there is clinical suspicion of aluminium intoxication. Indeed, most people undergoing dialysis in developed countries using aluminium free water have serum aluminium concentrations lower than 3.7 $\mu\text{mol/l}$.¹ Most are at risk of aluminium intoxication because of the phosphate binders that contain aluminium that they receive orally, and it appears to be difficult in these patients to repeat yearly bone biopsies to identify aluminium accumulation, as suggested by Dr Garrett. Moreover,

the probability of detecting an aluminium induced osteomalacia with a bone biopsy in these patients is low (7/33 patients in our selected group of patients, many of them (14/33) having hyperparathyroid bone lesions unrelated to aluminium intoxication). To assess bone aluminium overload in asymptomatic patients with low serum aluminium concentrations it seems more advisable to perform a desferrioxamine infusion test, which is less invasive than bone biopsy and therefore can be easily repeated. Desferrioxamine chelates aluminium and the peak serum aluminium after desferrioxamine infusion has been shown to be correlated with the bone aluminium content² and to be more sensitive than the baseline serum aluminium measurements in predicting aluminium deposits in bone.³

On the other hand, a much higher proportion of aluminium induced osteomalacia (12/23 patients (52%)) was found in patients with high serum aluminium values. Thus, we suggested performing bone biopsies as a priority in this latter group of patients—or in patients showing a high peak aluminium concentration after desferrioxamine—in an attempt to identify aluminium induced osteomalacia early. We believe that there is practical value in recognising aluminium induced osteomalacia according to well defined histomorphometric criteria. In fact, treatment of aluminium overload includes adequate treatment of the dialysis water, reduction or, when possible, withdrawal of aluminium containing gels, and repeated concentration after desferrioxamine infusions.

Whether it is beneficial for all patients who show stainable aluminium in bone without a mineralisation defect to be treated with desferrioxamine is not yet proved. However, in patients with aluminium induced osteomalacia we and others⁴ have observed very rapid and dramatic improvement of bone mineralisation on repeated bone biopsies under desferrioxamine, even when the patients continue to ingest moderate doses of aluminium hydroxide during the treatment. Thus, we think that patients with histologically proved aluminium related osteomalacia require treatment with desferrioxamine even when they are asymptomatic, to prevent bone fractures.

S CHARHON
P J MEUNIER

INSERM Pathologie des Tissus Calcifiés,
Faculté Alexis Carrel,
69008 Lyon,
France

- Simon P, Allain P, Ang KS, Cam G, Mauras Y. Prophylaxie et traitement de l'intoxication aluminium chez l'insuffisant rénal chronique. In: Crosnier J, Brentano JLF, Bach JF, Grunfeld JP, eds. *Actualités néphrologiques de l'Hôpital Necker*. Paris: Flammarion, 1984:343-414.
- Milliner DS, Nebeker HG, Ott S, et al. Use of deferoxamine infusion test in the diagnosis of aluminium-related osteodystrophy. *Ann Intern Med* 1984;101:775-80.
- Berland Y, Charhon SA, Olmer M, Meunier PJ. Predictive value of desferrioxamine infusion test for bone aluminium deposits in hemodialyzed patients. *Nephron* 1985;40:433-5.
- Malluche HH, Smith AJ, Abreo K, Faugere MC. The use of deferoxamine in the management of aluminium accumulation in bone in patients with renal failure. *N Engl J Med* 1984;311:140-4.

SIR,—Dr Sam A Charhon and others (1 June, p 1613) found that a single estimation of serum aluminium was of limited value in predicting the presence of aluminium staining or aluminium toxicity in bone. However, both of these features were predictable in a significant proportion of the patients studied on the basis of a serum aluminium of $>7 \mu\text{mol/l}$ ($189 \mu\text{g/l}$). While they point out that the population studied may well not have been in a steady state in relation to aluminium exposure, they conclude that, under steady state conditions, a serum aluminium concentration of $3.7 \mu\text{mol/l}$ ($100 \mu\text{g/l}$) should be the threshold beyond which biopsy of bone should be performed to enable early diagnosis and treatment of bone toxicity. Dr Peter Garrett (29 June, p 1986) questions their conclusions and is of the opinion that the only real

value of serum aluminium measurement (which he regards as difficult both to measure and to interpret) is in identifying recent aluminium loading. He concludes that patients at risk of aluminium toxicity should have yearly bone biopsies but does not indicate how patients at risk should be identified.

The regular measurement of plasma aluminium concentrations in patients undergoing haemodialysis in our unit has shown that the prevailing plasma aluminium concentration is a sensitive index of exposure from either dialysis fluid or the oral ingestion of aluminium; the plasma aluminium value rises with exposure and falls slowly after removal from exposure. When regular measurement of plasma aluminium is combined with similar monitoring of dialysis fluid, the source of exposure can be identified since, in the presence of a dialysis fluid aluminium $<1 \mu\text{mol/l}$ ($27 \mu\text{g/l}$), a rising plasma aluminium concentration indicates excessive oral intake of aluminium. Thus regular measurement of both plasma and dialysis fluid aluminium concentration identifies patients at risk of aluminium toxicity and enables the introduction of measures to minimise further exposure and so prevent toxicity.¹

The plasma aluminium concentration is also of considerable predictive value in clinical aluminium bone disease.¹ A plasma aluminium value consistently $>7 \mu\text{mol/l}$ ($100 \mu\text{g/l}$) indicates a 100% certainty of clinical bone disease either being present or developing unless exposure is reduced. By contrast, no patient under our care with a plasma aluminium value maintained $<5 \mu\text{mol/l}$ ($135 \mu\text{g/l}$) has developed clinical aluminium bone disease over the past six years. A single estimation of plasma aluminium will be of lower predictive value since exposure may be fluctuating and high levels may be missed. Indeed, a single measurement, if low, may be misleading.

Dr Charhon and colleagues suggest that the finding of bone aluminium deposits and histological features of aluminium bone disease may enable early treatment with desferrioxamine to prevent the development of osteomalacia but do not indicate whether any of their patients had clinical or radiological features of aluminium bone disease. We have found that resolution of clinical, radiological, and histological features of aluminium bone disease may occur simply with removal from exposure to aluminium. Particularly since recent exposure to aluminium could not be excluded in their patients, we would suggest that resolution might also occur in some of their patients with minimisation of further exposure. Since desferrioxamine is not without adverse effects, there is a need for studies of the clinical course of bone aluminium deposits and subclinical aluminium bone disease in patients in whom exposure to aluminium is minimised before desferrioxamine is routinely advocated for such changes.

We would suggest that the main emphasis should be on preventing aluminium toxicity in patients undergoing haemodialysis. Regular monitoring of plasma and dialysis fluid aluminium concentrations is a simple and non-invasive means of detecting patients at risk of aluminium toxicity, enabling the introduction of measures to minimise further exposure and the prevention of aluminium toxicity. Since the introduction of regular monitoring in 1979 we have seen only two patients with clinical aluminium bone disease, both of whom had a history of exposure and at one stage a plasma aluminium $>7 \mu\text{mol/l}$ ($100 \mu\text{g/l}$). While some of our patients may have undetected bone aluminium staining and aluminium induced disturbance of bone metabolism, there is no evidence of progressive bone disease over the five years during which exposure to aluminium has been maintained.

R J WINNEY
JOHN F COWIE

Medical Renal Unit, Royal Infirmary,
Edinburgh EH3 9YW

- Winney RJ, Cowie JF, Robson JS. What is the value of plasma/serum aluminium in patients with chronic renal failure. *Clin Nephrol* 1985;24(suppl 1):S2-8.

*This correspondence is now closed.—Ed, *BMJ*.

Economics of coronary artery bypass grafting

SIR,—Professor R J Jarrett will doubtless be pleased to know that I am very easily provoked, and perhaps still more pleased that I cannot answer all his points very satisfactorily. This is not entirely my fault, however, nor even that of my fellow health economists, nor even, for that matter, that of doctors, though I think we may all share some small part of the responsibility for the fact that in the present state of our knowledge we cannot answer the central fundamental question facing all health services all over the world—namely, how should the resources available for health care be deployed so as to maximise the health of the community?

Professor Jarrett tries to sidestep this issue in his last paragraph, believing erroneously that if more were spent on health the problem would disappear, whereas all that would happen is that it would present itself again in a different form in a different context. It is chastening to note that countries such as Sweden, West Germany, and the USA, which spend far more on health care than we do, also face the same kinds of problem.¹ Romantic escapism may be comforting but it is not helpful.

So proceeding on the assumption that he is now convinced that even if we had more "welfare" and less "warfare" hard choices would remain necessary, what is the alternative to cost per quality adjusted life year as the theoretically correct measure to aim at when establishing priorities? I see no answer to that key question in his letter. Instead I see only a rehearsal of the practical difficulties, and no one is denying that they exist in abundance. Indeed, I highlighted them myself. Professor Jarrett's position seems to be that until we can measure quality of life with the certainty with which we can measure mortality, then we had better ignore it, because it is too soft. This is tantamount to saying that although we know people value a good quality of life more highly than a bad one, since we are not sure exactly how much more highly they value it, we had better count it as of zero value, and assume that they care only about life expectancy. Surely it cannot be better to use the one valuation we know is wrong in preference to our best estimate of what is right?

It would, of course, be better if we had largescale survey data from impeccably designed trials using well validated instruments to measure outcomes descriptively in terms that relate to quality of life, and equally good data on how people value such health states relatively to each other, and I have been trying for many years to encourage people to do just that. But I have come to the view that until people are convinced that the methods themselves are correct in principle—that is, they answer the question posed, and deploy the relevant information, in an appropriate way—then it is unlikely that the methods will be taken seriously enough for the requisite data ever to be collected and deployed, so that their potential to enable us to grapple more effectively with our central policy problem will never be realised.

Although I regard adventurous exploration of this relatively untrodden territory as very important, I do not expect everyone to jump on to the quality of life bandwagon and abandon "hard endpoint" epidemiology, for there is still an important role for that kind of investigation if we want precise answers to more circumscribed questions. But that will not enable us to grapple with the choice between more total hip replacements and more hospital haemodialysis. So, if I may end on a mischievous note, I think the main difference between us is epitomised in the old story about the helpful passerby who found a man on his hands and knees under a lamp-post searching for his doorkey, and who kindly joined in the search.