CORRESPONDENCE

Death or dialysis

J S Cameron, FRCP; Mary G McGeown, FRCP, and others; L Griffiths, MB; K R Llewellyn, FRCP 236

Severe hypophosphataemia during recovery from acute respiratory acidosis

P Squara, MD, and others 237

Screening for small for dates fetuses

I Kennedy, FRCSED 237

High mean platelet volume after myocardial infarction

T M E Davis, MB; A Trowbridge, PHD, and others; R Sewell, MRCP, and others .. 238

Domiciliary immunisation for preschool child defaulters

R B McGucken, FRCP 239

Efficacy of facemask resuscitation at birth

A Mackie, FFARCS; A D Lawson, MB, and B Sweeney, FFARCS 239

Teaching terminal care

P M Kaye, MRCP 239

Hospital clinic care versus general practice care for type II diabetics

T M Hayes, FRCP, and J Harries, MRCPG .. 240

Further developments in psychogeriatrics in Britain

D J Jolley, FRCPsych 240

Management budgets in the NHS

Marguerite A R King, FRCPATH; P Davies, MSC, and M Prowle, MA 240

Infection with parvovirus during pregnancy

E P Wright, MRCPATH, and others 241

Pathways to complement activation during cardiopulmonary bypass

D Royston, FFARCS; J Ponte, FFARCS 241

Domperidone

Jenny E Leiser, MRCP, and D N Bateman, MRCP; G Lake-Bakaar, MD, and H A Cameron, MRCP 241

The link with zinc

G S Fell, MRCPATH 242

Defibrillators in general practice

R A Dixon, PHD, and E Wilkes, FRCP 242

Scandinavian model for eliminating measles, mumps, and rubella

A S Thornton, MRCPG; J Taranger, MD, and E Rabo, MD 242

Design and interpretation of clinical trials

A V Pollock, FRCS, and Mary Evans, BA .. 243

Nodular malignant melanoma in a patient treated by photochemotherapy

R A C Graham-Brown, MRCP, and others . 243

Hypoglycaemia during illness in children with congenital adrenal hyperplasia

A W McNinch, MRCP, and D C L Savage, FRCP 243

Weight gain after cholecystectomy

R M F Kwan, MB, and M A Mir, MRCP .. 243

Voice changes after thyroidectomy: role of the external laryngeal nerve

R E May, FRCS 243

Blood lead concentration and blood pressure

G Orssaud, MD, and others 244

On the state of the public ill health

D H Vaughan, FFCM 244

Doctors, drugs, and the DHSS

C M P Claoué, MB; J H Beaven, MB 244

One Man's Burden: setting the record straight

J Williams, FMA 245

Help for the Sudan

H M Lipman, MRCPG 245

Points Sleep helps healing (P G F Nixon); Cigarette lung (M C Kelly); "Tissue thyrotoxicosis" (S Taylor); Facilitating prevention in primary care (D Stephens); Factors affecting development of peritonitis in continuous ambulatory peritoneal dialysis (T H J Goodship; R Gabriel); "BNFs" for the Third World (R Philipp) 245

Because we receive many more letters than we have room to publish we may shorten those that we do publish to allow readers as wide a selection as possible. In particular, when we receive several letters on the same topic we reserve the right to abridge individual letters.

Letters critical of a paper may be sent to the authors of the paper so that their reply may appear in the same issue. We may also forward letters that we decide not to publish to the authors of the paper on which they comment.

Letters should not exceed 400 words and should be typed double spaced and signed by all authors, who should include their main degree.

Death or dialysis

SIR,—Mrs Elizabeth Ward's leading article (22-29 December, p 1712) is welcome in that it highlights yet again a national and international disgrace. She deals in some detail with a proposal for helping to cope with the problem but in a fashion far from complete: the true situation is both more complicated and more simple.

Mrs Ward suggests the adoption of contract dialysis to diminish the gross shortfall in facilities for centre dialysis in the United Kingdom for those patients who are unsuitable for transplantation or home peritoneal dialysis or haemodialysis. She has been surprised and saddened to find doctors caring for patients with end stage renal failure less than enthusiastic about these proposals, but the reasons are not difficult to find.

Regular dialysis at the figure of £60 per dialysis that Mrs Ward quotes—that is, £9360 a year for thrice weekly dialysis—is not such an improvement on what is already being achieved in many NHS dialysis units, and may be worse than in some. The real cost of a patient on regular dialysis, whether on haemodialysis or peritoneal dialysis, includes not only the raw cost of performing the dialysis (including staff costs) but also the cost of preparing the patient by creating a fistula or inserting an intraperitoneal catheter,

plus any revision of these; the cost of drugs (other than those such as heparin specifically required for the dialysis); the cost of hospital admissions, at £100-£200 a day, and more if intensive care is required; and the cost of a proportion of the salaries of the nurses and doctors who provide the back up and medical supervision of the patient. Costings in the United Kingdom¹ and elsewhere suggest that these extra costs represent an additional £3000-£5000, depending on the type of patient. In 1984 the DHSS quoted figures of £12 300 to £15 500 as average total costs for centre dialysis (A Hurst, personal communication), which cannot include more than about £9000-£10 000 for the dialysis alone.

In fact, an average figure for these extra costs would be a gross underestimate for those patients whom one hopes will benefit from an expansion of centre dialysis in the United Kingdom. Those who do not now receive treatment are, by and large, those aged over 60² and those with complicated diseases such as diabetes mellitus. Both groups make greater use of hospital facilities than the average patient on dialysis, and the hospital costs for a single such patient can run into tens of thousands of pounds in a year.

Thus, transfer of funds to a contractor, without allocating extra revenue to prevent

depletion of the necessary support services, would result in worse level of care for the needy, since no unit could afford to be responsible for a group of patients in a contracted out, freestanding facility performing only haemodialysis, which would telephone the hospital at the first sign of expensive trouble. Moreover, if North American experience is anything to go by, it would probably increase the total costs.

Almost all renal physicians in the United Kingdom would prefer that the number of NHS dialysis units should be expanded, the new units being placed in those regions with a poor record of treating end stage renal failure. It cannot be an accident that every other country in Western Europe, despite varying patterns of health finance, has opted for 3-4 units per million population, while the United Kingdom lags behind with only 1.1 per million.² A poorer, but effective, alternative would be to expand existing units. Renal services in the United Kingdom are all but saturated,³ with valuable plant lying idle up to half the time⁴—as in our own unit—because there is no money to staff the extra shifts. This money can come only from the regional health authorities, or from central allocation, either from cuts in other services or from an increase in total expenditure on health. In national

terms the amount needed—£30–£40m a year at most—is trivial; fortunately, chronic renal failure is rather rare.

There is no margin of inefficiency in the performance of dialysis which contractual dialysis can exploit; it comes down simply to the fact that until and unless the United Kingdom health service spends more on services for renal failure than some of those who should benefit from the treatment will die, and tinkering with the available budget in politically fashionable ways will not help them.

Finally, despite twofold variations in rates of treatment throughout the United Kingdom for terminal renal failure,⁴ few renal physicians actually “turn down” more than a handful of patients each year. Of course, there remains great variation in patients considered “suitable” in different parts of the country, and Mrs Ward is right to attack this, but a crucial point in the undertreatment of renal failure in the United Kingdom is underreferral.⁵ Why do many physicians and general practitioners not refer patients whom the dialysis units would regard as entirely treatable? There appears to be a subtle feedback from the attitudes of the local dialysis unit to the local physicians and practitioners, perhaps relating to events in the distant past, when treatment was much more experimental than it is today. This particularly applies to older patients—even in our own region (South West and South East Thames), where intake now approaches 50/million/year (over 300 patients/year), we are still referred few patients over 70 years of age, and almost none over 80.

STEWART CAMERON

Renal Unit,
Guy's Hospital,
London SE1 9RT

- 1 Pincherle G, Mancini PV. The costs of treating renal failure. In: Parsons FM, Ogg CS, eds. *Renal failure: who cares?* Lancaster: MTP Press, 1982: 25–33.
- 2 Wing AJ, Selwood NH. Achievements and problems in the treatment of end stage renal failure. In: Jones NF, Peters DK, eds. *Recent advances in renal medicine*. Edinburgh: Churchill Livingstone, 1982:103–19.
- 3 Jones NF, Goodwin FJ, Roberts AP. Manpower and workload in adult renal medicine in the United Kingdom 1975–1983. *Br Med J* 1984;288:992–4.
- 4 Davison AM, Read DJ, Lewins AM. Underutilised hospital haemodialysis resources. *Lancet* 1984;i: 723–5.
- 5 Challah S, Wing AJ, Bauer R, Morris RW, Schroeder SA. Negative selection of patients for dialysis and transplantation. *Br Med J* 1984;288:1119–22.

SIR,—We welcomed Mrs Elizabeth Ward's leading article on the continuing and scandalous shortage of treatment for patients with end stage renal failure. We agree with her that there is underprovision of facilities, that patients are being denied treatment, and that the fundamental issue of responsibility—government, Department of Health, regional health authority, or renal physician—is unresolved. At the sharp end, the renal physician continues to face a harrowing moral and ethical dilemma.

We disagree with Mrs Ward when she says that the most important factor in addition to lack of facilities is a lack of concern by renal physicians. The facts regarding the plight of Northern Ireland patients (including annual lists of names of patients turned down for treatment) have been repeatedly placed before the regional health authority and the health minister. The response has been sympathetic, but further finance to expand the service has not been forthcoming. It may be that something more dramatic on a national basis will be necessary before action results,

but it is nevertheless unhelpful, untrue, and offensive to malign the sincere efforts of renal physicians to solve the problem as constituting concealment of facts and a tacit acceptance of the present situation so that we can “sleep at night.”

MARY G MCGEOWN
JAMES F DOUGLAS
CIARAN C DOHERTY

Renal Unit,
Belfast City Hospital,
Belfast BT9 7AB

SIR,—On 8 January, in various radio and television items on the patient in Oxford who has had his renal dialysis discontinued (see p 249), I learnt his name and address; his history of earlier brain surgery; opinions on his present mental state; and details of his bowel and urinary functions, his raised blood pressure, lack of bodily cleanliness, and disturbed behaviour. One presumes that he had given his informed consent to the broadcast of these clinical details.

L GRIFFITHS

Llanrwst,
Gwynedd

SIR,—The lack of quality in Mr Derek Sage's life seems to have suggested that its prolongation was pointless, but he may turn out to be a seminal figure.

Compared with most countries of the West, Britain is inefficient, but the material resources available to us now are far greater than was the case at the start of the health service. On the demand side huge areas of disease have disappeared—tuberculosis, for example. The economies practised go far beyond the refusal to continue treatment of Mr Sage or even start treatment in many similar cases. I was horrified to learn the other day that the age at which amniocentesis can be offered in this area under the NHS has been raised from 35 to 38. My younger colleagues in many specialties tell me of falling standards in the NHS arising from inadequate resources.

Are we to move ever nearer to the American standards of morality in these things? Aneurin Bevan should have held out for a wholtime health service: the Labour party should make it an election plank even if it “walks it” for an election or two. Fighting the future battles of the health service will be a wholtime job. We may lose some of our best men to the flesh pots, but the fight will be too tough for part time soldiers.

K R LLEWELLIN

Chester CH1 6LG

Severe hypophosphataemia during recovery from acute respiratory acidosis

SIR,—We read with great interest Dr Tommy L Storm's article (25 August, p 456) about three patients with respiratory failure who showed profound hypophosphataemia during mechanical ventilation. These results strengthened a similar study of 36 patients with obstructive pulmonary disease that we recently reported.¹

When the patients were admitted serum phosphate concentrations were normal. After mechanical ventilation we observed a constant and significant fall in serum phosphate values (0.54 (SD) 0.14 mmol/l (1.7 (0.4) mg/100 ml) after the first 24 hours). In four cases there was profound hypophosphataemia (<0.30 mmol/l (0.9 mg/100 ml)).

The alkaline variation in pH was significantly related to the fall in serum phosphate values ($r=0.67$; $p<0.001$). Even when there was no respiratory alkalosis during mechanical ventilation there was a fall in phosphate concentrations when the pH became more. We also observed this phenomenon in patients with phosphate depletion (chronic alcoholism) and in those who were hyperphosphataemic at the time of their admission (renal failure). Then serum phosphate values tended to return to normal spontaneously and slowly without phosphate repletion. After extubation serum phosphate concentrations returned to normal within 36 hours. At the same time urinary phosphate excretion, which was normal when the patients were admitted, fell significantly during mechanical ventilation and returned to normal after extubation; this suggests a phosphorus shift into the intracellular compartment.

These observations must be connected with the work of Mostellar and Tuttle² and, more recently, with that of Brautbar *et al*,³ which show a cellular shift of phosphorus in connection with increased glycolytic activity when two essential factors are associated: respiratory alkalosis and glucose infusion.⁴ The main problem is to know what effect this hypophosphataemia has during respiratory failure. The change in left ventricular function, the fall of P_{50} , may alter spontaneous ventilation, especially after extubation. Moreover, we have shown that diaphragmatic contractility of the diaphragm is significantly impaired when serum phosphate concentrations are less than 0.40 mmol/l (1.2 mg/100 ml) (M Aubier *et al*, paper at Réunion de la Société de Langue Française des Maladies Respiratoires, Paris, 1984).

Although muscular weakness has been in phosphorus depleted patients or animals, it is possible that the intracellular shift in phosphate may induce a dysfunction of the cell muscle membrane, which may by itself impair diaphragmatic contractility. Further studies are needed to confirm whether the same mechanism occurs in hypophosphataemia due to phosphorus cellular shift, when the phosphorus supply is constant.

P SQUARA
G BLEICHNER
J P SOLLET
A PARENT

Service Réanimation Polyvalente,
Hôpital d'Argenteuil,
France

M AUBIER

Clinique Pneumologique,
INSERM U226,
Hôpital Beaujon,
Clichy,
France

- 1 Squara P, Bleichner G, Aubier M, Parent A, Sollet JP, Murcland D. Hypophosphorémie au cours de la ventilation mécanique des BPCO. Congrès de réanimation de langue française: Paris 1983. *Presse Médicale*. (In press.)
- 2 Mostellar ME, Tuttle EP. The effect of alkalosis in plasma concentration and urinary excretion of inorganic phosphate in man. *J Clin Invest* 1964;43: 138–49.
- 3 Brautbar N, Leibouich, Massry SG. On the mechanism of hypophosphatemia during acute hyper-ventilation: evidence for increased muscle glycolysis. *Mineral and Electrolyte Metabolism* 1983;9: 45–50.
- 4 Mansour TE. Studies on heart phosphofructokinase. *J Biol Chem* 1963;238:2285–92.

Screening for small for dates fetuses

SIR,—In their review of methods of detecting intrauterine growth retardation Dr J P Neilson and his colleagues dismiss the “symphysis-fundalists” because of only one unfavourable report¹ ignoring the many favourable ones^{2–5} and the work of Boddy in Edinburgh.

From their description, the method used by Rosenberg *et al* to measure symphysis-fundus height¹ was less careful and precise than that of the other authors, a point particularly emphasised by Gentz *et al*.⁶ It was also a