β -BLOCKADE AFTER MYOCARDIAL INFARCTION— A STATISTICAL VIEW

J.A. LEWIS

Imperial Chemical Industries PLC, Pharmaceuticals Division, Macclesfield SK10 4TF

During the past year several trials have been published which have increased the evidence in favour of a beneficial effect of treatment with a β -adrenergic blocking agent on mortality after myocardial infarction (Norwegian Multicentre Study Group, 1981; Hjalmarson *et al.*, 1981; Beta-blocker Heart Attack Study Group, 1981; Hansteen *et al.*, 1982). There are, however, such wide differences between the results obtained in the many trials published in the last 15 years or so that clinical interpretation of all the available information is difficult. Clinical reviews have been conducted in the past (Baber & Lewis, 1980; 1981; Baber, 1981; Hampton, 1981; Vedin & Wilhelmsson, 1979) using some statistical methods.

This paper attempts to clarify the present state of knowledge by using statistical techniques more fully.

Methods

Published trials

The literature was surveyed to identify the published trials of β -blockers in this indication, and 29 trials were identified. It is important in a statistical review to include *all* trials because of the possibility of bias in a selected subset. For example, negative trials are not always reported and hence

Table 1 Early entry studies

<i>Trial</i>	<i>Size</i>	<i>Drug</i>	<i>Dosing</i>	<i>Time from first symptoms to treatment</i>	<i>Length of follow-up</i>	<i>Infarct</i>
Wilcox <i>et al.</i> , 1980	473	Oxprenolol	Oral	Under 24 h	6 wks	Suspected
Norris <i>et al.</i> , 1968	454	Propranolol	Oral	Under 3 days	3 wks	Definite
Multicentre, Stephen, 1966	195	Propranolol	Oral	Under 48 h	4 wks	Definite
Andersen <i>et al.</i> , 1979	480	Alprenolol	Intravenous and oral	Median 6 h	1 yr	Suspected
Balcon <i>et al.</i> , 1966	114	Propranolol	Oral	Under 24 h	4 wks	Definite
Wilcox <i>et al.</i> , 1980	388	Propranolol	Oral	Under 24 h	1 yr	Suspected
		Atenolol				
Barber <i>et al.</i> , 1975	298	Practolol	Oral	Median 3.5 h	2 yrs	Definite
Barber <i>et al.</i> , 1967	99	Propranolol	Oral	Under 24 h	4 wks	Definite
Hjalmarson <i>et al.</i> , 1981	1395	Metoprolol	Intravenous and oral	Mean 11 h	3 mths	Suspected

Table 2 Late entry studies

<i>Trial</i>	<i>Size</i>	<i>Drug</i>	<i>Time from first symptoms to treatment</i>	<i>Length of follow-up</i>
Baber <i>et al.</i> , 1980	720	Propranolol	2-14 days	3-9 mths
BHAT, 1981	3837	Propranolol	5-21 days	30 mths
Hansteen <i>et al.</i> , 1982	560	Propranolol	4-6 days	12 mths
Wilhelmsson <i>et al.</i> , 1974	230	Alprenolol	About 6 wks	24 mths
Coronary Prevention Research Group, 1980	1134	Oxprenolol	Median 7 mths	Several yrs
Green <i>et al.</i> , 1977	3053	Practolol	1-4 wks	1-2 yrs
Pedersen <i>et al.</i> , 1981	1884	Timolol	1-4 wks	33 mths

even a review of all published trials might present an overoptimistic view of the effect of treatment. If any negative trials were missed for this reason, however, they were almost certainly small trials with little influence on the overall picture. Large trials are expensive, difficult to mount, and well known to the profession, so that publication is almost inevitable whether the results are positive or negative. Fortunately, the statistical review was dominated by large trials.

Selection of trials for critical analysis

The 29 trials were critically examined to establish whether they were suitable for inclusion in a pooled analysis and whether they were satisfactorily conducted by modern standards. The following criteria were used to select trials for further analysis: (a) more than 20 deaths recorded. The precision of a trial depends largely on the number of deaths, and imprecise trials reduce the power of some of the statistical tests used; (b) proper randomization. Randomization by date of birth or alternate entry to each group was not considered acceptable; (c) control. Trials of oral β -blockers were excluded if the control group did not receive placebo. In the case of acute intravenous treatment this condition was not imposed; (d) total mortality by 'intention to treat' recorded. The analysis of total mortality irrespective of cause by 'intention to treat' is well recognized as the proper pragmatic approach. It avoids potential bias due to differential mortality in withdrawn or defaulting patients or due to differential diagnosis.

Using these criteria I selected 16 trials (see Tables 1 and 2).

Classification of trials

Trials can be classified according to the clinical intention of either reducing the amount of myocardial damage of the index infarct by early intervention or reducing mortality after recovery from the event. Many studies originally intended to be early cannot now be regarded as such, as intervention occurred too late to influence the evolving infarct. This does not negate the importance of the classification for the purposes of analysis, as there are currently insufficient data to subcategorize the early intervention trials to investigate this point adequately.

A trial admission time of 48 hours after the acute event was taken as the cut-off between early and late intervention. This was found to give a clear division according to the intentions of the trials. The early and late trials are listed in Tables 1 and 2

with their important features. Seven were late and nine were early.

The included trials: additional information

Some trials need special comment.

1 Wilcox *et al.* (1980a) compared two β -blockers (atenolol and propranolol) with placebo in three corresponding groups of patients. The comparison of each β -blocker with placebo was included separately in the analysis, leading to a total of 17 comparisons.

2 In the paper by Andersen *et al.* (1979) the patients were split into those aged under 65 and those aged over 65 for the purpose of analysis. No other trial has confirmed a differential effect of β -blockade on postinfarct mortality according to age; nor is it entirely clear that the protocol of this trial specified that the beneficial effect was only to be sought in patients under 65. For these reasons the data were combined over age groups for this paper.

3 In the original report of the trial of oxprenolol conducted by the Coronary Prevention Research Group (1980) figures for total mortality were not provided. They were, however, later clarified by Hampton (1981).

4 Total mortality in the Multicentre International Study of practolol (1977) was available to me through my statistical work in the trial.

Statistical methods

Percentage treatment effects and their confidence limits

Figure 1 contains percentage treatment effects and their 95% confidence limits for the end-points of individual trials. The percentage effect of the treatment was defined as:

$$\frac{(\text{Control mortality rate} - \text{Mortality rate on } \beta\text{-blocker}) \times 100\%}{\text{Control mortality rate}}$$

The horizontal bars correspond to the 95% confidence intervals for this percentage effect. Thus the righthand end of each bar gives an upper limit for the beneficial effect of treatment and the lefthand end gives a lower limit—generally corresponding to a deleterious effect of treatment. The actual percentage effect of treatment on mortality is represented by the vertical line towards the centre of the bar. For sound technical reasons

Figure 1 Effect of β -blockers on post-infarction mortality. Difference in mortality expressed as a percentage of control rate in 17 controlled trials of β -blockers. 95% confidence limits based on odds ratio considerations. □ 'early' entry, ▨ 'late' entry.

the confidence limits were asymmetrically placed about the observed effect and this prevented the upper confidence limit exceeding 100% reduction in mortality, the maximum possible benefit. To take an example, in the Beta-blocker Heart Attack Trial (BHAT) the observed effect of propranolol on total mortality was a 26% reduction; 95% confidence limits ranged from a 9% reduction to a 41% reduction and these limits are likely to contain the 'true' effect of treatment intervention on mortality in this study. The statistical methodology for calculating such confidence limits can be found in Thomas & Gart (1977) and Baptista & Pike (1977).

Pooling groups of studies and contrasting groups of studies

Studies were pooled by fitting log-linear models (Nelder & Wedderburn, 1972) using the computer package GLIM (General Linear Interactive Modelling) (Baker & Nelder, 1978). The technique enabled the following types of questions to be investigated: (a) Is the proportional effect of treatment on mortality in a group of studies sufficiently consistent from study to study for any differences to be attributable to chance? (b) If so, is the effect of treatment statistically significant, combining the results from the group of studies? (c)

When two groups of studies are compared (early and late) is there a difference in the proportional effect of treatment on mortality? By combining the results of fitting log-linear models with the Thomas & Gart approach, it also proved possible to estimate the size of the pooled effect of treatment on a group of studies, together with 95% confidence limits for this effect. Such pooled intervals are shown in Figure 2.

Figure 2 Effects of β -blockers on post-infarction mortality. Pooled results for 17 trials with 95% confidence limits.

Figure 3 Thirty trials of β -blockers post infarction—effects on mortality and approximate relative weights. Weights based on reciprocals of asymptotic variances of percentage treatment effects. $\square$ excluded trial, $\blacksquare$ 'early' entry, $\square$ 'late' entry.

Relative importance of trials The width of the confidence intervals as calculated for Figure 1 gave only a rough idea of the relative size of the contribution made by each trial to the pooled results derived by the methods used for Figure 2. A more informative idea was gained by calculating weighting factors for the percentage treatment effects, as described by Tarone (1981). These weights (which were proportional to the reciprocals of the estimated asymptotic variances of the percentage treatment effects in each trial) were plotted against the observed treatment effect as shown in Figure 3.

Results

Figure 1 displays the percentage effects of treatment and their 95% confidence intervals for the 17 selected trials. Two features are worth noting: Firstly, there was an overall tendency for the observed percentage treatment effect to lie to the right of the zero line—that is, to correspond with a reduction in total mortality. Twelve results were positive and five were negative. Secondly, all trials were consistent with a treatment effect of about a 20% reduction in mortality.

The 17 selected trials were combined in a pooled analysis and there was a highly significant effect of β -blockade ($\chi^2_1=22.2$; $p<0.00001$). There was no evidence of heterogeneity of the treatment effect

($\chi^2_{16}=17.0$)—that is, the results of the separate trials were consistent statistically. The pooled effect on mortality was 21% with 95% confidence limits ranging from 13% to 28% (Figure 2).

Because of the sound clinical reasons for contrasting the 10 early- and the 7 late-entry trials, this was carried out despite the absence of statistical evidence of heterogeneity. The contrast between the two groups achieved the significance level of $p=0.045$, which is convincing evidence that early and late trials do differ in outcome.

The analysis of the 10 early-entry trials showed a small positive effect, but this had no statistical or clinical significance. (See Figure 2; the overall percentage reduction was 8%, $\chi^2_1=1.1$.)

Pooled analysis of the seven late-entry trials gave rise to a highly significant effect of β -blockade on total mortality ($\chi^2_1=25.1$, $p<0.00001$) with no evidence of heterogeneity ($\chi^2_6=5.5$). The pooled effect on mortality was 26% with 95% confidence limits ranging from 17% to 35% (Figure 2). This result was heavily dominated by the practolol multicentre trial, BHAT, and the Norwegian timolol trial. Of the remaining four trials, only one (Baber, 1981) showed a negative effect of treatment and this was a comparatively small trial with correspondingly wide confidence intervals.

Some support for the positive effect observed in the late-entry trials could be found in the early-entry trials with prolonged follow-up. Five of the

pooled early trials had a follow-up of three months or more: Wilcox (atenolol), Andersen, Barber (practolol), Hjalmarson, Wilcox (propranolol). Of these, four showed a positive trend and only one a negative trend. This contrasted with the other five pooled early-entry studies, where two showed a positive trend and three a negative trend. Hjalmarson and Andersen were the most precise of the early trials with long follow-ups, as evidenced by their narrow confidence intervals in Figure 2. In both these studies the favourable effect on mortality did not start to appear until a week or so after the infarct, despite the high early mortality.

The relative importance of the various trials is illustrated in Figure 3. All 30 comparisons are displayed in this figure, which can be interpreted in a similar way to a histogram. The observed percentage effect of treatment was divided into 5% intervals. Trials were assigned to the interval corresponding to the treatment effect in that trial, and their weights were accumulated vertically. The following features of Figure 3 are striking: (a) the bulk of the evidence contained in the trials centred on a 20% to 40% reduction in mortality associated with β -blockade; (b) the figure was dominated by three trials—BHAT, the multicentre timolol trial, and the multicentre practolol trial; (c) the trials excluded from the overall analysis had very little impact on the results; (d) most of the evidence favouring β -blockade was associated with late-entry trials.

Discussion and conclusions

There is a consistent and very highly significant reduction in total mortality of 26% (17%–35%) in late-entry trials. This conclusion would not be affected by decisions taken relating to trial exclusion. There is no statistical evidence that the effect varies from trial to trial—that is, that it is influenced by the choice of β -blocker or patient population. Such differences may exist, but if they do they are of secondary magnitude and it is likely to be very difficult to quantify them sufficiently accurately to be of clinical value.

There are a few late-entry trials of β -blockers given after infarction that are still in progress. Because of their expected size, these trials would need to produce some dramatically adverse effects of β -blockade on mortality to affect the conclusions of this paper. If there were any effect on the

conclusions, it would probably point to some peculiarity of the trials yet to be published. The positive effect of β -blockade on mortality is also observable in the early-entry trials with long-term follow-up but not during the early part of these trials. Indeed, a division of trials into those with long-term follow-up and those with only a few weeks follow-up would produce very similar results to the division carried out in this analysis. This suggests that the benefit is due to the effect of β -blockade on the outcome of new events rather than on any specific consequence of the original infarction.

This beneficial effect of long-term β -blockade must now be considered proved beyond any reasonable doubt and it remains only for the clinical impact of this finding to be realised in practice. Enhanced effects are generally observed in the mortality findings among patients who did not withdraw from treatment—for example, the Multicentre International Study (1977). Therefore, the benefit may be even larger in patients who comply with treatment, perhaps as much as a 30% to 35% reduction in mortality.

The conclusion for early intervention is still uncertain. Overall, the beneficial effect of β -blockade on early mortality in the trials published so far is certainly proportionately smaller than the effect on later mortality and may well be of no clinical value. There is, however, convincing evidence that intervention with an intravenous β -blocker sufficiently early after the onset of pain can limit infarct size (Yusuf *et al.*, 1980). Most of the patients in the trials described as early in this paper would not have achieved clinically valuable blood concentrations of the drug during the crucial hours when the infarct was developing. A corresponding effect on mortality and morbidity seems plausible and the outcome of trials to investigate this possibility is awaited with great interest.

This paper is based on the pooling of results of clinical trials, a procedure which is not without its opponents. I do not intend to debate the issue here. Instead I hope that the practical demonstration that information of great clinical value can be extracted by such a process will be its own defence.

I should like to thank Dr. N.S. Baber and Mr. S.H. Ellis with whom I have worked collaboratively on the medical and statistical material on which this paper is based. I should also like to thank Mr. R. Peto for many helpful discussions.

References

- AHLMARK, G. & SAETRE, H. (1976) Long term treatment with beta-blockers after myocardial infarction. *Europ. J. Clin. Pharm.*, **10**, 77-83.
- ANDERSEN, M.P., FREDERIKSEN, J., JURGENSEN, H.J., PEDERSEN, F., BEERSGAARD, P., HANSEN, D.A., NIELSEN, B., PEDERSEN-BJERGAARD, O. & RASMUSSEN, S.L. (1979). Effect of alprenolol on mortality among patients with definite or suspected acute myocardial infarction. *Lancet*, **2**, 865-868.
- BABER, N.S. (1981). Clinical experience with beta-adrenergic blocking agents in myocardial ischaemia: a dilemma and a challenge. *Pharmac. Ther.*, **13**, 285-320.
- BABER, N.S. & LEWIS, J.A. (1980). Beta-blockers in the treatment of myocardial infarction. *Br. Med. J.*, **1**, 59.
- BABER, N.S. & LEWIS, J.A. (1981). Post-infarction intervention studies with propranolol and atenolol: problems in design and interpretation. In: *Frontiers in Hypertension Research*. Eds. Laragh, Buhler and Seldin, New York, Springer-Verlag Inc., 485-9.
- BABER, N.S., WAINWRIGHT EVANS, D., HOWITT, G., THOMAS, M., WILSON, G., LEWIS, J.A., DAWES, P.M., HANDLER, K. & TUSON, R. (1980). Multicentre post-infarction trial of propranolol in 49 hospitals in the United Kingdom, Italy and Yugoslavia. *Br. Heart J.*, **44**, 96-100.
- BAKER, R.J. & NELDER, J.A. (1978). *General Linear Interactive Modelling*. Release 3, Oxford: NAG.
- BALCON, R.J., JEWITT, D.E., DAVIES, J.P.H. & ORAM, S. (1966). A controlled trial of propranolol in acute myocardial infarction. *Lancet*, **2**, 918-920.
- BAPTISTA, J. & PIKE, M.C. (1977). Exact two-sided confidence limits for the odds ratio in a 2x2 table. *Applied Statistics*, **26**, 214-220.
- BARBER, J.M., MURPHY, F.M. & MERRETT, J.D. (1967). Clinical trial of propranolol in acute myocardial infarction. *Ulster Med. J.*, **36**, 127-131.
- BARBER, J.M., BOYLE, McC. D., CHATUVERDI, N.C., SINGH, S. & WALSH, M.J. (1976). Practolol in acute myocardial infarction. *Acta Med. Scand.*, **586**, 213-219.
- BETA-BLOCKER HEART ATTACK STUDY GROUP. (1981). The beta-blocker heart attack trial. *JAMA*, **246**, 2073-2074.
- BRIANT, R.B. & NORRIS, R.M. (1970). Alprenolol in acute myocardial infarction: double-blind trial. *New Zealand Med. J.*, **71**, 135-138.
- CLAUSEN, J., FELSBY, M., SCHONAU JORGENSEN, F., LYAGER NIELSEN, B., ROIN, J. & STRANGE, B. (1966). Absence of prophylactic effect of propranolol in myocardial infarction. *Lancet*, **2**, 920-924.
- CORONARY PREVENTION RESEARCH GROUP. (1981). An early intervention secondary prevention study with oxprenolol following myocardial infarction. *European Heart J.*, **2**, 389-393.
- CORONARY PREVENTION RESEARCH GROUP. (1980). Secondary prevention studies with oxprenolol in coronary heart disease. In: Burley, D.M., Birdwood, G.F., eds. *The Clinical Impact of Beta-adrenoceptor Blockade*. 27-28. Ciba Laboratories Ltd.
- EVEMY, K.L. & PENTECOST, B.L. (1978). Intravenous and oral practolol in the acute stages of myocardial infarction. *Europ. J. Cardiol.* **7/5-6**: 391-398.
- FUCELLA, L.M. (1968) Report on the double-blind trial with compound CIBA 39089 (Trasicor) in myocardial infarction. Quoted by Sowton, E. in *Progress in Cardiovascular Diseases*, **10**, 561-574.
- HAMPTON, J.R. (1981). The use of beta-blockers for the reduction of mortality after myocardial infarction. *European Heart J.*, **22**, 259-268.
- HANSTEEN, V., MOINICHEN, E., LORENTSEN, E., ANDERSEN, A., STRØM, O., SØCLOND, K., DYRBEKK, D., RETSEIM, A-M., TROMSDAL, A. KNUDSEN, K., EIKA, C., BAKKEN T. Jnr., SMITH, P. & HOFF, P.J. (1982). One year's treatment with propranolol after myocardial infarction: preliminary report of Norwegian multicentre trial. *Br. Med. J.*, **284**, 155-160.
- HJALMARSON, Å., HERLITZ, J., MALEK, I., RYDEN, L., VEDIN, A., WALDENSTRÖM, A., WEDEL, H., ELIMFELDT, D., HOLMBERG, S., NYBERG, G., SWEDBERG, K., WAAGSTEIN, F., WALDERSTRÖM, J., WILHELMSEN, L. & WILHELMSSON, C. (1981). Effect on mortality of metoprolol in acute myocardial infarction. *Lancet*, **2**, 823-827.
- JOHANSEN, B.W. (1980). A comparative study of cardioselective beta-blockade and diazepam in patient with acute myocardial infarction and tachycardia. *Acta Med. Scand.*, **207**, 47-53.
- KAHLER, R.L., BRILL, S.J. & PERKINS, W.E. (1968). The role of propranolol in the management of acute myocardial infarction. In Kattus, A.A., Ross, G. and Hall, Y.E., eds. *Cardiovascular Beta-Adrenergic Responses*, 213-222. Univ. California Press, Los Angeles: California Press.
- LOMBARDO, M., SELVINI, A., BELLI, C., MOTOLESE, M. & PEDRONI, P. (1979). Beta-blocking therapy in 440 cases of acute myocardial infarction: a double-blind trial with oxprenolol. *Proceedings of Florence International Meeting on Myocardial Infarction*, May 8-12, **2**, 803-807.
- MULTICENTRE INTERNATIONAL STUDY. (1977). Supplementary Report. Reduction in mortality after myocardial infarction with long-term beta-blockade. *Br. Med. J.*, **2**, 419-421.
- MULTICENTRE TRIAL. (1966). Propranolol in acute myocardial infarction. *Lancet*, **2**, 1435-1438.
- NELDER, J.A. & WEDDERBURN, R.M.W. (1972). Generalised linear models. *JRSS Series A*, **135**, 370-384.
- NORRIS, R.M., CAUCHEY, D.E. & SCOTT, P.J. (1968). Trial of propranolol in acute myocardial infarction. *Br. Med. J.*, **2**, 398-400.
- THE NORWEGIAN MULTICENTRE STUDY GROUP. (1981). Timolol-induced reduction in mortality and reinfarction in patients surviving acute myocardial infarction. *New Engl. J. Med.*, **304**, 801-807.
- REYNOLDS, J.L. & WHITLOCK, A.M.L. (1972). Effects of a beta-adrenergic receptor blocker in myocardial infarction for one year from onset. *Br. Heart J.*, **34**, 252-259.
- SLOMAN, G. & STANNARD, M. (1967). Beta-adrenergic blockade and cardiac arrhythmias. *Br. Med. J.*, **4**, 508-512.

- SNOW, P.J.D. (1966). Treatment of acute myocardial infarction with propranolol. *Am. J. Cardiol.*, **18**, 458–462.
- TARONE, R.E. (1981). On summary estimators of relative risk. *J. Chron. Dis.*, **34**, 463–468.
- THOMAS, D.G. & GART, J.J. (1977). A table of exact confidence limits for differences and ratios of two proportions and their odds ratio. *JASA*, **72**, 73–76.
- VEDIN, J.A. & WILHELMSSON, C. (1979). A review of current beta-blocker trials in the world. *Hart Bulletin*, 180–182.
- WILCOX, R.G., ROWLEY, J.M., HAMPTON, J.R., MITCHELL, J.R.A., ROLAND, J.M. & BANKS, D.C. (1980a). Randomised placebo-controlled trial comparing oxprenolol with disopyramide phosphate in immediate treatment of suspected myocardial infarction. *Lancet*, **2**, 765–769.
- WILCOX, R.G., ROLAND, J.M., BANKS, D.C., HAMPTON, J.R. & MITCHELL, J.R.A. (1980). Randomised trial comparing propranolol with atenolol in immediate treatment of suspected myocardial infarction. *Br. Med. J.*, **280**, 885–888.
- WILHELMSSON, C., VEDIN, J.A., TIBBLIN, G., WILHELMSEN, L. & WERKÖ, L. (1974). Reduction of sudden death after myocardial infarction by treatment with alprenolol. *Lancet*, **2**, 1157–1163.
- YUSUF, S., RAMSDALE, D., PETO, R., FURSE, L., BENNETT, D., BRAY, C. & SLEIGHT, P. (1980). Early intravenous atenolol treatment in suspected acute myocardial infarction: preliminary report of a randomized trial. *Lancet*, **2**, 273–276.