DRUG HEPATOTOXICITY

JOHN A. TIMBRELL

Toxicology Unit, Department of Pharmacology, The School of Pharmacy,
29/39 Brunswick Square, London WC1N 1AX

Introduction

The liver is the main site of metabolism for drugs and other exogenous compounds. As most drugs are taken orally the liver is the portal to the tissues for such compounds following absorption from the gastrointestinal tract (G.I. tract). The liver is, therefore, a vulnerable organ, being exposed to both the parent drug carried from the G.I. tract via the portal vein and to any metabolites produced which then enter the systemic circulation via the hepatic vein. However, despite this vulnerability, the liver is not the major target for adverse drug reactions, only about 9.5% of these involve the liver (Davis & Williams, 1977). This may partly result from the large reserve functional capacity of the liver and the protective mechanisms present. However, although drug-induced hepatic damage may not be particularly common in general patient populations the case fatality rate is often high, and the severity of drug-induced hepatic injury is such that drugs are a major cause of hepatic failure (Zimmerman, 1978). The case fatality rate from halothane-induced hepatic failure may be as high as 50% (Inman & Mushin, 1974). In particular populations of patients, such as in geriatrics, hepatic injury may be more common, whereas in children it seems to be less frequent than in the general population.

Drugs and other exogenous compounds may affect the liver in various ways. Thus hepatic enzymes may be induced, protein synthesis and blood flow increased by certain barbiturates, enzymes may be inhibited such as by disulphiram, and bile secretion may be affected, as it is by rifampicin. Interference with haem biosynthesis may give rise to porphyria, as stimulated by alkyl substituted barbiturates. Some of these effects may be involved in hepatotoxicity or be factors altering sensitivity to hepatotoxic drugs.

In this review the principle types of drug hepatotoxicity are discussed. This includes (a) the features of hepatotoxic reactions in man, (b) the relevant characteristics of the drugs concerned and of the susceptible patients, (c) the mechanisms underlying the hepatotoxic effects where these are known. The animal data relevant to this have also been included to highlight the importance of such data in the anticipation of hepatotoxicity of new compounds. Such new medicinal agents clearly need to be evaluated for hepatotoxicity both in experimental animals and man.

The examples discussed indicate the variety of mechanisms underlying drug hepatotoxicity and the various manifestations that may occur. However detection and assessment of hepatotoxicity relies mainly on standard clinical chemistry and may not distinguish between drug induced hepatotoxicity and that due to other causes and may not always distinguish between the types of hepatotoxic response.

Knowledge and appreciation of the major types of drug hepatotoxicity therefore may aid interpretation of clinical and biochemical data with regard to new medicinal agents.

Types of hepatic damage

Drug-induced liver injury covers a variety of types, and includes most of the clinical and pathological expressions of liver damage (Zimmerman, 1981).

Various ways of classifying the types of liver damage have been used and Table 1 is a synthesis of these showing the spectrum of injury. However, drugs may produce more than one type of lesion and there is often overlap between them depending on the circumstances. Zimmerman (1978) has highlighted particular facets of drug-induced liver injury which are shown in Table 2.

It should also be noted that some of these lesions are dose-dependent and predictable whereas others are independent of dose and unpredictable. It is clear that some drug induced-liver injuries require metabolism of the drug whereas in other cases this is not so. The expression of liver damage depends on

Table 1 Types of hepatic damage

1. Interference with bilirubin uptake, excretion and conjugation.
2. Cytotoxic injury.
3. Cholestatic injury.
4. Mixed cholestatic and cytotoxic injury.
5. Fatty liver (steatosis).
6. Cirrhosis.
7. Phospholipidosis.
8. Liver tumours.
9. Vascular lesions.
10. Chronic active hepatitis.
11. Subacute hepatic necrosis.

Table 2 Facets of drug-induced hepatic injury

1. Character of injury.
 - a: Cytotoxic (Hepatocellular)
 - b: Cholestatic—Hepatocanalicular
—Canalicular
 - c: Mixed types.
2. Mechanism of toxicity.
 - a: Intrinsic—Direct
—Indirect
 - b: Idiosyncratic—Hypersensitivity
—Metabolic
3. Circumstances of exposure.
 - a: Iatrogenic—Therapeutic doses
 - b: Toxicologic—Overdoses

various factors as well as the size and duration of dosing. For example, hepatotoxicity may be influenced by genetic factors (Mitchell *et al.*, 1975a), by diet, (Angeli-Greaves & McLean, 1979) and the concomitant administration of other drugs (Wright & Prescott, 1973). Tissue type and other immunological factors may also be involved as is the case with other drug-induced diseases (Batchelor *et al.*, 1980).

Therefore, drug toxicity may range from a predictable acute response after an overdose, to an idiosyncratic reaction following chronic therapeutic doses. Any classification must attempt to take all of these factors into account. Hepatic damage may be recognised histologically in biopsy samples but normally hepatic dysfunction is detected biochemically (Zimmerman & Seef, 1970). The serum transaminases, aspartate aminotransferase (AST, SGOT) and alanine aminotransferase (ALT, SGPT) may indicate hepatic damage when at raised levels.

Elevated serum alkaline phosphatase levels generally indicate cholestatic lesions, particularly damage to the bile duct. When plasma levels of conjugated and unconjugated bilirubin are both elevated hepatic damage is normally indicated. However, the level of the unconjugated form may be raised without evident hepatic lesions, following the ingestion of certain drugs (see below). Various other biochemical tests are available for detecting the different forms of liver damage (Zimmerman & Seef, 1970) but the ones described are the most commonly used in human studies. However there are marked species differences in serum biochemistry following tissue damage (Woodman, 1981).

I shall now briefly discuss the types of liver injury as indicated in Table 1.

1. Interference with bilirubin transport and conjugation

This may be viewed as a variant of cholestatic toxicity (Zimmerman, 1978). A number of drugs cause this

type of effect. Rifampicin (Capelle *et al.*, 1974), flavaspidic acid (Kenright & Levi, 1974) and novobiocin (Hargreaves & Lathe, 1963) may all interfere with bilirubin transport giving rise to hyperbilirubinaemia but no other lesion.

2. Cytotoxic injury

This refers to overt damage to the parenchyma and is a much more serious effect than the above. It may be manifested as necrosis or steatosis (fatty liver). Necrosis may be zonal, such as that due to paracetamol (James *et al.*, 1975) and halothane (Moult & Sherlock, 1975) or diffuse, such as that caused by isoniazid and iproniazid (Mitchell *et al.*, 1976). Steatosis, which is similar to Reye's syndrome, may be microvesicular as after tetracycline (Kunelis *et al.*, 1965), or macrovesicular as caused by methotrexate (Dahl *et al.*, 1971).

3. Cholestatic injury

This type involves arrested bile flow and jaundice and may appear similar to biliary obstruction. It is less serious than cytotoxic injury, with a lower fatality rate. Cholestatic injury may involve bile stasis alone (canalicular) such as that caused by anabolic and contraceptive steroids (Zimmerman, 1974), or it may be more serious and involve both bile stasis and inflammation (hepatocanalicular) such as that caused by chlorpromazine (Ishak & Irey 1972) or erythromycin estolate (Havens, 1962).

4. Mixed cytotoxic and cholestatic injury

Cytotoxic hepatic damage may sometimes be combined with cholestasis, for instance occasionally seen after *p*-aminosalicylic acid therapy, (Zimmerman, 1974).

5. Fatty liver

Fatty liver (steatosis) may be considered as a type of cytotoxic injury (see 2), but it may also be a form of chronic hepatic damage.

6. Cirrhosis

Macronodular cirrhosis may directly follow acute hepatic damage and primary biliary cirrhosis may result from cholestatic jaundice.

7. Phospholipidosis

This may result from the use of drugs such as Coralgil, (4,4'-diethylaminoethoxyhexestrol dihydrochloride), and is characterised by hepatocytes full of lipid (Lullmann *et al.*, 1975). This lesion may develop into cirrhosis.

8. Liver tumours

Neoplastic lesions may result from drug use. Benign, liver cell adenomas have been associated with the use of contraceptive steroids (Neuberger *et al.*, 1981a) but liver cell carcinoma associated with these drugs is more uncertain (Zimmerman, 1978). Other suspect compounds, capable of producing malignant liver tumours in animals, are the anabolic steroids, griseofulvin, hycanthone and isoniazid (Zimmerman, 1978).

9. Vascular lesions

Hepatic vein occlusion, such as that due to the thrombogenic effect of contraceptive steroids, may result in hepatic damage. Centrilobular necrosis and fibrosis may follow and result in a congestive cirrhosis. Hepatic vein thrombosis has been described in users of the contraceptive pill (Zimmerman, 1978).

10. Chronic active hepatitis

This is a progressive necroinflammatory liver disease which may have many causes including drugs. This syndrome of portal inflammation, periportal necrosis, fibrosis and sometimes cirrhosis may result from prolonged ingestion of a hepatotoxin following an initial lesion. Oxyphenisatin has been implicated in this type of liver injury as have isoniazid, α -methyldopa, nitrofurantoin and dantrolene (Zimmerman, 1981).

11. Subacute hepatic necrosis

This syndrome consists of progressive, overt serious liver disease with jaundice and eventual cirrhosis. It develops more slowly than acute cytotoxic liver damage but more rapidly than chronic active hepatitis. α -Methyldopa may cause this type of lesion (Zimmerman, 1978).

Mechanisms of drug-induced liver damage

Certain drugs will produce predictable liver damage in the majority of cases, such as after overdoses, for instance. Others however will cause liver injury only rarely and unpredictably. There is thus a whole spectrum varying between these extremes.

In some cases the mechanism may involve the parent compound, in others a metabolite may be responsible. Direct cytotoxicity is known to be the underlying cause of liver damage in certain cases whereas in others immunological mechanisms or even a mixture of both cytotoxicity and immunogenicity may be involved.

The various mechanisms will now be discussed using specific examples, according to the type of injury.

1. Interference with bilirubin transport and conjugation

A number of drugs interfere with bilirubin transport and lead to elevated plasma bilirubin or hyperbilirubinaemia. Bilirubin normally diffuses across the hepatocyte cell membrane and is carried across the cell by a carrier protein such as the Y and Z proteins. Flavaspidic acid interferes with the uptake of bilirubin and its binding to the Z carrier protein (Kenright & Levi, 1974). This results in an elevated level of unconjugated bilirubin in the plasma.

Novobiocin inhibits UDP glucuronosyl transferase and may lead to elevated plasma levels of unconjugated bilirubin especially in neonates (Cox *et al.*, 1959).

Rifampicin. Rifampicin, the antibiotic used in the treatment of tuberculosis inhibits both uptake and excretion of bilirubin in a dose related manner, giving rise to elevated plasma levels of both conjugated and unconjugated bilirubin. This is due to blockade of uptake at the plasma membrane of the hepatocyte (Capelle *et al.*, 1972; Kenright & Levi, 1974).

As well as causing hyperbilirubinaemia, rifampicin may also cause overt hepatic damage (Zimmerman, 1978), although in many cases rifampicin is taken with other potentially hepatotoxic drugs such as isoniazid. Consequently the true incidence is difficult to establish, but rifampicin alone has been clearly implicated in some cases of hepatic damage. There is no indication of factors which might predispose to rifampicin induced injury, although preexisting liver disease has been suggested.

The lesion is characterised as hepatocellular with centrilobular necrosis which may be accompanied by cholestasis. Serum transaminases may be elevated but are not necessarily excessively so.

The mechanism of this hepatic damage is unknown at present, but clearly is not related to the effects on bilirubin transport which is dose related and predictable whereas the hepatocellular damage is idiosyncratic and not dose related (Zimmerman, 1978). It may be due to a hypersensitivity reaction, but systemic effects such as fever and rash are not common. There is no evidence for or against a toxic metabolite(s) being produced. Other effects of rifampicin include interference with DNA synthesis and the ability to induce hepatic microsomal enzymes and cause proliferation of the smooth endoplasmic reticulum. These effects, however, are not thought to be responsible for hepatic damage, although induction of the microsomal enzymes may well be a factor in the hepatotoxicity of rifampicin-isoniazid combinations (see below).

2. Cytotoxic injury

Direct, overt damage to hepatic parenchyma may be caused by a number of drugs. It may have a variety of underlying mechanisms, however, and several examples will be examined.

Paracetamol This minor analgesic causes predictable centrilobular hepatic necrosis in both experimental animals and man after overdoses (Boyd & Berezky, 1966; James *et al.*, 1975). Overdosing with paracetamol is an increasingly common means of suicide in Britain and consequently paracetamol has become one of the most common causes of hepatic necrosis (Prescott *et al.*, 1971). Consequently it is well characterised and the mechanism partially understood. Doses of 10 g may lead to liver damage, and doses greater than 15 g may be sufficient for fatal hepatic damage.

Liver damage may be detected as raised serum transaminases, AST and ALT (SGOT and SGPT) which may reach levels of 5000 i.u./l (Prescott *et al.*, 1971). However, bilirubin levels may be only moderately elevated.

Clinically, the course following overdosage consists of 3 phases. The first, immediately following poisoning, is characterised by nausea, vomiting, anorexia and shock. The second phase, with apparent subsidence of severe symptoms, is when raised serum enzymes may be detected. The third, final phase, is marked by the appearance of jaundice, with overt hepatic damage which may progress to fulminant hepatic failure. Myocardial injury and renal failure may also develop (Zimmerman, 1978).

The liver damage is predictable and is due to direct cytotoxicity of a metabolite as indicated by extensive studies on both experimental animals (Mitchell *et al.*, 1975b) and man (Mitchell *et al.*, 1974). Paracetamol is metabolised by three pathways, (Figure 1), two of which are conjugation reactions, and remove the major portion of the drug rapidly. A minor pathway accounting for about 5% of the dose, catalysed by the microsomal enzymes yields a reactive metabolite, thought to be *N*-acetyl-*p*-benzoquinone imine. Following normal doses this is detoxified by conjugation, both chemical and enzyme catalysed, with the ubiquitous tripeptide glutathione, and excreted as the *N*-acetylcysteine derivative in the urine. After overdoses however, the amount of reactive metabolite is sufficient to deplete the available hepatic glutathione. The reactive metabolite then reacts covalently with cellular macromolecules as revealed by measurement of covalent binding of radiolabelled paracetamol to hepatic protein and autoradiographic studies which indicate binding primarily in necrotic areas (Mitchell *et al.*, 1975b). The mechanism by which this binding causes cellular damage is unknown, but reducing the binding by supplying alternative

Figure 1 Metabolism of paracetamol showing proposed metabolic activation and its involvement in the toxicity.

nucleophiles and increasing glutathione levels protects against the hepatic injury (Mitchell *et al.*, 1974). These findings suggested a means of treatment, now successfully used (Prescott, 1981), which involves intravenous injection of *N*-acetylcysteine which, provided less than 10 h after overdosing has occurred, will generally minimize the hepatic damage. The likelihood of hepatic damage occurring can be estimated from the plasma level of paracetamol. Measurement of this level at 4 and 15 h after ingestion may be compared with a nomogram (Prescott, 1981). This reflects the ability of the liver to metabolise the drug and hence if this is reduced the half-life of paracetamol will be extended. Depending on the half-life, expected liver damage may be estimated as mild or absent, severe to mild and very severe.

Factors which may affect paracetamol hepatotoxicity include the activity of the hepatic microsomal enzymes and the concomitant administration of other drugs. Prior ingestion of phenobarbitone is known to increase metabolism to the toxic metabolite in man (Mitchell *et al.*, 1974). This presumably reflects induction of the microsomal enzymes. There is some evidence that patients previously using barbiturates have an increased risk of liver damage (Wright & Prescott, 1973).

Isoniazid and iproniazid

Both of these substituted hydrazine drugs may produce a hepatocellular, cytotoxic type of damage. However, this often resembles chronic active hepatitis or sub-acute hepatic necrosis and will be discussed below under these categories.

Halothane

This most widely used anaesthetic is now accepted as causing hepatic injury (Trowell *et al.*, 1975). The majority of cases of halothane induced injury (more than 80%) occur in patients after more than one exposure. Furthermore the severity of the injury is greater after multiple exposures as reflected in the increased case fatality rate. This may be as high as 63% after 4 or more exposures (Inman & Mushin, 1974). The actual incidence is hard to quantify and depends on circumstances of exposure as already indicated. For instance in one study 4 out of 18 patients (22%) had biochemical evidence of liver damage, subsequently confirmed by biopsy in 2 patients, following repeated halothane anaesthesia (Trowell *et al.*, 1975). The generally accepted incidence is one in several thousand but despite this low incidence, because of the large number of patients exposed, halothane induced injury accounts for perhaps 16–25% of cases of hepatic failure (Zimmerman, 1978). As already mentioned multiple exposure is a major factor which may predispose the patient to liver injury, particularly if reexposure occurs within 3 months. Obese patients and females seem more susceptible but children and young adults less so.

A clear syndrome can be ascribed to halothane induced liver injury (Zimmerman, 1978). There is normally a latent period between exposure to the anaesthetic and appearance of liver damage. The syndrome may include fever and myalgia but arthralgia and rash are rarer. Gastrointestinal symptoms such as anorexia, nausea and abdominal distress may also occur. These symptoms are followed by overt jaundice.

The hepatic damage is a zonal necrosis usually centrilobular with a sharp demarcation between necrotic areas and surrounding parenchyma, but may occasionally be massive. This is in contrast to viral hepatitis. Its biochemical features are similar to other instances of cytotoxicity, with markedly elevated serum levels of AST(SGOT) and ALT(SGPT). Serum bilirubin levels may be elevated, the extent reflecting the severity, and alkaline phosphatase may show a moderate (2–3 times) elevation. Recovery is generally complete and only rarely are there chronic changes.

Halothane-induced cytotoxicity is not predictable although it can be reproduced in experimental animals under certain conditions (Sipes *et al.*, 1981). However, mechanisms have been proposed which make this a particularly interesting and valuable drug toxicity to discuss.

Although isolated reports had appeared indicating that halothane-induced liver damage could be reproduced in experimental animals if these were pre-treated with phenobarbitone the lesion was clearly not readily reproducible. Halothane was

shown to be metabolised by a number of pathways and although the major metabolite, trifluoroacetic acid was not hepatotoxic, the influence of phenobarbitone pre-treatment on hepatotoxicity indicated that a metabolite was probably responsible (Sipes *et al.*, 1981). It became clear that halothane was metabolised by both oxidative and reductive pathways (Figure 2a,b). Furthermore, covalent binding of a halothane metabolite to cellular macromolecules was detected and this was enhanced under anaerobic conditions. Exposure of rats to halothane under conditions of low oxygen concentration increased serum fluoride levels (Sipes *et al.*, 1981). These findings led to the development of a model for halothane hepatotoxicity, involving (a) low oxygen concentrations, (b) the microsomal enzymes induced by pre-treatment with phenobarbitone, (c) the use of male rats, and (d) exposure to 1% halothane for 2 h (Sipes *et al.*, 1981). Increased microsomal metabolism and low oxygen tension direct metabolism via reductive pathways (Figure 2b). The lesion is dependent on dose, hepatic cytochrome P₄₅₀ content, oxygen concentration and duration of exposure. Thus at normal levels of oxygen and cytochrome P₄₅₀ activity, little if any liver injury occurs. It is suggested that reductive metabolism may produce a reactive metabolite, possibly a free radical, which binds covalently to liver macromolecules and leads to hepatic damage. There is a correlation between the hepatic covalent binding of a fluorine containing moiety and serum fluoride levels in humans (Sipes *et al.*, 1981). The reductive metabolism of halothane is also known to take place in humans, as the products, trifluorochloroethane and difluorochloroethylene, have been detected in the expired air (Maiorino *et al.*, 1979). An alternative or additional mechanism involves lipid peroxidation resulting from the formation of a reactive radical intermediate which may also covalently bind to cellular macromolecules (Figure 2). Lipid peroxidation has been detected after halothane anaesthesia in

Figure 2 Metabolism of halothane and proposed metabolic activation to a reactive metabolite.

both experimental animals and humans (Hempel *et al.*, 1981).

Although these data ascribe halothane hepatotoxicity to direct cytotoxicity of a reactive metabolite, the fact that serious liver damage is a rare event and occurs more commonly after multiple exposures indicates that other factors are at work. Clearly if reductive metabolism takes place in all patients, then the hepatic injury should be more common. It is likely, therefore, that this is a multifactorial problem. The susceptible patient will perhaps have a particularly high level of one particular form of cytochrome P₄₅₀, and the conditions of anaesthesia may be such as to result in a low oxygen tension.

However, it has been suggested that there are two forms of hepatic injury from halothane (Davis *et al.*, 1981). There is a mild form with slightly raised serum transaminases and perhaps mild necrosis, which occurs in up to 20% of patients, sometimes following one, but usually multiple, exposures (Trowell *et al.*, 1975). Then there is a more serious form occurring in possibly one in 22–35,000 patients which leads to massive hepatic necrosis and death in some cases (Strunin, 1977). It is suggested that this latter severe form has an immunological basis, whereas the more mild form is due to the direct cytotoxicity reproducible in experimental animals as described above. The immunological basis of the severe form is suggested by the occurrence of fever, rash, myalgia, eosinophilia and high titres of liver and kidney microsomal antibodies and high titres of antibodies to normal liver cell membrane components (Davis *et al.*, 1981).

Experimental studies revealed that human patients suffering the severe form of halothane induced hepatic injury had circulating antibodies against hepatocytes from rabbits treated with halothane. It was demonstrated that the antibodies were specifically against the hepatocyte membrane (Davis *et al.*, 1981). Furthermore, the lymphocytes from such patients were cytotoxic to the rabbit hepatocytes *in vitro*. The antibody, therefore, rendered the hepatocyte susceptible to cytotoxicity from normal lymphocytes. Further studies indicated that the immunological toxicity was the result of formation of an antigen via the oxidative rather than reductive pathway (Neuberger *et al.*, 1981b) (Figure 3).

The oxidative metabolic pathway predominates, however, and, therefore, other factors such as an abnormality in the immune system are presumably involved in the rare development of severe hepatic injury.

Thus the reductive metabolic pathway may be responsible for direct cytotoxicity and minor hepatic damage, and the oxidative pathway seems to be involved in the severe hepatic injury. It may be that various contributions of the two mechanisms give rise to the observed spectrum of clinical features, sometimes including both hepatic necrosis and peripheral lymphocytic infiltration for instance.

Figure 3 Postulated mechanisms for halothane hepatotoxicity.

3. Cholestatic injury

This type of hepatic toxicity may be seen with a number of drugs, and may be either the mild, canalicular form or the more severe hepatocanalicular variety.

Chlorpromazine This major tranquilizer is an important cause of drug-induced jaundice. The incidence is estimated at between 0.5 and 1% of recipients of therapeutic doses (Zimmerman, 1974) although up to 50% of patients receiving large doses may have minor abnormalities of liver function (Ishak & Irey, 1972). Jaundice commonly develops after 3 weeks of therapy, with often the development of fever, itching, abdominal pain, nausea and anorexia. This is often similar to extrahepatic obstructive jaundice, with elevated serum cholesterol and alkaline phosphatase levels more than 4 times normal. AST(SGOT) and ALT(SGPT) levels may also be moderately raised (Zimmerman, 1978).

The lesion is generally one of cholestasis particularly in the centrilobular region with bilirubin casts in distended canaliculi and bilirubin staining of hepatocytes and Kupfer cells. There may also be scattered areas of focal necrosis. An inflammatory response may occur in the portal areas.

Unlike cytotoxic hepatic injury, cholestatic damage has a very low fatality rate and usually subsides within 4 to 8 weeks in 50% of patients. Occasionally a syndrome similar to primary biliary cirrhosis has developed following prolonged cholestasis.

Certain elements of chlorpromazine-induced liver damage suggest a hypersensitivity reaction (Zimmerman, 1978). Thus symptoms appear 1–5 weeks after dosing has started, it occurs in less than 1% of recipients, and eosinophilia and an inflammatory response commonly occur. Furthermore, anti-nuclear antibodies can be detected in 40% of recipients, although the occurrence of these does not seem to be directly related to the hepatotoxicity.

However, chlorpromazine also seems to have intrinsic toxicity. *In vitro* chlorpromazine is cytotoxic to isolated cells in culture and liver slices in a dose related fashion (Zimmerman, 1978). Chlorpromazine is an amphipathic, cationic tertiary amine detergent, and effects on membranes are to be expected. It is highly concentrated in bile during biliary excretion and the concentration in bile may exceed that intrinsically toxic *in vitro* to cell membranes (Boyer, 1981; Ros *et al.*, 1979).

Chlorpromazine will intercalate with the inner half of the membrane lipid bilayer and interferes with membrane calcium transport (Boyer, 1981). Furthermore it inhibits Na^+, K^+ -ATPase and impairs Mg^{++} -ATPase activity. This inhibition is particularly associated with the ring hydroxylated metabolite, 7,8-dihydroxychlorpromazine (Boyer, 1981). Chlorpromazine also interacts with actin polymerisation, and hence interferes with the cellular cytoskeleton and microfilaments. Thus blebbing of the sinusoidal membrane is seen in the isolated perfused liver when concentrations of chlorpromazine are used which inhibit bile secretion (Boyer, 1981). Bile acid independent and dependent canalicular secretions are also inhibited, as may be the secretion of cholesterol and biliary phospholipid.

Chlorpromazine clearly has a number of effects on membranes and on transport and secretory mechanisms and it is probable that the primary site of injury is the bile ductules. The actual mechanism however remains to be discovered.

Anabolic steroids Steroids with an alkyl group at C_{17} (Figure 4) such as methyltestosterone may give rise to mild hepatotoxicity (Klatskin, 1975). This is normally a cholestatic injury generally without parenchymal damage (canalicular type). Thus alkaline phosphatase is only slightly raised (2 times normal) as are transaminases (AST) but jaundice may be marked, with elevated bilirubin levels (30 mg/100ml). The majority of recipients of therapeutic doses show some hepatic dysfunction, but only a few develop jaundice, usually after 2–5 months of dosing. The degree of injury and rate of development seem to be dependent on dose. Malaise, anorexia and nausea may be clinical features. Only occasionally is there parenchymal injury and inflammation is slight, arrested bile flow being the major change.

Figure 4 Basic structure of steroids related to testosterone. For testosterone $\text{R} = \text{H}$.

There are precise structural requirements for the steroids which produce this hepatic damage. Thus a C_{17} -methyl substituent is more active than an ethyl or vinyl group (DeLorimer *et al.*, 1965; Lennon, 1966). A keto group at C_3 confers greater activity than a hydroxyl group and saturation of the A ring of the steroid reduces its ability to cause hepatic dysfunction (Aldercreutz & Tenhunen, 1970; Lennon, 1966).

Although the exact mechanism of hepatotoxicity is obscure, anabolic steroids clearly cause a dose-related blockage of bile secretion (Zimmerman, 1978; Aldercreutz & Tenhunen, 1970). The excretion of conjugated bilirubin into the canaliculus is selectively impaired, and this may reflect a membrane effect. Thus the hepatocyte membrane Na^+, K^+ -ATPase is inhibited and permeability is changed (Wheeler, 1975). Anabolic steroids also interfere with the bile salt dependent and independent fraction of bile flow (Wheeler, 1975).

Clearly there is an effect on the excretory/secretory mechanisms and this can be seen as an impairment of bromosulphophthalein excretion which develops fairly rapidly following start of treatment with anabolic steroids (Zimmerman, 1978). This hepatic effect is not idiosyncratic as dysfunction may be detected in many recipients and may also be reproduced in experimental animals (Zimmerman, 1973).

4. Mixed cytotoxic/cholestatic injury

This type of liver injury covers damage with varying proportions of cytotoxic and cholestatic involvement. For example chlorpromazine may cause mixed hepatocanalicular jaundice with parenchymal injury as well as cholestasis. Conversely, *p*-aminosalicylic acid may cause mixed hepatocellular liver injury.

5. Fatty liver (steatosis)

Tetracycline This antibiotic may occasionally cause fatty liver after large (1.5 g/day) intravenous doses (Zimmerman, 1978; Davis, 1968; Dowling & Lepper, 1964). This toxic effect is very rare after oral doses, and occurs more commonly in females than males.

After 4–10 days of tetracycline administration nausea, vomiting and abdominal pain may be clinical features of a syndrome which may end in coma and death (Zimmerman, 1978; Davis, 1968). The prognosis is serious. Biochemically the changes may only be slight, with moderate elevations of AST(SGOT), usually less than 500 i.u.l, and smaller elevations of ALT(SGPT). Alkaline phosphatase levels are generally close to normal values and bilirubin values are usually only moderately elevated, being less than 5 mg/dl in 50% of patients (Zimmerman, 1978). Hypoglycaemia may occur (Dowling & Lepper, 1964) and serum amylase levels are elevated in 75% of cases, sometimes markedly (Schultz *et al.*, 1963).

Histologically hepatocytes are packed with small fat droplets. This microvesicular steatosis occurs initially in the centrilobular area. There is little inflammation, necrosis or cholestasis (Zimmerman, 1978).

This toxic effect of tetracycline is direct, predictable and dose dependent, and can be reproduced in experimental animals. The major effect seems to be inhibition of transport of lipid out of the hepatocyte, which can be detected within 30 min of dosing in experimental animals (Breen *et al.*, 1975). This effect may well be due to the inhibition of protein synthesis caused by tetracycline which will inhibit the production of the apolipoprotein complex involved in transport of the very low density lipoprotein (VLDL) out of the hepatocyte (Breen *et al.*, 1972, 1975). This is similar to the mechanism proposed for carbon tetrachloride and ethionine-induced fatty liver (Zimmerman, 1978). Alternative or additional mechanisms may involve decreased fatty acid oxidation, increased triglyceride uptake or increased fatty acid uptake (Breen *et al.*, 1975) (Figure 5).

Figure 5 Postulated mechanisms for tetracycline-induced fatty liver.

6. Chronic active hepatitis, cirrhosis and sub-acute necrosis

Chronic active hepatitis, sometimes associated with cirrhosis is associated with the use of a number of drugs such as oxyphenisatin α -methyl dopa, nitrofurantoin and isoniazid (Zimmerman, 1978). Isoniazid has been chosen as the example because the mechanism is at least partially understood.

Isoniazid Long term administration of the anti-tubercular drug isoniazid leads to hepatic dysfunction in a significant proportion of recipients (10–20%). This is usually mild and subclinical and subsides despite continued therapy (Mitchell *et al.*, 1976). However, some 0.1–1% of patients develop severe hepatic injury (Black *et al.*, 1975). Although this is

generally acute hepatocellular damage, in about 10% of cases chronic active hepatitis may develop with or without cirrhosis. Sub-acute necrosis may also occur following isoniazid (Zimmerman, 1978).

The syndrome of isoniazid-induced hepatic damage is similar to viral hepatitis with anorexia, fatigue, nausea and vomiting. Fever and rash are uncommon (Mitchell *et al.*, 1976). The syndrome develops typically 2 months, but in some cases up to 12 months after starting treatment (Mitchell *et al.*, 1976; Black *et al.*, 1975). AST(SGOT) and ALT(SGPT) values are elevated, (up to 10–20 times normal), but alkaline phosphatase levels are generally only moderately raised, except in about 10% of patients when mixed hepatocellular jaundice occurs, giving 4 times normal values for alkaline phosphatase (Black *et al.*, 1975). Histologically necrosis ranges from focal to massive, and cholestasis may at times be very significant. Serum bilirubin may therefore also be raised. In those cases when chronic active hepatitis, and necrosis occur, degenerative changes in periportal hepatocytes may be observed with fibrosis and inflammation in this region of the lobule (Mitchell *et al.*, 1976).

There are a number of predisposing factors for isoniazid-induced hepatic damage. Overt hepatic injury is rare in patients below the age of 20 and increases to 3% in patients over 50 (Mitchell *et al.*, 1976). Preexisting liver dysfunction, such as alcoholic cirrhosis, also increases susceptibility (Gronhagen-Riska *et al.*, 1978). Data from a large scale study in the U.S.A. has revealed that oriental males are more susceptible to isoniazid induced hepatic disease than white males, whereas black males are markedly less susceptible (Black *et al.*, 1975). There is no such ethnic difference between females, hence oriental males seem more, and black males less susceptible than females of the respective races (Kopanoff *et al.*, 1978). Orientals overall had the highest incidence over blacks and whites. However, black females showed a higher fatality rate than other groups in one study (Black *et al.*, 1975), as did elderly white females in another study (Gronhagen-Riska *et al.*, 1978). Finally, a factor around which some controversy exists is the acetylase phenotype. Isoniazid is metabolised by several routes, acetylation being a major one (Figure 6). There are well known genetic (Evans, 1968) and ethnic (Lunde *et al.*, 1977) effects on this reaction. Thus human subjects may be divided into rapid and slow acetylators depending on the amount of acetylated product excreted. American blacks and whites each comprise approximately 50% rapid acetylators whereas orientals are mainly (90%) rapid acetylators. The greater susceptibility of orientals to isoniazid-induced hepatic damage (Black *et al.*, 1975) first suggested a possible relationship with the acetylase phenotype. A retrospective study carried out in the U.S.A. (Mitchell *et al.*, 1975a) of non-

oriental patients, suffering overt hepatic damage following prophylaxis with isoniazid alone, revealed a preponderance of rapid acetylators (88%) when the expected incidence was approximately 50%. This apparently increased susceptibility in those of the rapid acetylator phenotype has not been found in other studies. Indeed, discrepancies abound, with some studies revealing a greater susceptibility in slow acetylators, whereas others have shown no difference (Girling, 1978). More recently, a study revealed that slow acetylators may be more susceptible to the more severe form of hepatic damage, both phenotypes being equally susceptible to the milder form (Gronhagen-Riska *et al.*, 1978).

However, a number of relevant points should be noted regarding these discrepancies. All the studies except that showing a greater susceptibility in rapid acetylators (Mitchell *et al.*, 1975a) have been carried out on patients receiving a mixture of anti-tubercular drugs. All the studies, except those by Mitchell *et al.* (1975a) and Gronhagen-Riska *et al.* (1978) did not differentiate between the mild and severe forms of liver damage. These two factors may be causes of the discrepancy as clearly other anti-tubercular drugs such as rifampicin may well be hepatotoxic, and there seem to be two forms of isoniazid hepatic damage, sub-clinical and overt.

Isoniazid hepatotoxicity occasionally has features suggesting an allergic response with fever, eosinophilia, rash and immediate response to challenge (Black *et al.*, 1975). Although anti-nuclear antibodies may be detected (Mitchell *et al.*, 1975c) these do not correlate with hepatic injury. The majority of cases of isoniazid induced liver damage therefore do not conform to the allergic type of mechanism. Furthermore minor hepatic dysfunction is detected in 10–20% of patients, hardly an idiosyncratic, hypersensitivity

reaction. Also isoniazid hepatotoxicity may be reproduced in experimental animals (Mitchell *et al.*, 1976).

The mechanism of isoniazid induced hepatic damage involves production of a toxic metabolite. The findings of Mitchell *et al.* (1975a) suggested that rapid acetylation might be a predisposing factor as it was reasoned that this would produce more of the metabolite acetylisoniazid and hence more acetylhydrazine (Figure 6). Acetylisoniazid and especially acetylhydrazine are extremely hepatotoxic causing centrilobular hepatic necrosis, in experimental animals in which the microsomal enzymes are induced by phenobarbitone (Mitchell *et al.*, 1976; Bahri *et al.*, 1981).

The hepatotoxicity of acetylisoniazid depends on its metabolism to acetylhydrazine. This metabolite in turn is metabolically activated by the microsomal enzymes to a reactive acylating species which reacts covalently with liver protein (Timbrell *et al.*, 1980).

The role of the acetylator phenotype in this is complex, as studies in human volunteers indicated that although rapid acetylators produced more potentially toxic acetylhydrazine, this was then further acetylated to non-toxic diacetylhydrazine (Timbrell *et al.*, 1977). The relative activities of several pathways are therefore crucial and the eventual toxicity will depend on a number of factors. One such factor may be the nature of the microsomal enzymes, as animal studies have suggested that a specific form of cytochrome P-450 is required for the toxic metabolic activation (Fleet & Timbrell, unpublished observations). There are, however, still a number of unanswered questions relating to isoniazid hepatotoxicity. Acetylhydrazine is a metabolite in all human individuals (Timbrell *et al.*, 1977) and yet only a few develop hepatic damage.

Figure 6 Metabolism of isoniazid showing proposed metabolic activation and its involvement in the toxicity.

Perhaps only these few possess sufficient of the enzyme or isoenzyme required for the metabolic activation.

Certain similarities with halothane-induced hepatic damage suggest that direct cytotoxicity may be the result of a toxic metabolite, whereas the much less frequent severe hepatic damage might involve a different, possibly allergic mechanism or perhaps a mixture of both mechanisms. However, there is no doubt that isoniazid is metabolised in man to metabolites which are directly hepatotoxic in experimental animals.

The relatively high incidence of hepatic dysfunction associated with isoniazid has curtailed its prophylactic use in certain countries. Although its use against active tuberculosis is justified, awareness of the problems should allow reduction in risk. Thus monitoring of ALT(SGPT) has been suggested (Zimmerman, 1978) as 10% of patients may develop jaundice without prior symptoms. The case fatality rate for such overt hepatic damage may be as high as 64% when this develops more than 2 months after the start of treatment and bilirubin values are greater than 20 mg/100ml (Black *et al.*, 1975).

7. Phospholipidosis

This syndrome may be caused by a number of different drugs, and various organs may be affected. Hepatic phospholipidosis has been caused by the drug Coralgil, a coronary dilator, in Japan (Lullmann *et al.*, 1975).

The features of this form of hepatic damage are an accumulation of phospholipids in hepatocytes, bile duct proliferation and inflammation in the portal area. AST(SGOT) and ALT(SGPT) values may be elevated. The lesion may progress to liver cirrhosis and have a fatal outcome (Lullmann *et al.*, 1975). The mechanism is thought to involve the formation of complexes between lipid micelles or liposomes, and the drug in question (Lullmann & Lullmann-Rauch, 1977). Compounds such as Coralgil which may cause phospholipidosis are amphiphilic cations, having both hydrophobic and hydrophilic parts to the positively charged molecule. The interaction between phospholipids and the drug is believed to alter the surface charge of the phospholipid micelle or liposome in such a way that the ability of phospholipases to break them down is impaired (Lullmann & Lullmann-Rauch, 1977). Consequently the phospholipids accumulate in the cell and are stored in the lysosomes forming lamellated and sometimes crystalloid inclusions

in the liver and other organs and tissues (Lullmann *et al.*, 1975; Lullmann & Lullmann-Rauch, 1977).

8. Liver tumours

Anabolic steroids have been implicated as responsible for primary hepatocellular carcinomas and adenomas (Zimmerman, 1978). Similarly use of contraceptive steroids has been connected with liver tumours, particularly the oestrogenic components (Neuberger *et al.*, 1981a).

The mechanism(s) is unknown although interference with the metabolism of foreign compounds or bile salt derivatives so as to increase their tumourigenicity has been suggested (Goldfarb, 1976). 1976).

9. Vascular lesions

Anabolic steroids and contraceptive steroids may also be associated with rare vascular lesions such as peliosis hepatis (Zimmerman, 1978). The development of this hepatic lesion may be associated with the occurrence of adenomas. The mechanism is unknown, but may be related to damage to the membranes of the hepatocyte and sinusoid.

Conclusion

It is clear that drugs may cause a wide variety of hepatic lesions. In some cases these may be indistinguishable pathologically and biochemically from lesions due to other causes. Drug-induced hepatic damage ranges from the unpredictable and non-dose related to that occurring predictably after overdoses. Drug hepatotoxicity may involve metabolism to toxic, reactive intermediates and covalent interactions with cell constituents, interference with membrane transport or with cellular biochemistry such as protein synthesis, or immunological mechanisms. Furthermore several of these mechanisms may operate for one drug, giving rise to a variety of clinical and pathological features.

The occurrence of hepatic damage may be modified by differences in immune responsiveness and genetic, dietary and other factors.

The combination of such factors may give rise to incidences of hepatic toxicity varying from one in several thousands up to 10 or 20% of recipients.

As concluded by Williams *et al.* (1981) human susceptibility to drug-induced liver injury is clearly often multifactorial, and future approaches to the problem must take this into account.

References

ADLERCREUTZ, H. & TENHUNEN, R. (1970). Some aspects of the interaction between natural and synthetic

female sex hormones and the liver. *Am. J. Med.*, **49**, 630-648.

- ANGELI-GREAVES, M. & McLEAN, A.E.M. (1979). The effect of diet on the toxicity of drugs. In *Drug Toxicity*, ed. Gorrod, J.W., pp. 91-100. London: Taylor and Francis.
- BAHRI, A.K., CHIANG, C.S. & TIMBRELL, J.A. (1981). Acetylhydrazine hepatotoxicity. *Tox. appl. Pharmac.*, **60**, 561-569.
- BATCHELOR, J.R., WELSH, K.I., MANSILLA-TINOCO, R., DOLLERY, C.T., HUGHES, G.R.V., BERNSTEIN, R., RYAN, P., NAISH, P.F., ABER, G.M., BING, R.F. & RUSSELL, G.I. (1980). Hydralazine-induced systemic lupus erythematosus; influence of HLA-DR and sex on susceptibility. *Lancet*, **i**, 1107-1109.
- BLACK, M., MITCHELL, J.R., ZIMMERMAN, H.J., ISHAK, K.G. & EPLER, G.R. (1975). Isoniazid associated hepatitis in 114 patients. *Gastroenterology*, **69**, 289-302.
- BOYD, E.M. & BEREZKY, G.M. (1966). Liver necrosis from paracetamol. *Br. J. Pharmac.*, **26**, 606-614.
- BOYER, J.L. (1981). Membrane events in mechanisms of bile formation and drug-induced cholestasis. In *Drug Reactions and the Liver*, eds. Davis, M., Tredger, J.M. & Williams, R., pp. 64-71. London: Pitman Medical.
- BREEN, L., SCHENKER, S. & HEIMBERG, M. (1972). The effect of tetracycline on the hepatic secretion of triglyceride. *Biochem. Biophys. Acta*, **270**, 74-80.
- BREEN, K.J., SCHENKER, S. & HEIMBERG, M. (1975). Fatty liver induced by tetracycline in the rat; dose-response relationship and effect of sex. *Gastroenterology*, **69**, 714-723.
- CAPELLE, P., DHUMEAUX, D., MORA, M., FELDMAN, G. & BERTHELOT, P. (1972). Effect of rifampicin on liver function in man. *Gut*, **13**, 366-371.
- COX, R.P., FOLTZ, E.L., RAYMOND, S. & DREWYER, R. (1959). Novobiocin jaundice. *New Engl. J. Med.*, **261**, 139-141.
- DAHL, M.G.C., GREGORY, M.M. & SCHEUER, P.J. (1971). Liver damage due to methotrexate in patients with psoriasis. *Br. med. J.*, **1**, 625-630.
- DAVIS, J.S. (1968). Liver damage due to tetracycline and its relationship to pregnancy. In *Drug Induced Diseases*, eds. Meyler, L. & Peck, H.M., pp. 103-110. Amsterdam: Excerpta Medica Foundation.
- DAVIS, M. & WILLIAMS, R. (1977) Hepatic Disorders. In *Textbook of Adverse Drug Reactions*, ed. Davies, D.M., pp. 146-172. Oxford: Oxford University Press.
- DAVIS, M., VERGANI, D., MIELI-VERGANI, G., EDDLESTON, A.L.W.F., NEUBERGER, J. & WILLIAMS, R. (1981). Immunological studies on the pathogenesis of halothane associated hepatitis. In *Drug Reactions and the Liver*, eds. Davis, M., Tredger, J.M. & Williams, R., pp. 237-244. London: Pitman Medical.
- DeLORIMIER, A.A., GORDON, G.S., LOWE, R.C. & CARBONE, J.V. (1965). Methyltestosterone, related steroids and liver function. *Arch. intern. Med.*, **116**, 289-294.
- DOWLING, H.F. & LEPPER, M.H. (1964). Hepatic reactions to tetracycline. *J. Am. med. Ass.*, **188**, 307-309.
- EVANS, D.A.P. (1968). Genetic variation in the acetylation of isoniazid and other drugs. *Ann. N.Y. Acad. Sci.*, **151**, 723-733.
- GIRLING, D.J. (1978). The hepatic toxicity of anti-tuberculous regimens containing isoniazid, rifampicin and pyrazinamide. *Tubercle*, **59**, 13-32.
- GOLDFARB, S. (1976). Sex hormones and hepatic neoplasia. *Cancer Res.*, **36**, 2584-2588.
- GRONHAGEN-RISKA, C., HELLSTROM, P.E. & FROSETH, B. (1978). Predisposing factors in hepatitis induced by isoniazid-rifampin treatment of tuberculosis. *Am. Rev. resp. Dis.*, **118**, 461-466.
- HARGREAVES, T. & LATHE, G.H. (1963). Inhibitory aspects of bile secretion. *Nature*, **200**, 1172-76.
- HAVENS, W.P. (1962). Cholestatic jaundice in patients treated with erythromycin estolate. *J. Am. med. Ass.*, **180**, 30-32.
- HEMPEL, V., REMMER, H., MAY, R. & DIAZ DE TORANZO, E.G. (1981). Halothane associated hepatitis: Role of lipid peroxidation in man. In *Drug Reactions and the Liver*, eds. Davis, M., Tredger, J.M. & Williams, R., pp. 232-236. London: Pitman Medical.
- INMAN, W.H.W. & MUSHIN, W.M. (1974). Jaundice after repeated exposures to halothane: An analysis of reports to the Committee on Safety of Medicines. *Br. med. J.*, **1**, 5-10.
- ISHAK, K.G. & IREY, N.S. (1972). Hepatic injury associated with the phenothiazines: Clinico-pathologic and follow up study of 36 patients. *Arch. Path.*, **93**, 283-304.
- JAMES, O., ROBERTS, S.H. & DOUGLAS, A.P. (1975). Liver damage after paracetamol overdose. Comparison of liver function tests, fasting serum bile acids and liver histology. *Lancet*, **ii**, 579-581.
- KENRIGHT, S. & LEVI, A.J. (1974). Sites of competition in the selective uptake of rifamycin-SV, flavaspidic acid, bilirubin and bromosulphalein. *Gut*, **15**, 220-226.
- KLATSKIN, G. (1975). Toxic and drug-induced hepatitis. In *Diseases of the Liver*, 4th ed., ed. Schiff, L., pp. 604-710. Philadelphia: J.P. Lippincott.
- KLATSKIN, G. & SMITH, D.P. (1975). Halothane-induced hepatitis. In *Drugs and the Liver*, eds. Gerok, W. & Sickinger, K.F., pp. 289-296. Stuttgart: F.K. Schattauer Verlag.
- KOPANFF, D.E., SNIDER, D.E. & CARAS, G.J. (1978). Isoniazid-related hepatitis. *Am. Rev. resp. Dis.*, **117**, 991-1001.
- KUNELIS, C.T., PETERS, R.L. & EDMONDSON, H.A. (1965). Fatty liver of pregnancy and its relationship to tetracycline therapy. *Am. J. Med.*, **38**, 359-377.
- LENNON, D.H. (1966). Effects of various 17 α -alkyl substitutions and structural modifications of steroids on sulfobromophthalein (BSP) retention in rabbits. *Steroids*, **7**, 157-170.
- LENNON, D.H. (1966). Relative effects of 17 α -alkylated anabolic steroids on sulfobromophthalein (BSP) retention in rabbits. *J. Pharmac. exp. Ther.*, **151**, 143-150.
- LULLMAN, H., LULLMANN-RAUCH, R. & WASSERMANN, O. (1975). Drug induced phospholipidosis. *CRC Crit. Rev. Tox.*, **4**, 185-218.
- LULLMANN, H. & LULLMANN-RAUCH, R. (1977). Drug-induced lipidosis. In *Drug Design and Adverse Reactions*, eds. Bundgaard, H., Juul, P. & Kofod, H., pp. 29-34. Copenhagen: Munksgaard.
- LUNDE, P.K.M., FRISLID, K. & HANSTEEN, V. (1977). Disease and acetylation polymorphism. *Clin. Pharmacokinet.*, **2**, 182-197.
- MAIORINO, R.M., SIPES, I.G., GANDOLFI, A.J. & BROWN, B.R. (1979). Quantitative analysis of volatile halothane metabolites in biological tissues by gas chromatography. *J. Chromatog.*, **164**, 63-72.
- MITCHELL, J.R., LONG, M.W., THORGIERSSON, U.P. & JOLLO, D.J. (1975c). Acetylation rates and monthly liver function tests during one year of isoniazid prevent-

- ive therapy. *Chest*, **68**, 181–190.
- MITCHELL, J.R., POTTER, W.Z., HINSON, J.A., SNODGRASS, W.R., TIMBRELL, J.A. & GILLETTE, J.R. (1975b). Toxic drug reactions. In *Concepts of Biochemical Pharmacology* vol. XXVIII/3, eds. Gillette, J.R. & Mitchell, J.R., pp 383–419. New York: Springer-Verlag.
- MITCHELL, J.R., THORGIERSSEN, U.P., BLACK, M., TIMBRELL, J.A., SNODGRASS, W.R., POTTER, W.Z., JOLLOW, D.J. & KEISER, H. (1975a). Increased incidence of isoniazid hepatitis in rapid acetylators: Possible relation to hydrazine metabolites. *Clin. Pharmac. Ther.*, **18**, 70–79.
- MITCHELL, J.R., THORGIERSSEN, S.S., POTTER, W.R., JOLLOW, D.J. & KEISER, H. (1974). Acetaminophen induced hepatic injury: Protective role of glutathione in man and rationale for therapy. *Clin. Pharmac. Ther.*, **16**, 676–684.
- MITCHELL, J.R., ZIMMERMAN, H.J., ISHAK, K.G., THORGIERSSEN, U.P., TIMBRELL, J.A., SNODGRASS, W.R. & NELSON, S.D. (1976). Isoniazid liver injury: Clinical spectrum, pathology, and probably pathogenesis. *Ann. intern. Med.*, **84**, 181–192.
- MOULT, P. & SHERLOCK, S. (1975). Halothane-related hepatitis. *Quart. J. Med.*, **44**, 99–114.
- NEUBERGER, J.M., DAVIS, M. & WILLIAMS, R. (1981a). Clinical aspects of oral contraceptive associated liver tumours. In *Drug Reactions and the Liver*, eds. Davis, M., Tredger, J.M. & Williams, R., pp 271–283. London: Pitman Medical.
- NEUBERGER, J., TREDGER, J.M. & DAVIS, M. (1981b). Oxidative metabolism of halothane in the production of hepatocyte neoantigens. In *Drug Reactions and the Liver*, eds. Davis, M., Tredger, J.M. & Williams, R., pp 245–249. London: Pitman Medical.
- PETERS, J.H., MILLER, K.S. & BROWN, P. (1965). Studies on the metabolic basis for the genetically determined capacities for isoniazid inactivation in man. *J. Pharmac. exp. Ther.*, **150**, 298–304.
- PRESCOTT, L.F. (1981). Clinical use of protective agents after paracetamol overdose. In *Drug Reactions and the Liver*, eds. Davis, M., Tredger, J.M. & Williams, R., pp 157–163. London: Pitman Medical.
- PRESCOTT, L.F., WRIGHT, N., ROSCOE, P. & BROWN, S.S. (1971). Plasma paracetamol half-life and hepatic necrosis in patients with paracetamol overdose. *Lancet*, **i**, 519–522.
- ROS, E., SMALL, B.M. & CAREY, M.C. (1979). Effects of chlorpromazine hydrochloride on bile salt synthesis, bile formation and biliary lipid secretion in the rhesus monkey: a model for chlorpromazine induced cholestasis. *Eur. J. clin. Invest.*, **9**, 29–41.
- SCHULTZ, J.C., ADAMSON, J.S., WORKMAN, W.W. & NORMAN, T.D. (1963). Fatal liver disease after intravenous administration of tetracycline in high dosage. *New Engl. J. Med.*, **269**, 999–1004.
- SIPES, I.G., GANDOLFI, A.J. & BROWN, B.R. (1981). Halothane associated hepatitis: Evidence for direct toxicity. In *Drug Reactions and the Liver*, eds. Davis, M., Tredger, J.M. & Williams, R., pp 219–231. London: Pitman Medical.
- STRUNIN, L. (1977). *The liver and anaesthesia*. London: W.B. Saunders Co.
- TIMBRELL, J.A., WRIGHT, J.M. & BAILLIE, T.A. (1977). Monoacetylhydrazine as a metabolite of isoniazid in man. *Clin. Pharmac. Ther.*, **22**, 602–608.
- TIMBRELL, J.A., MITCHELL, J.R., SNODGRASS, W.R. & NELSON, S.D. (1980). Isoniazid hepatotoxicity: The relationship between covalent binding and metabolism *in vivo*. *J. Pharmac. exp. Ther.*, **213**, 364–369.
- TROWELL, J., PETO, R. & SMITH, A.C. (1975). Controlled trial of repeated halothane anaesthetics in patients with carcinoma of the uterine cervix treated with radium. *Lancet*, **ii**, 821–824.
- WHEELER, H.O. (1975). Secretion of bile. In *Diseases of the Liver*, 4th ed. ed. Schiff, L., pp 87–110. Philadelphia: J.P. Lipincott.
- WILLIAMS, R., TREDGER, J.M. & DAVIS, M. (1981). Introduction In *Drug Reactions and the Liver*, eds. Davis, M., Tredger, J.M. & Williams, R., pp 1–3. London: Pitman Medical.
- WOODMAN, D.D. (1981). Plasma enzymes in drug toxicity. In *Testing for Toxicity*, ed. Gorrod, J.W., pp 145–156. London: Taylor and Francis.
- WRIGHT, N. & PRESCOTT, L.F. (1973) Potentiation by previous drug therapy of hepatotoxicity following paracetamol overdosage. *Scot. med. J.*, **318**, 56–58.
- ZIMMERMAN, H. (1974). Hepatic injury caused by therapeutic agents. In *The Liver*, ed. Becker, F.F., pp 225–302. New York: Marcel Dekker.
- ZIMMERMAN, H.J. (1978). *Hepatotoxicity*, New York: Appleton-Century-Crofts.
- ZIMMERMAN, H.J. (1981). Drug Hepatotoxicity: Spectrum of clinical lesions. In *Drug Reactions and the Liver*, eds. Davis, M., Tredger, J.M. & Williams, R., pp 35–53. London: Pitman Medical.
- ZIMMERMAN, H.J. & SEEFF, L.B. (1970). Enzymes in hepatic disease. In *Diagnostic Enzymology*, ed. Coodley, E.L., pp 1–38. Philadelphia: Lea and Febiger.