BIOCHEMICAL PHARMACOLOGY OF PARADOXICAL SLEEP

J.-M. GAILLARD

Clinique psychiatrique de l'Université de Genève, Clinique de Bel-Air, 1225 Chene-Bourg, Switzerland

1 The role of noradrenergic cells in the regulation of paradoxical sleep is still controversial, and experimental data have given rise to contradictory interpretations.

2 Early investigations focused primarily on chemical neurotransmissions. However, the process of information transmission between cells involves many other factors, and the cell surface is an important site for transduction of messages into modifications of the activity of postsynaptic cells.

3 α -adrenoceptors are believed to play an important role in the control of wakefulness and paradoxical sleep. Experimental evidence suggests that physiological modulation of receptor sensitivity, possibly by specific neuro-modulators, may be a key mechanism in synaptic transmission.

4 In the investigation of the mechanisms involved in paradoxical sleep regulation, lesions of the locus coeruleus have given equivocal results. Collateral inhibition, probably mediated by α_2 -adrenoceptors, appears to be a powerful mechanism. The exact temporal relationship between noradrenergic cell activation and paradoxical sleep production is not established, but 5-HT appears to be involved. Differences between paradoxical sleep and waking may be related to a physiological modulation of α_2 -adrenoceptor sensitivity.

Keywords Paradoxical sleep α_2 -adrenoceptor noradrenaline 5-HT

Introduction

The application to the brain of the main concepts of neurotransmission developed for peripheral organs allowed major advances to be made during the decade 1960-70 in understanding sleep regulation. Two main factors were recognized during this period: first, the critical role of two groups of nuclei, the raphe nuclei and the locus coeruleus complex, in the brain stem, and, second, the possibility of modifying the activity of their transmitters, 5-hydroxytryptamine (5-HT) and noradrenaline (NA) respectively (Jouvet, 1972). Jouvet introduced the concept of wet neurophysiology, combining classical neurophysiological techniques with the biochemistry of neurotransmitters, and this emphasized the fact that the brain is not only a collection of interconnected wires, but also a chemical organ.

The monoamine hypothesis of sleep regulation inspired numerous experiments, but it was not long before problems emerged. Some results clearly supported the monoamine theory, whereas others equally clearly contradicted it. Thus, in some experiments 5-HT appeared to be a prerequisite for the maintenance of NREM-sleep, while in others it appeared irrelevant (Gillin *et al.*, 1978; Puizillout *et al.*, 1981). NA cells were seen by some as necessary for the occurrence of paradoxical sleep (PS), but other results indicated that their activation prevented

its appearance. In this confusing atmosphere, some authors even concluded that monoamines probably have, after all, nothing to do with sleep (Ramm, 1979).

Early enthusiasm declined and the investigation of sleep mechanisms has been less active during the last few years. However, when the early results are examined more closely it appears that the contradictions which occurred were caused by naive interpretations which failed to take into account the enormous complexity of the central nervous system.

Logicians know that when a theory leads to contradictions it is hopeless to try to solve them within the framework of the theory itself. On the contrary, it is necessary to build a wider theory which embraces the first and considers it as a particular case. In sleep research, this means that in practical terms it is useless to adopt a position either in favour of or against the involvement of 5-HT. It is more productive to consider new facts and observations collected outside the previous theory and to try to integrate them all from a new point of view. It is not yet possible to build a complete overall theory of sleep regulation because a number of results are still lacking. Nevertheless, interesting new avenues of research can be identified and experiments can be designed to collect the missing information.

Until now, the investigation of sleep regulation, as well as that of other functions of the central nervous system, has focused primarily on chemical neurotransmission. The amount of transmitter present in the synaptic cleft, as determined by the relative rate of synthesis, release and inactivation, is often cited as the primary factor governing the transmission of information from one cell to the next. However, it is becoming increasingly clear that much more complex mechanisms are involved and the concept of neuromodulation has recently emerged. A neuromodulator is a substance released by nerve terminals, unable by itself to trigger the postsynaptic cell but capable of modifying the effect of the main transmitter. The role of neurotransmitters and neuromodulators can be better understood by taking into account new data concerning the cellular biology of the neuronal systems under consideration. The cell surface is particularly important as a site at which messages from the outside are transduced into modifications of cell activity. Some data have been reviewed here, although several of the mechanisms have not yet been demonstrated in nerve cells. It is likely, however, that they are general properties shared by many different categories of cell, and taking the background into consideration may help to elaborate new hypotheses of sleep regulation.

It is evident also that the old principle of one neurone—one transmitter has to be reconsidered. The simultaneous presence of up to three transmitters within a single neuron has been demonstrated (Johansson *et al.*, 1981). It is therefore conceivable that one is the main transmitter while the others are released as neuromodulators.

Greater attention should also be paid to the modulation of receptor sensitivity. It is a well-established fact that deafferentation induces receptor hypersensitivity, but it is often believed that, under physiological conditions, receptors are stable units the activity of which depends mainly on the available transmitter. Recent developments in cellular biology indicate that the plasma membrane, as well as the receptors embedded in it, is subject to rapid changes over time. Not only are receptors constantly renewed, but mechanisms of rapid sensitivity modification, perhaps controlled by neuromodulators, can be envisaged. The study of receptor metabolism and sensitivity modulation opens up exciting new perspectives.

The present report is devoted primarily to identifying the usefulness of pharmacology as a tool in the investigation of sleep regulation. Great progress has been made in understanding the mode of action of pharmacological substances, although some controversies still persist. The main advantage of pharmacology is that it allows a molecular dissection of the processes under investigation, much finer than that achieved by other methods such as electrical stimulation or surgical lesion. In addition, it can be applied to

man and particularly to psychiatric conditions for which no consistent animal models have yet been found.

The cell surface

Several different models of membrane structure have been proposed in the past. Thermodynamic considerations indicated, however, that one of them, the fluid mosaic model described by Singer & Nicolson (1972), was more likely than others to be viable. Since membranes are composed of roughly half protein and half lipid, with a small proportion of carbohydrate, it was difficult at first to see which one of these components made up the matrix of the membrane. The fluid mosaic postulates that phospholipids form a two-dimensional fluid in which proteins are, in principle, free to move laterally. As we shall see later, this lateral mobility is restricted in a number of ways. Nevertheless, this model has gained wide acceptance and has been supported by a number of experimental data.

Membrane lipids

There is now considerable evidence that lipids in biological membranes are arranged in bilayer form, with the polar heads of the molecules facing outwards and the apolar tails kept away from water contact on the inside of the membrane (Gitler, 1976). It is important to realise that the viscosity of this fluid is relatively low, approximately comparable to that of edible oil, so that the protein molecules embedded in it are not restricted by viscosity from moving laterally. This holds true at temperatures above the lipid phase transition though at lower temperatures the proteins can be immobilized (Cherry, 1979). In this respect it may be relevant to recall that the phase transition of some biological lipids is close to 37°C. Several techniques are now available for measuring the lateral mobility of proteins; one of them, widely used, is that of fluorescence photobleaching recovery (Cherry, 1979).

The composition of lipids in the cell membrane differs from place to place and there is evidence for a predominance of polyunsaturated fatty acids of phospholipids in the inner leaflet (Nicolson, 1976). This asymmetry may conceivably result in differential fluidity of the two leaflets, with the inner more fluid than the outer. Among other functional consequences, this may facilitate ionic interactions, notably between Ca^{++} and proteins associated with the cytoplasmic side of the membrane. Calmodulin (CaM), a Ca^{++} -binding protein, could be involved in this mechanism, and seems in certain states to be associated with the cytoplasmic side of the membrane.

Some data also suggest that phosphatidylethanolamine is more abundant in the inner and phosphati-

dylcholine in the outer leaflet of the membrane (Hirata *et al.*, 1980). Two enzymes are responsible for the methylation of phosphatidylethanolamine, and during this process the phospholipids are transferred from the inner to the outer leaflet by a 'flip-flop' movement. The data above imply that the flip-flop rate of phospholipids is much higher than was previously believed (Hirata *et al.*, 1980). Methylation of phospholipids affects membrane fluidity and it is the first step in the methylation of phosphatidylethanolamine which enhances primarily the viscosity of membranes (Hirata *et al.*, 1980). However, it should be kept in mind that most of these results have been obtained in erythrocytes and some caution should be used in applying them to any other plasma membrane.

In addition to the asymmetry between the inner and outer leaflets, there are local heterogeneities in the phospholipid composition of the membranes. Several mechanisms by which microdomains of lipids differing in composition from their surroundings can be formed have been discussed by Nicolson (1976). The existence of these microdomains has been visualized by electron microscopy of freeze-fracture preparations using antibiotics which induce local membrane deformity as a consequence of their binding to lipids (Bearer & Friend, 1980). Even more important is the distinction between bulk lipids and boundary lipids. Boundary lipids are those which lie in close contact with proteins and are often extracted together with them (Gitler, 1976).

It is currently believed that apolar lipid-protein interactions are especially relevant, from a thermodynamic point of view, in anchoring the protein within the membrane. A considerable expenditure of energy would be necessary to move a protein along an axis normal to the plane of the membrane because the hydrophobic part of the protein would have to cross the polar region of the head groups of the phospholipids. Many enzymes require the presence of specific phospholipids for activity. It is likely that the polar interactions between the protein and specific head groups of phospholipids are relevant to this function (Singer & Nicolson, 1972; Gitler, 1976). Nevertheless, apolar interaction may also be involved since, for instance, the photochemical transformation of octopus rhodopsin is affected by modifications of the membrane lipid composition and particularly by the amount of unsaturated hydrocarbon chains present (Tsuda & Akino, 1981). This fact may be of significance also in the case of ligand-receptor interactions.

Membrane proteins

Membrane proteins have been classified into two broad categories, peripheral and integral proteins. Peripheral proteins are relatively weakly bound to the membrane from which they can be removed by mild treatments such as high ionic strength or metal ion

chelating agents. In fact, they are bound to integral proteins and appear to modulate or regulate some specific membrane functions. One example is spectrin, a peripheral protein of the erythrocyte, situated on the cytoplasmic side of the plasma membrane. Spectrin molecules are bound to integral glycoproteins and their function may be that of restricting their lateral mobility (Singer, 1974). They are also responsible for the biconcave shape as well as for the mechanical properties of these cells. Spectrin is a typical erythrocyte protein and it is unfortunate that this cell, the most readily available in large homogeneous quantities, is atypical and that its membrane cannot be considered as a model for other plasma membranes.

Peripheral proteins have the properties of soluble proteins. In the case of nerve cells, it is conceivable that some soluble proteins or peptides can bind in a similar way to the outer side of the cell membrane and possibly play a role in neuromodulation.

Integral proteins are much more tightly bound to membranes and can be removed by hydrophobic bond breaking agents, such as organic solvents or detergents. Sodium dodecyl sulphate (SDS) is an example, and polyacrylamide gel electrophoresis of SDS treated membranes is a useful tool in the analysis of integral proteins. These proteins have an amphipathic structure in intact membranes. Their hydrophobic part is not necessarily constituted of continuous stretches of apolar amino acids, but the folding of the molecule can bring together apolar amino acids separated by several polar residues along the chain. The inside of the molecule can thus be hydrophilic without violating the thermodynamic considerations mentioned above. This is probably the case for some ion channels. However, it is worth recalling that many integral proteins form complex aggregates in the membrane, e.g. the chloride channel-GABA receptor-benzodiazepine receptor. Some integral proteins span the entire membrane with a hydrophilic portion on either side of it, whereas others extend only partially within the thickness of the membrane.

Mobility of proteins and receptors within membranes

Three kinds of movement can be considered for integral proteins: rocking movements, floating movements along an axis normal to the plane of the membrane and lateral displacements. Rocking movements in which a binding site, for instance for a cation, might cross the entire thickness of the membrane in order to carry its ligand from the outside towards the inside of the cell have been postulated in the past. However, they are very unlikely to exist from a thermodynamic point of view and in some experimental models, such as bacteriorhodopsin, they have been ruled out (Cherry, 1979).

An X-ray diffraction study of the frog retinal

receptor disc membrane indicated that the rhodopsin molecule is only partially embedded in the lipid core; before bleaching, the molecule appeared to protrude into the aqueous layer, whereas bleaching caused it to sink into the lipid core by about one sixth of its diameter (Blasie, 1972). A reinvestigation of this model concluded, on the contrary, that the rhodopsin molecule probably spans the whole thickness of the membrane and that the bleaching-induced movements previously detected were artefacts (Chabre, 1975). Thus, floating movements, up and down with respect to the plane of the membrane, remain doubtful in the case of rhodopsin. However, Borochoy & Shinitzky (1976), using fluorescence labelling of erythrocyte membrane proteins have detected fluorescence changes following cholesterol depletion or enrichment, most likely caused by vertical displacements of protein. Shinitzky & Rivnay (1977) have shown that the degree of exposure of membrane proteins increases with the cholesterol/phospholipid ratio. When the microviscosity increases the proteins become more exposed to the aqueous medium, whereas fluidity enhancement causes them to sink into the membrane (Shinitzky & Henkart, 1979).

The fluid mosaic model predicts that lateral movements of protein molecules in the plane of the membrane are by far the most important (we do not consider rotational movements separately). A large body of experimental data supports this view. Redistribution of molecules within the membrane has been observed in several conditions. One of them, the patching and capping phenomenon, described first in the lymphocyte, has been observed subsequently in many other cells (but not in erythrocytes!). The addition of multivalent cross-linking agents such as immunoglobulins or lectins leads, within minutes, to aggregation of the binding sites into patches, followed by a gathering of these aggregates into a single cap (Taylor *et al.*, 1971; Lackie, 1980). The subsequent internalization of the cap by endocytosis involves, at least partly, the formation of coated pits and coated vesicles (Goldstein *et al.*, 1979; Salisbury *et al.*, 1980). These vesicles appear to be involved in more general processes of membrane recycling and may be an essential feature of internalization of ligand-receptor complexes (Anderson *et al.*, 1977). It has been shown recently that the formation of patches is accompanied by an accumulation of clathrin on the cytoplasmic side of the cell membrane opposite to the ligand-receptor clusters (Salisbury *et al.*, 1980). Clathrin is a large polypeptic (M.w. 180 000), the major component of coated vesicles (Pearse, 1976). In addition to clathrin accumulation, capping is also accompanied by an intracellular redistribution of CaM, which becomes concentrated in regions of the cytoplasm below the cap (Salisbury *et al.*, 1981). Trifluoperazine, a phenothiazine derivative used as neuroleptic drug, binds with high affinity to CaM in

the presence of Ca^{++} (Weiss & Levin, 1978). It inhibits CaM stimulation of phosphodiesterase and adenylate cyclase (Weiss & Levin, 1978). This agent does not block cap formation but inhibits clathrin recruitment to clustered receptor complexes. Under its effect, the capped region of the cell membrane remains uncoated and does not enter the pathway of endocytosis (Salisbury *et al.*, 1980). This suggests that clathrin recruitment may be CaM dependent (Salisbury *et al.*, 1981). The activity of CaM is regulated by Ca^{++} fluxes across the membrane and CaM controls the cytoplasmic movements of Ca^{++} (Cheung, 1980); therefore CaM may be a critical factor in the control of lateral protein motion in the plane of the membrane and particularly in receptor metabolism. Whether this mechanism can contribute to up-regulation of α -adrenoceptors under the influence of neuroleptic drugs remains to be seen.

It is, however, evident that in many cells of complex organisms integral proteins are not entirely free to move randomly within the fluid lipid matrix. It has been shown, for instance, that patching and capping is not independent of the cytoskeleton (see Nicolson, 1976). Capping is inhibited from a negligible to a moderate degree by cytochalasin B, while drugs disrupting microtubules have little effect. However, combined treatment of lymphocytes with cytochalasin B and colchicine or vinblastine markedly prevents capping (de Petris, 1974; Ryan *et al.*, 1974). Prior binding of concanavalin A inhibits Ig patching and capping, but this inhibition is suppressed by colchicine or vinblastine (Yahara & Edelman, 1973). Thus, in certain conditions, drugs which disrupt elements of the cytoskeleton affect drastically lateral movements of membrane proteins.

Frye & Edidin (1970) induced cultured cells of human and mouse origin to fuse by exposure to the Sendai virus. They used fluorescent dyes and appropriate antibodies to label respectively human and mouse antigenic components of the cell surface. They observed that, shortly after fusion, human and mouse antigenic components were largely segregated, whereas after 40 min at 37°C, the components were intermixed. Temperature lowering below 15°C markedly decreased this intermixing process. This experiment was considered an excellent demonstration of membrane fluidity. Later it was shown that the Sendai virus not only allows membranes to fuse and form heterokaryons, but also enhances the mobility of several surface components probably related to the cytoskeleton (Maeda *et al.*, 1979), by disrupting a restraint system.

The electrical charge of the cell membrane can affect the lateral mobility of components embedded in it. Charged molecules can be segregated if the potential drop across the length of the cell is of the order of a few millivolts. This is not uncommon in physiological conditions and may represent a surface

electrophoresis contributing to the distribution of components in cell membranes (Cherry, 1979).

In cultured embryonic muscle cells of *Xenopus laevis*, acetylcholine (ACh) receptors are distributed randomly over the cell surface. Application of an electrical field of 10 V/cm increases the sensitivity to microiontophoretically applied ACh on the cathode pole of the cells, indicating a redistribution of ACh receptors. Even more interesting is the fact that, under these conditions, ACh receptors form aggregates which are stable over time and not affected by colchicine or cytochalasin B (Poo, 1979).

Receptors of neurotransmitters are a class of (presumably) integral proteins of particular interest to us. The formation of contacts between nerve terminals and muscle cells during synaptogenesis induces the aggregation of ACh receptors in the muscle membrane and, conversely, denervation is followed by an increase of extrajunctional receptors (see Farmbrough, 1979). The effect of surgical denervation is mimicked by chronic blockade of receptors with α -bungarotoxin (Pestronk *et al.*, 1980). It has been shown recently that factors from the cell surroundings can affect the membrane distribution of receptors. When lyophilized material from neuroblastoma and glioma hybrid cell culture is added to rat primary myotubes, some of the mobile ACh receptors become immobilized, perhaps by anchorage to a cytoskeletal structure. Similar material from pure glioma cell culture is ineffective (Axelrod *et al.*, 1981). An effect in the reverse direction, i.e. an influence of postsynaptic structures over presynaptic endings also exists since extrajunctional ACh receptors play a role in the sprouting of ACh terminals (Pestronk & Drachman, 1978).

Some experimental data are available suggesting that membrane fluidity may affect either binding or function of receptors. Thus, fluidity of platelet membranes affects their sensitivity to the aggregating agent adrenaline (Insel *et al.*, 1978). Cholesterol depleted platelets have increased membrane fluidity and decreased sensitivity to adrenaline, whereas the reverse is seen in cholesterol enriched platelets. This effect is not due to a change in number or affinity of α -adrenoceptors for adrenaline and is unrelated to adenylate cyclase activity. The alteration of platelet response to adrenaline is distal to the ligand-receptor interaction and membrane fluidity appears to be critical in this effect. Fluidity of mouse brain membranes affects 5-HT binding by modifying $B_{\max}$ and K_D of both low and high affinity binding sites. Under conditions of high lipid fluidity, mainly high affinity 5-HT receptors are detected, whereas, in low fluidity, low affinity receptors predominate. There is no linear variation of binding as a function of microviscosity but a more complex relationship exists (Heron *et al.*, 1980). These results are interpreted in terms of vertical displacement of a membrane protein, with low

and high affinity sites located at different levels along the vertical axis of the receptor molecule. However, this is not the only possible interpretation; there may be more than two binding sites and these sites may be situated on different molecules. Moreover, the protein-lipid interactions are probably critical (Loh & Law, 1980). Nevertheless, the important point is the marked effect of membrane fluidity on 5-HT binding. Treatment of synaptosomal membranes with cholesterol, an agent which modulates membrane fluidity, results in a decrease of 5-HT uptake as well as in a considerable reduction of membrane associated adenylate cyclase (Paraphilis & Deliconstantinos, 1980).

Receptors within synapses

Although proteins embedded in the plasma membrane are free to move laterally in the lipid matrix, there are local arrangements of molecules in which mobility is restricted by bridges formed between them. A particular example is that of the synaptic junction. Smith & Loh (1978) have concluded from cross-linking experiments that proteins are free to move laterally in the synaptosomal plasma membrane, whereas they are ordered in arrays in the synaptic junction. Similar observations have been made with concanavalin A labelled receptors (Matus *et al.*, 1973). The use of cross-linking reagents such as bisimides, which have two reactive groups separated by various distances in the molecule, permitted gauging of the distance separating the cross-linked species. Reagents with reactive groups of 50 nm apart resulted in extensive aggregation of proteins in synaptosomal plasma membranes. The critical effect of this reagent cannot be accounted for by random collisions of the reactive species but indicates that they exist in ordered arrays in their native form. In these observations, there was also noted a minor participation of freely mobile proteins, probably located outside the synaptic junctions (Smith & Loh, 1979).

The ratio of carbohydrates and lipids to proteins in synaptosomal plasma membrane is higher than in other non-neuronal membranes, which probably reflects a high concentration of receptors and cell recognition sites. Most glycoproteins in the synaptosome are orientated externally (Smith & Loh, 1977) and an appreciable proportion of the proteins in this membrane can be isolated by alkaline extraction, a relatively mild treatment, suggesting that they are probably not very deeply embedded in it (Smith & Loh, 1977). Of the total protein molecules present in synaptosomes, roughly one third are external, one third are integral (probably spanning the whole thickness of the membrane), and one third are in an intermediate category, corresponding to integral proteins partially embedded in the membrane. Finally, it is worth emphasizing that a great similarity in composi-

tion exists between the membrane of nerve endings and of the remainder of the neuron. Also the smooth endoplasmic reticulum has SDS gel profiles very similar to those of synaptosome: and neurotransmitter receptors in particular are found in those intracellular membranes (see Smith & Loh, 1979).

The usual procedures for isolation of synaptosomes leave the post-synaptic membrane attached to the nerve ending, suggesting that relatively strong bonds hold the two together. The exact nature of these bridges is unknown, but they are sensitive to proteolytic digestion by trypsin or protease, while collagenase and neuraminidase are ineffective (Gordon Weeks *et al.*, 1981).

Insertion of proteins into membranes

If we consider the thermodynamic rules governing the relationships of proteins to the lipid matrix, it is apparent that proteins require special mechanisms for insertion into the membrane. It is believed that large portions of the cell membrane are constantly renewed (Holtzman & Mercurio, 1980, for review) and the synthesis of membranes seems to occur in the endoplasmic reticulum (Singer, 1974). Parts of it are released into the cytoplasm in the form of vesicles, which, after an intermediate step in the Golgi apparatus, become attached to and eventually fuse with the plasma membrane. Thus, the entire material of the vesicle is added to the membrane in such a way that the portions of protein molecules protruding into the inside of the vesicle come to be situated at the external surface of the cell. At the same time material contained by the vesicle (soluble proteins or smaller molecules) is released outside the cell. It is supposed that a similar mechanism exists at nerve endings (Smith & Loh, 1979).

One of the mechanisms for protein insertion into membranes has been elucidated by studies of the capsid protein of stomatitis virus and of Semliki forest virus (Rothman & Lodish, 1977; Rothman *et al.*, 1980). It is known as the signal hypothesis. The first 40 or so amino acids synthesized comprise a signal sequence which provokes the attachment to the rough endoplasmic reticulum of the full complex ribosome-messenger RNA and nascent polypeptide. The polypeptide is then pushed progressively through the membrane into the lumen of the endoplasmic reticulum as its synthesis proceeds. For glycoproteins, the carbohydrate residues are attached to the polypeptide chain as soon as it protrudes from the membrane. This process is consistent with the predominance of glycoproteins at the external surface of the cell. However, the existence of other mechanisms for protein insertion, and particularly post-translation insertion, is supported by experimental evidence (Wickner, 1980). Thus, proteins can penetrate the membrane by a helical hydrophobic hairpin struc-

ture which separates spontaneously from the aqueous cytoplasm into the hydrophobic bilayer (Engelman & Steitz, 1981). Such a mechanism does not require the presence of a special apparatus in the membrane and it might explain why some polypeptide chains cross the bilayer several times and/or have their COOH-terminus exposed on the external surface of the cell. The reverse would be expected according to the signal hypothesis, since the polypeptide synthesis begins at the NH₂-terminus. Of these possibilities, that which constitutes the exact method of receptor insertion into the membrane is not known at present.

As suggested by studies of virus synthesis, it is likely that, after their incorporation into the membrane, proteins can travel in the plane of the membrane randomly or as driven by elements of the cytoskeleton. They are not removed from the same site at which they are inserted (Lackie, 1980). It is possible, but has not been demonstrated, that neurotransmitter receptors are incorporated into the synaptic densities in this way. The synthesis of receptors probably follows the common pathway of protein synthesis outlined above, and thus their presence in membranes of the endoplasmic reticulum is to be expected.

There is good evidence that in nerve cells many membrane proteins, and particularly glycoproteins, are conveyed by fast axonal transport (Rambourg & Droz, 1980). There is a continuity between rough endoplasmic reticulum and axonal smooth endoplasmic reticulum. Proteins, and probably receptors in particular, are synthesized in the rough endoplasmic reticulum and transported down the axons embedded in the membrane of the smooth endoplasmic reticulum. The latter must be seen not as a system of pipes but 'as a dynamic system of membranes in permanent remodelling' (Rambourg & Droz, 1980). Contacts between smooth endoplasmic reticulum and axolemma can be observed under electron microscopy and allow the exchange of material between the two membranes. It is possible that axolemmal proteins are turned over in this way. Similarly, the smooth endoplasmic reticulum participates in the synaptic vesicle formation and in the renewal of membrane components in synaptic endings.

Neurotransmission and neuromodulation

The observation that electrical synapses exist in primitive animals, whereas more differentiated organisms rely exclusively on chemical synapses, may seem puzzling at first glance: direct electrical transmission from one cell to the next may appear more efficient than transmission by an intermediate and complicated chemical process. However, it is obvious that chemical synapses provide many more possibili-

ties for regulation. The postsynaptic cell is less strictly dependent on the activity of the presynaptic cell and can be modulated by many factors, including substances not released by the presynaptic ending but penetrating the synaptic cleft. The presynaptic terminal can release more than one transmitter with different effects on the electrical potential or the metabolism of the postsynaptic cell. This provides a means for highly differentiated neuromodulation.

In the classical scheme of neurotransmission, the transmitter is synthesized in the presynaptic cell. After release, it diffuses into the synaptic cleft, binds to receptors on the postsynaptic membrane and triggers a physiological effect. Finally it is inactivated, mainly by reuptake in the case of monoamines, but also by catabolism and by diffusion. Early studies of the influence of monoaminergic neurotransmission on sleep attempted to increase or decrease the amount of transmitter available. Thus *p*CPA-induced insomnia was discovered (Jouvet, 1972). However, unexpected results were sometimes obtained due to the lack of specificity of the transmitters and to secondary regulation of the nervous tissue. For instance, levodopa is decarboxylated not only in catecholaminergic but also in 5-hydroxytryptaminergic cells (Gaillard *et al.*, 1974); and the administration of α -methylparatyrosine to animals gave contradictory results which were not understood until recently (Kafi *et al.*, 1977, and see below).

α - and β -adrenoceptors

The classification of adrenoceptors into α and β types was made by Ahlquist (1948). Relatively little attention has been paid to the role of β -adrenoceptors in sleep regulation and drugs acting on them have, in general, limited effects (Dunleavy *et al.*, 1972; Hilakivi *et al.*, 1978). This impression of their relative lack of importance may prove false, but at present more is known about the involvement of α -adrenoceptors in this field.

An important advance has been achieved with the discovery of presynaptic receptors (Langer, 1977). They are powerful regulators of transmitter release: when stimulated by endogenous transmitter, e.g., noradrenaline (NA), they inhibit its further release. By contrast, when the transmitter is present in the synaptic cleft at very low levels, or when the receptors are blocked by an antagonist, transmitter release is enhanced. They in fact set the gain of the synapse by adjusting the per impulse release to the amount of transmitter already present outside the cell. The involvement of these receptors in sleep regulation has been shown in animals (Kafi & Gaillard, 1978) and in man (Gaillard & Kafi, 1979).

Postsynaptic and presynaptic receptors have been termed α_1 - and α_2 -adrenoceptors respectively (Langer, 1977). Recently however, the matter has

become more complicated and α_2 -adrenoceptors can no longer be considered synonymous with presynaptic receptors. Indeed, postsynaptic α_2 -adrenoceptors have been identified in several systems (Timmermans & Zwieten, 1981). Similar receptors have been detected at non-catecholaminergic terminals, including those in the cerebral cortex. In the latter structure they modulate the release of 5-HT; therefore they are probably situated on 5-HT terminals (Gothert *et al.*, 1981). Presynaptic α_1 -receptors have also been postulated (Kobinger & Pichler, 1980).

α_1 - and α_2 -adrenoceptors have different pharmacological properties. Antagonists with different selectivity for one or the other type are known, and some have mixed affinity for both types. In binding experiments, clonidine and WB 4101 are used as specific ligands for α_2 - and α_1 -adrenoceptors respectively (Taube *et al.*, 1977; Titeler & Seeman, 1980; Tanaka & Starke, 1980; Kobinger & Pichler, 1981). As emphasized by Ariens & Simonis (1976), there is no chemical relationship between α -adrenoceptor agonists and antagonists and they have very different profiles of selectivity. Most α -adrenoceptor antagonists interact with many different types of receptor: cholinergic, dopaminergic, α -adrenergic, histaminergic and 5-hydroxytryptaminergic. It can be assumed that agonists and antagonists do not bind to receptors in the same way. Agonists probably act by polar interaction, whereas antagonists bind to accessory sites adjacent to the receptor. It is likely that the lipid environment of the receptor plays a critical role in this interaction. The hydrophobic ring of many of these compounds may interact with the lipid annulus surrounding the receptor and thus interfere with its normal binding capacity for agonists. They may also interfere by different mechanisms with α -adrenoceptors, for instance, by altering their position within the membrane or their lateral mobility. In general, highly flexible molecules are not selective, probably because they can accommodate the different shape of various receptors. Conversely, rigid molecules are more specific (Ariens & Simonis, 1976).

The existence of presynaptic β -adrenoceptors has been postulated but not yet definitively established. Nevertheless, intriguing results point to an interaction between α_2 - and β -adrenoceptors, particularly at the prejunctional level. In rat atria incubated with [3 H]-NA, the β -adrenoceptor agonist isoprenaline enhanced the efflux of radioactivity in the presence of phentolamine, but decreased it in the presence of NA. The effect of β -adrenoceptor stimulation was thus inversely related to the degree of α -activation (Majewski & Rand, 1981). Other results indicate that in cortex slices the β -adrenoceptor agonist isoprenaline increases the number of α_2 -adrenoceptor binding sites while decreasing the number of β -binding sites (Maggi *et al.*, 1980a). Other types of interaction have been shown between α -adrenergic receptors and

muscarinic cholinergic receptors which seem to share the same Ca^{++} channels in rat parotid acinar cells (Putney, 1977; Davis *et al.*, 1980).

On the basis of their pharmacological properties, β -adrenoceptors have been subdivided into β_1 - and β_2 types. β_1 -receptors are innervated receptors, whereas β_2 -receptors mediate responses to circulating catecholamines (Ariens & Simonis, 1976; Bryan *et al.*, 1981). β -receptors activate the catalytic unit of adenylate cyclase (Minneman *et al.*, 1981). The enzyme is probably not permanently coupled to the receptor; both components float freely in the lipid sea and the activation of the enzyme takes place during the encounter with the receptor occupied by the ligand (Rimon *et al.*, 1978). GTP is involved in the coupling of β -receptors to the cyclase. The fluidity of the membrane can affect this coupling and also regulates the number of receptors accessible to binding. Methylation of the membrane phospholipids, and particularly the synthesis of phosphorylcholine, decreases membrane fluidity and enhances the number of binding sites for [^3H]-dihydroalprenolol without any change in affinity, thus uncovering hidden receptors (Strittmatter *et al.*, 1979).

The mechanisms involved in α -receptor activation have received less attention. Interaction of α_1 -adrenoceptors with an agonist causes mobilization of intracellular calcium, and, at least in some tissues, influx of extracellular Ca^{++} ions through specific channels (Exton, 1980). It is possible that phospholipid-inositol breakdown is the direct response to receptor occupation and triggers the intracellular modification of Ca^{++} distribution (Michell & Kirk, 1981), perhaps by synthesis of phosphatidic acid which may function as a Ca^{++} ionophore (Putney, 1981). This constitutes an amplification step, which is not, however, specific for α -adrenoceptors. Histamine (H_1)-, 5-HT $_1$ - and muscarinic-receptors share this mechanism. Activation of α_2 -adrenoceptors reduces NA liberation by decrease of Ca^{++} availability (Dismukes *et al.*, 1977). These receptors also inhibit, at least in some tissues, adenylate cyclase when it is activated by another transmitter.

Pre- and postsynaptic receptors also exist in 5-HT systems and function in a similar way (Haigler & Aghajanian, 1977). As for α_2 -adrenoceptors, pre-synaptic 5-HT receptors are situated not only in nerve terminals but also on the cell body. They mediate collateral inhibition in dorsal raphe nuclei (Wang & Aghajanian, 1978), a mechanism similar to that demonstrated in the locus coeruleus system (Aghajanian *et al.*, 1977; Cedarbaum & Aghajanian, 1978; Watabe & Satoh, 1980). LSD and other hallucinogenic substances are potent agonists of pre-synaptic 5-HT receptors, but no effective antagonist for postsynaptic 5-HT receptors active in all brain areas is known.

Modulation of receptor sensitivity

Modification of receptor sensitivity is a major mechanism regulating and adjusting synaptic transmission. Continuous stimulation by agonists, or excess of endogenous transmitter, decreases receptor sensitivity (down regulation). Conversely, the absence of endogenous transmitter, or continuous blockade with antagonists, leads to enhanced sensitivity (denervation hypersensitivity, up regulation (see Schwartz *et al.*, 1978)). More recently, modifications of sensitivity of β and α_2 -adrenoceptors have been demonstrated as a consequence of increased availability of the transmitter, itself a consequence of reuptake blockade by antidepressant drugs (Banerjee *et al.*, 1977; Crews & Smith, 1978; Spyraiki & Fibiger, 1980; Smith *et al.*, 1981). The α_2 -adrenoceptor agonist clonidine was also shown to down regulate these receptors (Engberg *et al.*, 1982). Receptor hypersensitivity induced by 6-hydroxydopamine (6-OHDA) is suppressed by continuous administration of NA (Biswas & Jonsson, 1981). On the other hand, numerous experiments have shown enhanced dopaminergic and nor-adrenergic sensitivity after chronic treatment with α -adrenoceptor antagonists.

The most frequently considered mechanism of sensitivity modification is alteration of the number of receptors. The state of hypersensitivity corresponds to more receptors present in the membrane whereas decrease in receptor numbers leads to hyposensitivity. However, more attention should be paid to the possible ways in which the receptor number is changed. As we have seen, receptors are synthesized using the protein synthesis machinery of the cell and inserted into the membrane during or after translation. They can move laterally in the plane of the membrane, with some restrictions, and are finally destroyed after internalization. Evidently this metabolism is controlled by a number of factors, some of them identified in particular models. However, caution should be exerted in generalizing from these results, as variations are possible from one system to the next. Thus, binding of the ligand to traditional hormone receptors is essential for the regulation of the number of receptors (Kaplan, 1981), whereas low-density lipoprotein (LDL) receptors are continuously recycled at a rapid pace, regardless of the presence or absence of the specific ligand (Goldstein *et al.*, 1979; Basu *et al.*, 1981). In class I receptors, according to Kaplan (1981), ligand-receptor interaction is followed by internalization and the presence of the ligand regulates the catabolism of receptors while their synthesis is unaffected. A similar mechanism could be responsible for up-regulation induced by α -adrenoceptor antagonists; by preventing the ligand-receptor interaction, the antagonist would interfere with receptor catabolism. If their synthesis

were unaltered, receptors would accumulate in the membrane. In addition, and as a consequence of their lipophilic properties, α -adrenoceptor antagonists could prevent receptor clustering, a necessary step before internalization for a number of proteins. Conversely, the presence of an agonist in excess would facilitate the catabolism of receptors not replaced rapidly enough by unmodified synthesis. This mechanism has not yet been established for α -adrenoceptors, but good evidence suggests that β -adrenoceptors may be regulated in this way. In frog erythrocytes, β -adrenoceptor agonists induce receptor clustering, probably mediated by transglutaminase and followed by internalization (Chuang *et al.*, 1980). However, this process does not involve the full receptor complex, but only the recognition site, since the catalytic units of adenylate cyclase were not found to be internalized in conditions of desensitization (Chuang *et al.*, 1980). Similar results have been obtained using cell fusion techniques: desensitization of erythrocytes with isoprenaline is followed by a decrease of recognition sites, whereas the catalytic unit of adenylate cyclase and the nucleotide regulatory protein remain unaltered (Pike & Lefkowitz, 1980).

In the case of α -receptors, the presence of agonist does not always, or not immediately, induce a desensitization. Preincubation of blood platelets with adrenaline leads, as is well known, to a decrease of receptor numbers as well as a decrease of adrenaline-induced aggregation. However, brief exposure of the platelets to the agonist results in an increase in receptor binding (Cooper *et al.*, 1978). The contractile response of rat vas deferens to adrenaline is enhanced by prior exposure to the same agent. In this particular experiment, an increase in the number of α -receptors as well as an enhancement of maximal response to adrenaline was observed (Hata *et al.*, 1980). The mechanism involved in these effects is not known.

Much experimental evidence supports the concept of spare receptors. For instance, the ED_{50} of positive inotropic effect induced by isoprenaline on cat papillary muscle is dose-dependently enhanced after prior exposure to the antagonist NHNB-NBE, whereas the maximal response remains the same regardless of the dose of the antagonist (Venter, 1979). It is admitted that, in physiological conditions, the number of receptors which must be occupied in order to elicit a maximal response is generally of the order of 10 to 30% of the total store of receptors present in the membrane. In this way the sensitivity of the cell may be regulated without modifications of the maximal response. Although verified in a number of systems, this concept may be an oversimplification since modifications of the maximal response are observed in some conditions, such as after reserpine administration (Schwartz *et al.*, 1978).

Experimental evidence suggests that the regulation

of cell sensitivity is more complex and not always related to a change in the number of receptors. Thus, chronic administration of the antidepressant drugs mianserin and zimelidine causes a decrease in the sensitivity of the cyclic AMP response to NA in the cortex of rats without any modification of the density of β -adrenergic receptors (Mishra *et al.*, 1980). Chlorimipramine and fluoxetine, potent inhibitors of 5-HT uptake, presumably enhance the extracellular concentration of 5-HT, but do not affect 5-HT binding in the brain (Wirz-Justice *et al.*, 1978; Maggi *et al.*, 1980b). These discrepancies suggest that changes in cell responsiveness are not related exclusively to variations in the absolute number of recognition sites. However, it is difficult to ascertain whether the number of binding sites accessible to the natural neurotransmitter *in vivo* is accurately reflected by *in vitro* experiments.

Several different mechanisms of receptor sensitivity modulation can coexist in some systems, as was shown for the rat parotid acinar cells in which two mechanisms with different time courses have been described (Davis *et al.*, 1980). The same experiments also indicated that cross-desensitization may exist between α -adrenergic and cholinergic receptors (Davis *et al.*, 1980).

The properties of cell membranes outlined above indicate that many different factors may participate in the regulation of receptor sensitivity. There are indications suggesting that clustering, microaggregation or cross-linking may be critical steps in receptor activation. Thus, incubation of neuroblastoma cells with rhodamine-enkephalin results in the slow appearance of patches which are not internalized. Prolonged contact of the culture with morphine does not result in down regulation and the patches remain stable for long periods of time. The receptors probably contain sulphhydryl or disulphide groups essential for the formation of patches, since fluorescence remains diffuse over the cell surface if the cells are preincubated with the sulphhydryl reagent dithiothreitol (Hazum *et al.*, 1981). We have seen above that ACh receptors presumably cluster under the effect of an electrical field. It is possible that receptor clustering is necessary for the efficient transmission of messages within the membrane and into the cell.

In pathological cases, cross-linking of receptors may lead to their accelerated degradation. In myasthenia gravis, the number of ACh receptors is decreased. This decrease is secondary to interactions with antireceptor antibodies, which results in blockade, cross-linking and accelerated degradation of receptors (Lindstrom *et al.*, 1976; Drachman *et al.*, 1977, 1978; Appel *et al.*, 1977; Pollet & Levey, 1980).

Experimental evidence is not yet abundant, but suggests clearly that modulation of receptor sensitivity is a key mechanism in synaptic transmission. It is

even perhaps more important than the simple availability of the transmitter. This modulation can involve the number of receptors, their disposition (both in the plane and normal to the plane of the cell membrane), their cross-linking, their coupling to the intracellular chain of events they normally trigger and, finally, their interaction with membrane lipids. Several factors can affect one or another of these mechanisms: the transmitter itself (agonist), a co-transmitter (a substance released by the nerve terminal together with the main transmitter) or a substance coming from the extracellular space and released in a more remote location. Such neuromodulators may induce optimal disposition of recognition sites in the membrane (clustering?) or may be necessary to maintain together the parts of the receptor complex required for signal transduction. Based on our present knowledge of the biology of membranes, other mechanisms are probably conceivable and amenable to experimental investigation.

The problem of paradoxical sleep

The early experiments of Jouvet pointed to a positive involvement of brain NA neurons, especially those of the locus coeruleus (LC), in the maintenance of paradoxical sleep (PS) (see Jouvet, 1972). Some results also indicated that 5-hydroxytryptaminergic neurons may play a role in the priming of this stage. The existence of a cholinergic link in this complex chain of events was suggested by the observation of a PS-suppressing effect of parenteral atropine both in normal and pontine cats (Jouvet, 1952). It has been shown also that cholinergic stimulation of the brain stem reticular formation induces PS (George *et al.*, 1964; Mitler & Dement, 1974; Amatruda *et al.*, 1975; Silberman *et al.*, 1980). Some modifications of PS observed in human pathology have been ascribed to changes in the sensitivity of muscarinic receptors (Sitaram *et al.*, 1978; Gillin *et al.*, 1979).

While a cholinergic participation in the control of PS appears very likely, the exact role of dopaminergic (DA) and NA neurons is still a highly disputed issue. A large number of contradictory data have appeared in the literature, giving the impression that whatever the manipulation aimed at catecholaminergic (CA) neurons, the results obtained are random and unpredictable. The confusing picture which emerged from these reports led to the conclusion that brain CA systems are not involved in any critical aspect of the regulation of PS, but may reflect at most a more general neuromodulating function of LC neurons (Ramm, 1979).

Although it had been considered for some time that NA promotes sleep (Mandell & Spooner, 1968), it is now generally accepted that this transmitter is involved in sustaining wakefulness (Torda, 1968;

Gaillard *et al.*, 1969; Petitjean & Jouvet, 1970; Jones, 1972; Panksepp *et al.*, 1973; Lidbrink, 1974). NA seems to be more specifically responsible for the EEG activating aspects of wakefulness, whereas DA regulates waking behaviour (Jones *et al.*, 1969). Subtotal depletion of brain CA completely suppresses waking (Keane *et al.*, 1976). PS is also a state of intense activation of the central nervous system, and, despite obvious differences, has several aspects in common with wakefulness. Therefore it seemed reasonable to suppose that catecholamines are positively involved in at least some aspects of the mechanisms controlling PS.

Lesions of the locus coeruleus

Coagulation of the ascending dorsal NA bundle in the region of the ponto-mesencephalic junction (the isthmus) and lesions of the anterior third of the LC induce a decrease of wakefulness, perhaps by disinhibition of 5-HT neurons (Petitjean & Jouvet, 1970). Coagulation of the caudal two thirds of the LC decreases by 30 to 40% the NA content of the telencephalon and suppresses the occurrence of PS, without preventing completely the appearance of PGO waves. Total lesions of the LC and subcoeruleus nuclei cause a more significant reduction of NA in the forebrain and lead to decrease of waking as well as to a permanent suppression of PS and PGO waves (Buguet, 1969). These observations and their interpretation have been challenged by others who have shown that, in the cat, extensive destruction of the LC (69%), giving 85% decrement of NA content in the cortex, did not prevent PS reappearing within 48 h and recovering rapidly to normal levels. It was concluded that LC was not essential for the occurrence of PS, although it may play a role in the control of its phasic components, the PGO waves (Jones *et al.*, 1977). It has been claimed that in the cat lesions destroying the pontine gigantocellular tegmental field but sparing the LC resulted in a complete suppression of PS and particularly of the muscular atonia normally present in this stage (Jones, 1979). FTG cells were believed to discharge selectively during PS and therefore to represent the best candidate for PS generator neurons (Hobson *et al.*, 1974; Steriade & Hobson, 1976). More recently, Sastre *et al.* (1981) have demonstrated that total destruction of FTG cells with kainic acid was not followed by any significant alteration of PS, thus ruling out the critical role of these cells in the generation of this stage.

Obviously, the results of electrolytic or surgical lesions are difficult to interpret, especially in the cat, an animal in which the neurones of the coerulean complex are not tightly packed. Among the numerous reasons for discrepancies are uncertainties about the maximum size of the lesions the LC could tolerate without marked disruption of its functional capa-

bility, the difficulty of precise limitation of the lesion to coerulean neurones without severing neighbouring structures, the fact that the LC contains two types of cell (Groves & Wilson, 1980a) (one of them being possibly cholinergic and equally destroyed by coagulation) and the possible interruption of passing fibres connecting other structures. Another reason for the difficulty in drawing conclusions from this type of experiment resides in the anatomical disposition of NA cells in the brain stem. These cells may be subdivided into three major groups: an LC group, including the A4 and A6 group of Dahlstrom & Fuxe (1964), a dorsal medullary group (A2) and a lateral tegmental group (A1, A3, A5, A7) (Grzanna & Molliver, 1980). Some of the NA cells, at least in the rat, are densely packed in the A6 nucleus, and constitute the main part of the LC complex. The remaining cells, however, are much more loosely distributed in large areas of the brain stem, dorsally, centrally, caudally and rostrally with respect to the LC. A lesion extensive enough to destroy all these cells would certainly be incompatible with the survival of the animal.

Unit activity of the locus coeruleus

The second technique used in the investigation of the relationships between LC neurones and the sleep-waking cycle is the study of their spontaneous discharge by unicellular recordings. Earlier data were obtained only in the cat (Chu & Bloom, 1974; Hobson *et al.*, 1975; Sakai, 1980), in which species NA cells are intermingled with non-NA cells. In this animal, several units in the LC region tended to discharge faster during PS than during slow wave sleep. These units were identified by histofluorescence as NA cells (Chu & Bloom, 1974). Further investigations suggested that most, if not all, NA cells in this region were PS-off, that is they stopped firing in PS (Hobson *et al.*, 1975; McCarley & Hobson, 1975; Sakai, 1980). A more recent report, based on unicellular recordings in the rat, indicates a high rate of discharges in active wakefulness, a lower rate in quiet sleep and almost total silence in PS (Aston-Jones & Bloom, 1981). It can be argued that only nine cells were recorded in PS, representing a rather small sample of the roughly 1,600 NA cells present in the rat LC, and that they were not identified by antidromic stimulation in order to calculate their conduction velocity characteristics. Nevertheless, we believe that this investigation reflects accurately the true situation. It is worthwhile stressing that these recordings were obtained from the most dense part of the LC complex, where the NA nature of the cells could be established without ambiguity. Moreover, these authors made the interesting observation that cells situated at the edge of the nucleus tended to discharge faster during all three states of alertness, although the difference was significant only for wakefulness (Aston-Jones & Bloom, 1981).

6-Hydroxydopamine and α -methyl-p-tyrosine

Until now, neuropharmacological investigations of PS regulation have been carried out mainly using neurotoxic compounds such as 6-OHDA, α -methyl-p-tyrosine (AMPT) and various pharmacological agents acting on α -adrenoceptors. Reserpine and other compounds have been used as well, but will not be discussed here because of their lack of specificity.

Intraventricular injection of 6-OHDA in the cat results in a dose-dependent decrease of PS, even if 5-hydroxytryptaminergic neurones are protected by concomitant treatment with chlorimipramine (Laguzzi *et al.*, 1972). A similar decrease is seen in the rat, followed by a gradual recovery to nearly normal levels after about 8 days (Matsuyama *et al.*, 1973). However, the nonspecific effect of 6-OHDA caused objections to be raised and different results were obtained by other investigators (Hartmann *et al.*, 1971). The results of intraventricular injections can be difficult to interpret, because the pattern of penetration of the drug into the brain is generally unknown. If some structures are preferentially affected, a modification of equilibrium can occur in an unexpected direction. Intracerebral injection of 6-OHDA in the region of the ventral NA pathway resulted in an increase in PS, but this is difficult to interpret for the reasons mentioned above (Panksepp *et al.*, 1973).

AMPT, an inhibitor of tyrosine hydroxylase, has generally been shown to enhance PS in the rat (Hartmann *et al.*, 1971), in the cat (Iskander & Kaebbling, 1970; King & Jewett, 1971; Henriksen & Dement, 1972; Stern & Morgane, 1973; Stein *et al.*, 1974) and in man (Wyatt *et al.*, 1971, 1972). These data were regarded as supporting the idea of a negative relationship between PS and brain CA activity. In some studies, however, AMPT was without effect (Marantz *et al.*, 1968; Branchey & Kissin, 1970) whereas a few reports described a decrease of PS in the rat (Torda, 1968) and in the monkey (Weitzman *et al.*, 1969). A difficulty posed by AMPT is its toxicity, which prevents its being given as a single injection in sufficiently high dosage. In almost all the experiments quoted above, the dosage used, probably for this reason, was between 50 and 100 mg/day. Histofluorescence studies revealed that 50 mg/kg in the cat induced only a minimal decrease of endogenous CA; after 100 mg/kg, the decrease was obvious in cell bodies, but still limited in terminals (Kafi *et al.*, 1977). A method of treatment consisting of several injections of 75 mg/kg every 4 h allowed the avoidance of the toxic effects of AMPT and resulted in complete disappearance of endogenous CA fluorescence after a total dose of 525 mg/kg. The animals appeared sedated but recovered well. Sleep recordings revealed that after total doses of 75 or 150 mg/kg, the production of PS was enhanced, whereas it was decreased after higher doses administered in this way

(Kafi *et al.*, 1977). This result showed that both increase and decrease of PS could be obtained in the same animal depending on the dose of AMPT. It also indicated that when the depletion of endogenous CA was sufficient, a depression of PS was observed, whereas a minimal decrease of CA may remove some inhibitory influence while leaving enough transmitter to ensure normal NA synaptic activity. The exact mechanism of this effect was not understood at the time.

Agonists and antagonists of α -adrenoceptors

Other pharmacological tools are available for the study of α -adrenoceptors in PS regulation. Clonidine is a potent and specific agonist at α_2 -adrenoceptors (Starke & Altmann, 1973; Starke & Montel, 1973; Starke *et al.*, 1974). It binds specifically to α_2 -adrenoceptors in the cortex (Titeler & Seeman, 1980), and as a consequence of α_2 -activation it reduces turnover of NA (Anden *et al.*, 1970; Braestrup, 1974). Given systemically or by microiontophoresis, clonidine inhibits the firing of NA cells in the LC, whereas DA cells are not influenced (Svensson *et al.*, 1975).

Clonidine reduces PS in the rat and in the cat (Kleinlogel *et al.*, 1975; Leppavuori & Putkonen, 1980). A similar decrease has been observed in man with a dose roughly five times smaller than that used in the rat (Gaillard & Kafi, 1979). We have recently examined the dose-effect relationships of PS inhibition in the rat (Gaillard *et al.*, 1982). With increasing doses, the latency of PS was enhanced and total PS was decreased, mainly by shortening of the PS episodes. The calculation of the general trend, a function describing the evolution of the stage with respect to sleep time (Gaillard, 1979), revealed that after the initial period of complete inhibition, PS was subsequently produced at a higher rate than in control conditions, suggesting a rebound phenomenon. This rebound did not appear if the animal had been treated with AMPT 24 h earlier, a pretreatment presumably enhancing the sensitivity of α -adrenoceptors (Kafi & Gaillard, 1981).

The relationships between the latency of PS and the log-dose of clonidine was not linear but parabolic, suggesting that clonidine acted at more than one site (Gaillard *et al.*, 1982). One possibility is that at low dose, clonidine may act predominantly in the locus coeruleus, where α_2 -adrenoceptors are more densely packed, a factor likely to facilitate drug-receptor interaction. At higher doses an effect of clonidine on other receptors, probably situated on more remote NA terminals, is added progressively.

The possibility of PS rebound after clonidine was also investigated in man (Gaillard *et al.*, 1983). Sixteen normal young adults were recorded in two laboratory sessions, using standard sleep monitoring techniques. In one session, they underwent one

habituation night followed by two baseline nights, and, in the other, the habituation night was followed by one night under the influence of clonidine (0.4 μ g/kg) and one placebo night. The order of the sessions was balanced so that the sequence clonidine-placebo was exactly comparable to the sequence of the two baseline nights, in order to control for a possible effect of the sequential order of the nights spent in the laboratory. Under these conditions, clonidine did not induce any significant modification of PS, but on the next night a highly significant increase was manifest. Examination of the individual results indicated that four subjects of the sample had a decrease of PS under clonidine, whereas the remaining 12 had no modification. Interestingly, the four affected subjects had no PS increase on the following night as compared to the baseline, whereas the 12 subjects showing no direct effect of clonidine experienced a marked subsequent increase (Table 1). This is the first experimental model in which an increase of PS was observed after a manipulation that by itself does not induce any change. We consider that the mechanism of this increase is identical to that responsible for PS rebound in other experimental situations, such as that following instrumental deprivation or withdrawal of antidepressant drugs; therefore, it represents a rebound without previous debt. The mechanism of PS enhancement on the placebo night is probably related to clonidine-induced desensitization of α_2 -adrenoceptors.

Another interesting observation was made during this experiment. The comparison of baselines revealed that the group of 12 subjects with PS rebound had, on average, about twice as much stage 4 sleep as the group without rebound. These subjects also had shorter latencies of stages 3 and 4. This suggests that the mechanisms underlying slow wave sleep production were weaker in the subjects who had no PS rebound after clonidine. This observation points to a possible link between the structures responsible for slow wave sleep generation and PS regulation.

In another series of experiments, we investigated the effects of three antagonists of α -adrenoceptors: chlorpromazine (CPZ), yohimbine (YOH) and piperoxane (PIP) (Kafi & Gaillard, 1978; Gaillard & Kafi, 1979; Kafi & Gaillard, 1980; Gaillard *et al.*, 1983). CPZ is a potent antagonist of α_1 -adrenoceptors, but also has a significant effect on α_2 -adrenoceptors. The reverse is true for YOH, which has a higher affinity for α_2 - than for α_1 -adrenoceptors. PIP has an even higher affinity for α_2 -adrenoceptors, while α_1 -antagonism only appears at very high doses.

In the rat, doses of CPZ below 1 mg/kg facilitated PS production, whereas doses higher than this threshold induced a dose-related decrease. If a low dose of CPZ was preceded by pretreatment with AMPT (150 mg/kg), not only did the PS facilitation no longer appear but it was replaced by a marked

Table 1 Parameters of paradoxical sleep (PS) in 16 normal young adults under and after CLN.

Group I: Twelve subjects without PS decrease under clonidine, with significant increase in the following placebo night.

Group II: Four subjects with a decrease of PS under clonidine, without significant increase over baseline in the following placebo night

	Baseline	Clonidine (0.4 μ g/kg)	Placebo
Group I (<i>n</i> = 12)			
PS (min)	101 $\pm$ 21	105 $\pm$ 22	116 $\pm$ 21*
PS (%)	21.3 $\pm$ 3.2	21.9 $\pm$ 4.2	24.6 $\pm$ 4.0***
PS latency (min)	101 $\pm$ 31	101 $\pm$ 43	85 $\pm$ 24
Group II (<i>n</i> = 4)			
PS (min)	112 $\pm$ 11	74 $\pm$ 22**	121 $\pm$ 17
PS (%)	23.9 $\pm$ 2.2	15.1 $\pm$ 2.6***	25.2 $\pm$ 2.8
PS latency (min)	88 $\pm$ 23	139 $\pm$ 56	90 $\pm$ 41

P* < 0.05; *P* < 0.01; ****P* < 0.005 (two-tailed Student's *t*-test).

decrease. This dose of AMPT given alone and in a single injection, in contrast to the effect of two injections of 75 mg/kg described above, induced a small and non-significant decrease of PS production. The same pretreatment before a dose of CPZ, which by itself diminished PS, induced an even more important decrease (Gaillard *et al.*, 1982). It was also shown that a very small dose of clonidine, ineffective by itself, could suppress the PS facilitation following a small dose of CPZ (Gaillard & Kafi, 1979). These data support the view of a preponderant effect of CPZ at low doses on presynaptic α_2 -adrenoceptors, facilitating the release of NA and compensating for the blockade of α_1 -adrenoceptors. This blockade is revealed only when AMPT prevents the increase of NA turnover, which is known to occur as a consequence of α -adrenoceptor antagonism (Bartholini *et al.*, 1973). With increasing doses of CPZ, the α_2 effect cannot increase to the same extent as the α_1 effect, and the latter becomes predominant. It has therefore to be assumed that both effects exist whatever the dose, but that the result on PS which we see depends on the algebraic sum of these prejunctional and postjunctional influences. It is also possible that CPZ acts proportionally more on postsynaptic than on presynaptic α_2 -adrenoceptors at the site of collateral inhibition, a condition necessary for PS facilitation to occur (see below).

The results of YOH supported the same interpretation (Kafi & Gaillard, 1980). A small dose of this compound (0.03 mg/kg) induced a facilitation of PS production, whereas a dose 100 times larger tended to decrease PS. Thus, the effect occurred in the expected direction, but was smaller than anticipated. Indeed, the better specificity of YOH for α_2 -adrenoceptors led us to expect a greater PS facilitation than was actually observed. We began to suspect that

another mechanism was acting as a brake and maintaining PS modifications within relatively narrow limits.

Combination of YOH with AMPT indicated that, as expected, α_1 -antagonism occurred only at high doses of the compound. Pretreatment with AMPT before a low dose of YOH suppressed the facilitation seen with YOH alone and gave rise to a PS production only slightly below control levels; in contrast, a high dose of YOH after AMPT showed a CPZ-type effect, that is a marked decrease of PS compared to YOH alone (Gaillard *et al.*, 1982).

Because of the high selectivity of PIP for α_2 -adrenoceptors (Anden *et al.*, 1976; Franklin & Herbert, 1977; Brown *et al.*, 1980), we expected in our first experiments with this compound a PS facilitation much larger than with CPZ for comparable doses. As it happened, the results showed a dose-response relationship exactly opposite to our predictions. The dose of 1 mg/kg caused a significant decrease of PS (Table 2). After doses between 1 and 20 mg/kg a progressive facilitation of PS was observed, reaching a maximum after 20 mg/kg and decreasing slightly for higher doses. After 20 mg/kg, the percentage of PS was not significantly different from control values, but the calculation of the general trend yielded significant parameters (level: *P* < 0.05; slope: *P* < 0.01; *n* = 3) to indicate that the PS facilitation at this dose was statistically significant. For doses below 1 mg/kg, no facilitation of PS was seen.

The combination of AMPT with 1 mg/kg of PIP did not modify the effect obtained after PIP alone, suggesting that the decrease of PS was not related to blockade of α_1 -adrenoceptors. However, AMPT given 30 min before 20 mg/kg of PIP suppressed the PS facilitation elicited by the same dose of PIP alone.

It is thus evident that the effects of PIP bear some

Table 2 Effect of various doses of piperoxane on the main parameters of wakefulness and paradoxical sleep in the rat. The animals received the drug i.p. at 08.00 h and were recorded for the next 11 h. Waking was calculated as a percentage of total sleep recording time; PS was calculated as a percentage of total sleep.

Sleep latency is the interval in min between the injection and sleep onset.

PS latency is the interval between sleep onset and the first onset of paradoxical sleep.

	n	Waking (%)	PS (%)	Sleep latency	PS latency
Piperoxane 0 (mg/kg)	16	31 ± 6	16 ± 3	10 ± 12	13 ± 6
0.2	4	32 ± 2	15 ± 3	16 ± 5	12 ± 6
0.46	5	34 ± 7	16 ± 2	21 ± 8	15 ± 7
1	4	51 ± 9***	12 ± 1*	35 ± 9***	13 ± 8
2	4	46 ± 8***	14 ± 2	48 ± 40***	20 ± 6
4.6	3	35 ± 1	15 ± 2	35 ± 3***	30 ± 9***
10	3	38 ± 1*	17 ± 4	24 ± 11	21 ± 5
20	3	36 ± 4	18 ± 1	31 ± 20**	14 ± 11
46	4	41 ± 5**	17 ± 3	29 ± 7**	17 ± 6

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.005$ (two-tailed Student's *t*-test).

resemblance to those of YOH, particularly when the combination with AMPT is considered, and are opposite to those of CPZ. At low doses, CPZ facilitates, whereas PIP inhibits, PS; if the animals have been pretreated with AMPT, CPZ is followed by a marked decrease whereas the effect of PIP is not modified. After high doses, CPZ decreases whereas PIP facilitates PS. Pretreatment with AMPT enhances markedly the CPZ-induced inhibition of PS and only suppresses the PIP-induced facilitation.

The role of collateral inhibition

The effects of CPZ and YOH can be accounted for by a balance between presynaptic and postsynaptic α -antagonism. The effect of PIP is more difficult to understand. Before attempting an interpretation, it is necessary to report other experiments in which we investigated the possible role of collateral inhibition in the decrease of PS induced by antidepressant drugs. In the LC, the presence of CA nerve terminals (Hökfelt, 1967; Groves & Wilson, 1980a, b), as well as dendro-dendritic connections (Sladek & Parnavelas, 1975), has been established by histochemical and electron microscopic studies. The axons of the LC neurons possess collateral branching visible in Golgi preparations (Shimizu & Imamoto, 1970; Swanson, 1976). Results of single unit recordings support collateral inhibition in the LC, allowing NA neurons to exert homo- and contralateral inhibitory control on each other (Aghajanian *et al.*, 1977; Cedarbaum & Aghajanian, 1978; Watabe & Satoh, 1980). Thus anatomical and electrophysiological data are consistent with the view that LC neurons send collaterals possibly to themselves, certainly to neighbouring cells and to cells in the contralateral nucleus, inhibiting their discharge rate. Microiontophoretic studies

suggest that this inhibition is mediated by α_2 -adrenoceptors (Aghajanian *et al.*, 1977; Cedarbaum & Aghajanian, 1977). α_1 -adrenoceptors have not been detected in the LC, and it is reasonable to suppose that α_2 -adrenoceptors have both a presynaptic and a postsynaptic location in this structure.

Desipramine (DMI), a classical antidepressant with specific inhibition of NA reuptake and very slight anticholinergic activity, induced a dose-dependent decrease of PS production in the rat (Justafre & Gaillard, 1981) and in man (Justafre & Gaillard, unpublished results). Rats treated with 150 mg/kg of AMPT and 30 min later with 4 mg/kg of DMI showed significantly less decrease of PS than rats receiving the same dose of DMI alone (Justafre & Gaillard, 1981). This observation suggests that DMI potentiates collateral inhibition, probably by enhancing the extracellular concentration of NA and thus stimulating post-synaptic α_2 -adrenoceptors situated on the cell bodies of NA nuclei of the brain stem. The pretreatment with AMPT decreases slightly the endogenous supply of NA, thereby decreasing the PS depressing effect of DMI.

The facilitation of PS induced by a small dose of AMPT reported above can be accounted for in the same way, but we have to take into consideration an additional factor. Indeed if a small decrease of endogenous NA reduces collateral inhibition slightly, it is at first glance difficult to understand why it should not reduce simultaneously the NA synaptic transmission in the effector region, and notably in the cortex, thus re-establishing the balance between collateral inhibition acting as a brake and remote NA terminals facilitating PS. It has been shown recently that extrasynaptic diffusion is an important factor in inactivation of the extracellular transmitter (Baumann & Koella, 1980). In regions where activated noradren-

ergic synapses are densely spaced, extracellular concentration of NA may rise to levels interfering with diffusion of the transmitter away from neighbouring synapses. It is also possible that NA diffuses from one synapse to the next, producing a phenomenon of cross-talk between them (Baumann & Koella, 1980). This is less likely in regions in which synapses are less densely spaced. It has been estimated that the incidence of NA synapses in the cortex is less than 1 per 1,000 total synapses (Lapierre *et al.*, 1973), whereas as many as 4 out of 100 synapses in the LC may be noradrenergic (Groves & Wilson, 1980b). If these estimates are correct, they mean that NA synapses are considerably more densely spaced in the LC than in the cortex, with a ratio of 40 to 1. It follows that we can expect more cross-talk between synapses in the LC. Therefore, a slight decrease of the endogenous transmitter is likely to have a more critical effect in the LC than in the cortex or other remote regions of NA appositions, thereby releasing partially the collateral inhibition of NA cells. This imbalance explains the PS facilitation observed with low doses of AMPT, whereas after higher doses NA transmission is impaired in the effector region as well, leading to PS decrease. For the same reason, the effect of DMI is more important on collateral inhibition than in the effector region, leading to PS decrease.

With these data at hand it is now possible to propose an interpretation for the effect of PIP. This drug has a high affinity for α_2 -adrenoceptors and enhances the release of NA from presynaptic terminals, at the site of collateral inhibition within the NA nuclei of the brain stem and in the effector region of NA appositions. As a result, the extracellular concentration of NA rises and facilitates NA synaptic activity at these two sites. For 1 mg/kg of PIP, the balance between the different mechanisms (enhanced NA release, reuptake and cross-talk) simultaneously at work maintains an extracellular concentration of NA which favours collateral inhibition, thereby decreasing PS. This decrease is more limited than that following administration of a drug such as DMI because reuptake is a powerful inactivation mechanism and partially compensates for the enhanced NA release. For higher doses of PIP, the blockade of postsynaptic α_2 -adrenoceptors within the LC and other NA nuclei of the brain stem is enhanced proportionally more than that of the presynaptic receptors. In other words, with increasing doses, progressive competition between PIP and endogenous NA takes place at the level of α_2 -adrenoceptors, resulting in a decrease of collateral inhibition. The balance is shifted in favour of the effector region and PS facilitation is observed. This facilitation is suppressed if the synthesis of NA is limited, indicating that it is dependent on the amount of the endogenous transmitter present.

Variations in collateral inhibition can also be responsible for the rebound without previous debt

which we have described after a small dose of clonidine. Collateral inhibition becomes less effective as a consequence of clonidine induced desensitization of α_2 -adrenoceptors and PS partially escapes its control.

Possible relationships between NA cell activity and PS

The data reviewed up to this point confront us with a major contradiction. Electrophysiological recordings of unit activity in the LC indicate that NA cells become silent during PS, which seems to rule out their active participation in the executive mechanism of this stage. In contrast, the results of pharmacological experiments suggest that when NA synaptic transmission is impaired, the production of PS is hindered, whereas when this transmission is facilitated, the production of PS is enhanced. Other investigations have also reached the same conclusion (Leppavuori & Putkonen, 1980; Leppavuori, 1980; Radulovacki *et al.*, 1980; Radulovacki *et al.*, 1981). Non-specific toxic effects could still be invoked, to explain, for instance, the decrease of PS obtained after high doses of AMPT, but no support is provided by available evidence (Kafi *et al.*, 1977). The modifications of PS induced by the drugs reported here could also be due to another and still unknown effect. However, within the limits of our present knowledge, the only common mechanism of action of these components involves alpha-receptors.

Two main lines of thinking emerged from the data reviewed here. Historically, the first proposed that the activity of brain CA systems, and probably mainly NA cells, is related to the maintenance of PS; PS executive NA neurons of the LC were said to be triggered by another mechanism in which 5-HT is involved, with a probable cholinergic link (Jouvet, 1969, 1972). Since this interpretation did not appear to be supported by later evidence, a second theory was proposed. In its most typical formulation, this view is diametrically opposed to the first and states that there is a negative correlation between PS and brain NA activity. According to this view, PS is inhibited by an activation of NA neurons and can appear only when they decrease or stop firing (Hartmann *et al.*, 1971; Hartmann & Schildkraut, 1973; Oswald *et al.*, 1975; Hobson *et al.*, 1976). An attenuated form of this interpretation maintains that the activity of brain NA neurons is inessential for PS (Marantz *et al.*, 1968; Branchey & Kissin, 1970). Another development along the same line of reasoning proposed that the function of PS is actually to maintain the availability of the neurotransmitter supposedly depleted during wakefulness and thus to repair synaptic fatigue due to CA depletion (Stern & Morgane, 1974). To summarize, unit activity of NA cells in the LC suggests a negative relationship with PS production, whereas pharmacological data suggest a positive relationship. No definitive interpretation

can account at present for these conflicting data. It is not possible simply to discard one of these alternatives; the results of electrophysiological and pharmacological experiments probably represent the same facts observed from different view points.

The particular anatomical distribution of NA cells in the brain stem may be critical for the regulation of wakefulness and PS. A majority of cells is densely packed in the lateral part of the pontine tegmentum and forms the core of the LC, whereas the remaining cells are more loosely dispersed over extensive regions from the medulla to the midbrain (Grzanna & Molliver, 1980). It can reasonably be supposed that in the core of the nucleus the number of NA terminals per unit volume is higher, and therefore that the extracellular concentration of the transmitter rises to higher levels than in other locations where cells are scattered throughout the tissue. The consequence is a higher degree of collateral inhibition where cells are dense than where they are dispersed. It is conceivable that this inhibition is sufficient to stop completely the activity of NA cells in the LC during PS, as has been observed by unit activity recording (Aston-Jones & Bloom, 1981). However, authors presenting the latter data noted that at the edge of the nucleus the rate of unit discharges tended to be higher than at its centre. Although they gave no interpretation of this finding, it can be proposed that cells in this marginal situation are less efficiently controlled by collateral inhibition. This is probably not yet enough for PS generation and it may well be that NA cells in the entire LC are essentially inactive during this stage. In contrast, this may not be the case for NA cells dispersed outside the LC in the other NA nuclei of the brain stem, in which they may escape more successfully collateral inhibition. Therefore, these cells may be active during PS, although still under partial control of collateral inhibition, as suggested by pharmacological observations. According to this view, the degree of inhibition could vary as a function of the topographical density of NA cells. The activity of these cells during PS could be a reciprocal function of inhibition: densely packed cells are silent whereas scattered cells only are active.

This interpretation could reconcile electrophysiological and pharmacological data. It could also bring into agreement phylogeny and ontogeny. It is well known that PS is especially developed in mammals and much less represented or absent in lower species. However, this is an instance in which ontogeny does not apparently recapitulate phylogeny since in human as well as in many mammalian new-borns, PS is much more abundant than in the adult individuals of the same species (Roffwarg *et al.*, 1966; Petre-Quadens, 1967). Since animals less mature at birth tend to have more PS than others, it is possible that in them the pathways or receptors mediating collateral inhibition are not fully developed. In the early period of life, the

maturation of these systems progressively enslaves PS until the adult tight control is acquired.

The proposed role of collateral inhibition in PS regulation can help in understanding the rebound phenomenon which has remained mysterious until now. It is well known that withdrawal of anti-depressant drugs results in marked increase of PS production (Dunleavy *et al.*, 1972; Passouant *et al.*, 1972). It has been established recently that these drugs down-regulate α_2 -adrenoceptors (Crews & Smith, 1978; Spyraiki & Fibiger, 1980; Smith *et al.*, 1981; Charney *et al.*, 1981; Siever *et al.*, 1981). Hypo-sensitivity of these receptors could conceivably decrease the efficiency of collateral inhibition leaving PS free partially to escape its control.

There are similarities between wakefulness and PS. Both represent a state of intense activation of the central nervous system. The EEG in these two states is very similar in most animal species and shows little difference in man. The activity of many units throughout the central nervous system is at the same level in wakefulness and in PS. In the course of evolution, wakefulness and PS may have proceeded by differentiation of a common primitive state of arousal and, accordingly, mammalian PS could be regarded as a remainder of this primitive wakefulness now embedded in sleep and devoted to another function. Collateral inhibition is perhaps a critical factor in this differentiation and its anatomical substrate may reach its full development only in mammals. Comparative topographical studies of the NA cells of the brain stem in animals at different levels of evolution could indicate whether this hypothesis is reasonable.

One important issue remains unexplained by this hypothesis: the ultradian PS rhythm. In addition, if it is true that a few NA cells are active during PS (Aston-Jones & Bloom, 1981; Sakai, personal communication) it is also evident that their activity is considerably lower than in any other state of alertness, and possibly insufficient to sustain the realization of PS. Future research will have to look for the activity of dispersed NA cells in the brain stem, carefully identified by their conduction times or by precise histofluorescence techniques.

We have indicated that α -adrenoceptors are also present on non-NA terminals. Accordingly, the α -adrenoceptor agonists or antagonists could modify PS by influencing non-NA cells. This is, however, not very likely since several of these effects are modified by alteration of the supply of endogenous transmitter or by limitation of the transmitter synthesis capacity of the cell. Therefore, the activity of at least some alpha-antagonists depends on the presence of endogenous NA.

Another hypothesis, not incompatible with the role of collateral inhibition and the topographical variation in NA cell density within the brain stem, could in addition account for the ultradian rhythm of PS. The

activity of NA cells could be critical for PS generation, not during the stage itself, but before it occurs. In this case, NA would not be directly responsible for PS realization, but nevertheless positively related to it and participating in a chain of events ultimately leading to PS appearance.

This hypothesis is not incompatible with, but does not necessarily require, the idea of a permissive role of NA neurons in PS realization. However, the precise relationship frequently observed between the onset of PS and the cessation of firing of units with conduction velocity compatible with NA cells (Sakai, personal communication) lends some weight to this possibility. Accordingly, PS preparation would be positively linked to NA cell activity, but actual PS realization would be negatively related to this activity.

The role of indolaminergic neurones

Recent experiments in the laboratory of Jouvet also support the idea of a factor, still unidentified, necessary for PS realization and synthesized before PS onset. It is known that treatment of cats with *p*CPA results in total insomnia lasting for several days; during recovery, PS reappears progressively without rebound (Mouret *et al.*, 1967). Peripheral or intraventricular administration of 5-HTP reverses *p*CPA insomnia (Mouret *et al.*, 1967; Koella *et al.*, 1968). The time course of this reversal is characteristic: 5-HTP blocks immediately *p*CPA-induced PGO waves and restores quiet sleep in about 30 min and PS only after 1 h. However, *p*CPA does not prevent the PS rebound which normally follows instrumental deprivation, but suppresses quiet sleep, thus inducing a narcoleptic syndrome in the cat. Cerebrospinal fluid (CSF) of PS deprived cats transferred to other cats with *p*CPA-induced insomnia restores PS. These data are compatible with the existence in CSF of a hypnogenic factor active in the absence of 5-HT. The synthesis of this factor may be under 5-HT control but its direct effects on PS realization do not require the presence of 5-HT (Sallanon, 1981).

Chloramphenicol induces a total suppression of PS in the cat (Petitjean *et al.*, 1979) and in the mouse (Kitahama & Valatx, 1975). This agent does not prevent, but delays by about 6 h, the PS rebound following deprivation by the flower pot technique (Sallanon, 1981). Thus, chloramphenicol does not prevent synthesis and accumulation of the hypothetical PS factor, but probably interferes with its utilization.

The choroid plexus may participate in the synthesis of this factor since its metabolism is high during sleep (Bobillier *et al.*, 1980) and the intraventricular injection of extracts of choroid plexus restores PS in *p*CPA treated animals, though with a longer latency than that following the transfer of CSF (Sallanon, 1981).

Numerous 5-HT terminals are found in the ependyma (Richards *et al.*, 1973; Lorez *et al.*, 1975) and 5-HT receptors have been identified in the choroid plexus (Meibach *et al.*, 1980).

The rebound of PS following a small dose of CLN depends on the abundance of slow wave sleep. This could suggest that if a PS factor really exists, its synthesis could be linked partially to the activity of the slow wave generating system. On the other hand, the hypothesis of a PS factor also requires that it is consumed during PS realization. An analysis of PS cycle in terms of appetitive or consummatory phase has been proposed by Tissot (1966), and could help to elucidate which is the component of quiet sleep more specifically involved in PS preparation. This component could be related to stage 2 since this author found an antagonism between slow wave sleep and PS, but a synergy between stage 2 and PS.

Differences between PS and waking: possible modulation of α_2 -adrenoceptor sensitivity

If NA cells participate in the regulation of both PS and waking, they appear to function differently in these two states of alertness. Their rate of firing is high in waking and very low or completely absent in PS; field potentials are associated with firing in wakefulness, but dissociated from unit activity during PS. This dissociation may reflect strong tonic inhibition of NA neurones preventing soma discharges (Aston-Jones & Bloom, 1981). These authors postulate that these cells receive strong inhibitory afferents during PS. We postulate further that α_2 -adrenoceptors are involved in this inhibition, which may be related to collateral inhibition, without excluding other inhibitory influences. A modulation of the sensitivity of α_2 -adrenoceptors mediating collateral inhibition could account in part for the different activity of NA cells in waking and in PS. Conceivably, if the sensitivity of α_2 -adrenoceptors is high, collateral inhibition is powerful and maintains NA cells activity under tight control; in contrast, if this sensitivity is low, collateral inhibition is much weaker and may release the activity of these cells.

Recent experiments in our laboratory support this view. The effect of clonidine over a large dose range was examined in the rat (Table 3) (Kafi & Gaillard, unpublished). The experiments extend previous results reported elsewhere (Gaillard *et al.*, 1982). Clonidine induced a simple dose-dependent decrease of PS, with significant modification after doses equal to or higher than 0.01 mg/kg. Neither at this dose nor at 0.02 mg/kg was waking influenced. After 0.04 mg/kg, however, waking was significantly enhanced; after the next two doses, 0.08 and 0.16 mg/kg, this parameter was still enhanced, but not significantly so because of larger variability of individual values. Finally, after 0.32 mg/kg the amount of waking was

Table 3 Effect of various doses of clonidine on waking and paradoxical sleep in the rat. The animals received the drug i.p. at 08.00 h and were recorded for the next 11 h.

Waking and PS were calculated as in Table 2

	n	Waking (%)	PS (%)
Clonidine 0 (mg/kg)	16	31 ± 6	16 ± 3
0.0025	7	30 ± 6	17 ± 2
0.005	3	32 ± 3	17 ± 2
0.01	8	33 ± 9	12 ± 3**
0.02	6	31 ± 6	9 ± 2***
0.04	3	40 ± 4*	5 ± 2***
0.08	3	42 ± 15	6 ± 7***
0.16	3	37 ± 5	2 ± 3***
0.32	3	29 ± 9	0

Waking and PS were calculated as in Table 2.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.005$ (two-tailed Student's *t*-test).

identical to control values (Table 3). A more precise picture was obtained when the general trend of waking was examined. After 0.04 mg/kg of clonidine, the production of waking was facilitated (i.e. the slope of the function was steeper than in controls) in the first 150 min of the recording; subsequently, this production returned to control value. This particular evolution conferred to the general trend a parabolic shape with a negative curvature. After 0.08 and 0.16 mg/kg, a similar facilitation was seen, but it occurred later in the recording period. After the dose of 0.32 mg/kg, the production of waking was initially lower than in controls, but right at the end of the recording a facilitation appeared, bringing the final point, which represents the total amount of waking for the full recording period, exactly to the control value. Thus, not only was waking facilitated after doses of clonidine between 0.04 and 0.16 mg/kg, but for higher doses this facilitation appeared later during the recording period. This particular dose-effect relationship suggests that waking facilitation may be related to a critical concentration of clonidine in the brain: it only occurs when this critical concentration is reached. This concentration is attained shortly after administration of 0.04 mg/kg of clonidine. If larger doses are given, the brain concentration of CLN presumably rises initially to higher levels and drops only later to the critical wake facilitating concentration. After the dose of 0.32 mg/kg, the initial concentration of clonidine in the brain is enough to depress the production of waking and reaches the concentration critical for waking facilitation only about 8 h later.

The fact that the doses of clonidine modifying PS are lower than those modifying waking suggests that α_2 -adrenoceptors are more sensitive in the former than in the latter stage. According to our model, clonidine probably acts simultaneously at three different sites: (1) on presynaptic α_2 -adrenoceptors in

NA nuclei, decreasing NA release and lowering collateral inhibition; (2) on postsynaptic α_2 -adrenoceptors on NA cell bodies, decreasing the firing of NA cells; and (3) on presynaptic α_2 -adrenoceptors in remote NA appositions. We have seen above the reason why it is likely that CLN acts in NA nuclei at lower doses than on remote NA terminals. In PS, the sensitivity of pre- and postsynaptic α_2 -adrenoceptors is probably equilibrated in such a way that CLN does not induce any PS increase for any dose. In wakefulness, in addition to overall decrease, α_2 -adrenoceptor sensitivity may be proportionally lower at postsynaptic than at presynaptic α_2 -adrenoceptors in NA nuclei of the brain stem. Thus, the facilitation of waking may result from decrease of collateral inhibition, whereas the decrease observed after higher doses suggests that this dose is necessary to stimulate postsynaptic α_2 -adrenoceptors in NA nuclei.

Experiments with various doses of piperoxane also support the conclusion that PS is under tighter control than waking by a mechanism in which α_2 -adrenoceptors are involved (Gaillard & Kafi, 1983; see also Table 2).

The properties of cell membranes which we have reviewed above provide ample possibilities for physiological modulation of receptor sensitivity. The modulators supposedly involved are unknown; however, this system may not function symmetrically in both directions. If increase of sensitivity is slower than decrease, we could understand why it takes more time to fall asleep than to wake up.

Conclusion

A complete theory of PS regulation cannot be proposed at this time. The data reviewed here suggest the possible involvement of three factors: collateral

inhibition on NA cells, variation of topographical density of NA cell bodies in the brain stem and modulation of the sensitivity of α_2 -adrenoceptors. If pharmacological experiments suggest a positive relationship between NA activity and PS production, the temporal relationship between these two events may not be as simple as previously believed.

NA may have a mechanism of activity other than that usually considered and may not be an executive neurotransmitter like acetylcholine at the neuromuscular junction. It may participate in PS generation as a link in a chain of events that ultimately leads to PS realization. The results of experiments are

sometimes difficult to understand because this chain probably consists of many links and temporal relationships between causes and effects may be difficult to establish. Finally, the amount of a neurotransmitter may not be the only factor of importance. Complex mechanisms controlling receptor sensitivity and metabolism can be expected to play a crucial role in the regulation of the states of alertness. The interpretation of experimental data is still tentative, but it is hoped that this review will inspire future experimentation aimed at testing the possible participation of the factors suggested here.

References

- AGHAJANIAN, G.K., CEDARBAUM, J.M. & WANG, R.Y. (1977). Evidence for norepinephrine-mediated collateral inhibition of locus coeruleus neurons. *Brain Res.*, **136**, 570–577.
- AHLQUIST, R.P. (1948). A study of the adrenergic receptors. *Am. J. Physiol.*, **153**, 586–600.
- AMATRUDA, T.T., BLACK, D.A., MCKENNA, T.M., MCCARLEY, R.W. & HOBSON, J.A. (1975). Sleep cycle control and cholinergic mechanisms: Differential effects of carbachol injections at pontine brain stem sites. *Brain Res.*, **98**, 501–515.
- ANDEN, N.-E., CORRODI, H., FUXE, K., HOKFELT, B., HOKFELT, T., RYDIN, C. & SVENSSON, T. (1970). Evidence for a central noradrenaline receptor stimulation by clonidine. *Life Sci.*, **9**, 513–523.
- ANDEN, N.-E., GRABOWSKA, M. & STRÖMBOM, U. (1976). Different alpha-adrenoreceptors in the central nervous system mediating biochemical and functional effects of clonidine and receptor blocking agents. *Naunyn-Schmiedeberg's Arch. Pharmacol.*, **292**, 43–52.
- ANDERSON, R.G.W., BROWN, M.S. & GOLSTEIN, J.L. (1977). Role of the coated endocytic vesicle in the uptake of receptor-bound low density lipoprotein in human fibroblasts. *Cell*, **10**, 351–364.
- APPEL, S.H., ANWYL, R., McADAMS, M.W. & ELIAS, S. (1977). Accelerated degradation of acetylcholine receptor from cultured rat myotubes with myasthenia gravis sera and globulins. *Proc. Nat. Acad. Sci. USA*, **74**, 2130–2134.
- ARIENS, E.J. & SIMONIS, A.M. (1976). Receptors and receptor mechanisms. In *Beta-adrenoceptor blocking agents*, eds. Saxena, P.R. & Forsyth, R.P., pp. 3–27. Amsterdam: North Holland Publishing Co.
- ASTON-JONES, G. & BLOOM, F.E. (1981). Activity of norepinephrine-containing locus coeruleus neurons in behaving rats anticipates fluctuations in the sleep-waking cycle. *J. Neurosci.*, **1**, 876–886.
- AXELROD, D., BAUER, H.C., STYA, M. & CHRISTIAN, C.N. (1981). A factor from neurons induces partial immobilization of nonclustered acetylcholine receptors on cultured muscle cells. *J. cell. Biol.*, **88**, 459–462.
- BANNERJEE, S.P., KUNG, L.S., RIGGI, S.J. & CHANDA, S.K. (1977). Development of beta-adrenergic receptor subsensitivity by antidepressants. *Nature (Lond.)*, **268**, 445–456.
- BARTHOLINI, G., KELLER, H.H. & PLETSCHER, A. (1973). Effect of neuroleptics on endogenous norepinephrine in rat brain. *Neuropharmacology*, **12**, 751–756.
- BASU, S.K., GOLDSTEIN, J.L., ANDERSON, R.G.W. & BROWN, M.S. (1981). Monensin interrupts the recycling of low density lipoprotein receptors in human fibroblasts. *Cell*, **24**, 493–502.
- BAUMANN, P.A. & KOELLA, W.P. (1980). Feedback control of noradrenaline release as a function of noradrenaline concentration in the synaptic cleft in cortical slices of the rat. *Brain Res.*, **189**, 437–448.
- BEARER, E.L. & FRIEND, D.S. (1980). Anionic lipid domains: correlation with functional topography in a mammalian cell membrane. *Proc. Nat. Acad. Sci. USA*, **77**, 6601–6605.
- BISWAS, B. & JONSSON, G. (1981). Reversal of noradrenaline denervation-induced increase of beta-adrenoceptor binding in rat neocortex by noradrenaline infusion. *Eur. J. Pharmacol.*, **69**, 341–346.
- BLASIE, J.K. (1972). The location of photopigment molecules in the cross-section of frog retinal receptor disc membranes. *Biophys. J.*, **12**, 191–204.
- BOBILLIER, P., SEGUIN, S., PETITJEAN, F., BUDA, C., SALVERT, D., JANIN, M., CHOUVET, G., SOUCHIER, C. & JOUVET, M. (1980). Physiologie animale.—Augmentation de la consommation cérébrale locale de glucose dans les plexus choroïdes au cours du sommeil lent chez le chat. *CR Acad. Sci. Paris*, **291**, 91–96.
- BOROCHOV, H. & SHINITZKY, M. (1976). Vertical displacement of membrane proteins mediated by changes in micro-viscosity. *Proc. Nat. Acad. Sci. USA*, **73**, 4526–4530.
- BRAESTRUP, C. (1974). Effects of phenoxybenzamine, aceperone and clonidine on the level of 3-methoxy-4-hydroxyphenylglycol (MOPEG) in rat brain. *J. Pharm. Pharmacol.*, **26**, 139–141.
- BRANCHEY, M. & KISSIN, B. (1970). The effect of alpha-methyl-paratyrosine on sleep and arousal in the rat. *Psychol. Sci.*, **19**, 281–282.
- BROWN, J., DOXY, J.C. & HANDLEY, S. (1980). Effects of alpha-adrenoceptor agonists and antagonists and of antidepressant drugs on pre- and postsynaptic alpha-adrenoceptors. *Eur. J. Pharmacol.*, **67**, 33–40.
- BRYAN, L.J., COLE, J.J., O'DONNELL, S.R. & WANSTALL, J.C. (1981). A study designed to explore the hypothesis

- that beta-1 adrenoceptors are 'innervated' receptors and beta-2 adrenoceptors are 'hormonal' receptors. *J. Pharmac. exp. Ther.*, **216**, 395-400.
- BUGUET, A. (1969). Monoamines et sommeil. V. Etude des relations entre les structures monoaminergiques du pont et les pointes ponto-geniculo occipitales du sommeil. *Thèse Médecine, Lyon*, Tixier Ed., 214 pp.
- CERDARBAUM, J.M. & AGHAJANIAN, G.K. (1977). Catecholamine receptors on locus coeruleus neurons: Pharmacological characterization. *Eur. J. Pharmac.*, **44**, 375-385.
- CERDARBAUM, J.M. & AGHAJANIAN, G.K. (1978). Activation of locus coeruleus neurons by peripheral stimuli: modulation by a collateral inhibitory mechanism. *Life Sci.*, **23**, 1383-1392.
- CHABRE, M. (1975). X-Ray diffraction studies on retinal rods. I. Structure of the disc membrane, effect of illumination. *Biochim. biophys. Acta*, **383**, 322-335.
- CHARNEY, D.S., MENKES, D.B. & HENINGER, G.R. (1981). Receptor sensitivity and the mechanism of action of antidepressant treatment. *Arch. gen. Psychiat.*, **38**, 1160-1180.
- CHERRY, R.J. (1979). Rotational and lateral diffusion of membrane proteins. *Biochim. biophys. Acta*, **559**, 289-327.
- CHEUNG, W.Y. (1980). Calmodulin plays a pivotal role in cellular regulation. *Science*, **207**, 19-27.
- CHU, N.-S. & BLOOM, F.E. (1974). Activity patterns of catecholamine-containing pontine neurons in the dorso-lateral tegmentum of unrestrained cats. *J. Neurobiol.*, **5**, 527-544.
- CHUANG, D.-M., KINNIER, W.J., FARBER, L. & COSTA, E. (1980). A biochemical study of receptor internalization during beta-adrenergic receptor desensitization in frog erythrocytes. *Mol. Pharmac.*, **18**, 348-355.
- COOPER, B., HANDIN, R.I., YOUNG, L.H. & ALESANDER, R.W. (1978). Agonist regulation of the human platelet alpha-adrenergic receptor. *Nature (Lond.)*, **274**, 703.
- CREWS, F.T. & SMITH, C.B. (1978). Presynaptic alpha-receptor subsensitivity after long-term antidepressant treatment. *Science*, **202**, 322-324.
- DAHLSTROM, A. & FUXE, K. (1964). Evidence for the existence of monoamine neurons in the central nervous system. I. Demonstration of monoamines in the cell bodies of brain stem neurons. *Acta physiol. scand.*, **62** (suppl.) 232.
- DAVIS, J.N., OLENDER, E., MAURY, W. & McDANIEL, R. (1980). Alpha-adrenergic regulation of cholinergic responses in rat parotid acinar cells. *Mol. Pharmac.*, **18**, 356-361.
- DE PETRIS, S. (1974). Inhibition and reversal of capping by cytochalasin B, vinblastine and colchicine. *Nature*, **250**, 54-56.
- DISMUKES, K., DE BOER, A.A. & MULDER, A.H. (1977). On the mechanism of alpha-receptor mediated modulation of ³H-noradrenaline release from slices of rat brain neocortex. *Naunyn-Schmiedeberg's Arch. Pharmac.*, **299**, 115-122.
- DRACHMAN, D.B., ANGUS, C.W., ADAMS, R.N., MICHELSON, J.D. & HOFFMAN, G.J. (1978). Myasthenic antibodies cross-link acetylcholine receptors to accelerate degradation. *New Engl. J. Med.*, **298**, 1116-1122.
- DRACHMAN, D.B., KAO, L., ANGUS, C.W. & MURPHY, A. (1977). Effect of myasthenic immunoglobulin on acetylcholine receptors of cultured muscle. *Ann. Neurol.*, **1**, 504.
- DUNLEAVY, D.L.F., BREZINOVA, V., OSWALD, I., McLEAN, A.W. & TINKER, M. (1972). Changes during weeks in effects of tricyclic drugs on the human sleeping brain. *Br. J. Psychiat.*, **120**, 663-672.
- DUNLEAVY, D.L.F., MacLEAN, A.W. & OSWALD, I. (1971). Debrisoquine, guanethidine, propranolol and human sleep. *Psychopharmacologia (Berl.)*, **21**, 101-110.
- ENGBERG, G., ELAM, M., & SVENSSON, T.H. (1982). Clonidine withdrawal: activation of brain noradrenergic neurons with specifically reduced alpha₂-receptor sensitivity. *Life Sci.*, **30**, 235-243.
- ENGELMAN, D.M. & STEITZ, T.A. (1981). The spontaneous insertion of proteins into and across membranes: the helical hairpin hypothesis. *Cell*, **23**, 411-422.
- EXTON, J.H. (1980). Mechanisms involved in alpha-adrenergic phenomena: role of calcium ions in actions of catecholamines in liver and other tissues. Editorial Review, *Am. physiol. Soc.*: E3-E12.
- FARMBROUGH, D.M. (1979). Control of acetylcholine receptors in skeletal muscle. *Physiol. Rev.*, **59**, 165-227.
- FRANKLIN, K.B.J. & HERBERT, L.J. (1977). Presynaptic alpha-adrenoceptors: The depression of self-stimulation by clonidine and its restoration by piperoxane but not by phentolamine or phenoxybenzamine. *Eur. J. Pharmac.*, **43**, 33-38.
- FRYE, C.D. & EDIDIN, M. (1970). The rapid intermixing of cell surface antigens after formation of mouse-human heterokaryons. *J. cell. Sci.*, **7**, 313.
- GAILLARD, J.-M. (1979). Temporal organization of human sleep: general trends of sleep stages and their ultradian cyclic components. *Encéphale*, **V**, 71-93.
- GAILLARD, J.-M., BARTHOLINI, G., HERKERT, B. & TISSOT, R. (1974). Involvement of 5-hydroxytryptamine in the cortical synchronization induced by L-DOPA in the rabbit. *Brain Res.*, **68**, 344-350.
- GAILLARD, J.-M., IORIO, G., KAFI, S. & BLOIS, R. (1983). Paradoxical sleep rebound without previous debt: the effect of minute doses of clonidine in man. *Sleep*, **6**, 60-66.
- GAILLARD, J.-M. & KAFI, S. (1979). Involvement of pre- and postsynaptic receptors in catecholaminergic control of paradoxical sleep in man. *Eur. J. clin. Pharmac.*, **15**, 83-89.
- GAILLARD, J.-M. & KAFI, S. (1983). Effect of alpha-adrenoceptor antagonist piperoxane on rat waking and paradoxical sleep. *Sixth European Congress of Sleep Research*, March 23-26, 1982, Zürich, 149-150. Basel: Karger.
- GAILLARD, J.-M., KAFI, S. & JUSTAFRE, J.-C. (1982). On the role of brain alpha-adrenergic systems in the production of paradoxical sleep. *Encéphale*, **8**, 413-434.
- GAILLARD, J.-M., SCHAEPI, R. & TISSOT, R. (1969). Potentialisation des effets centraux de la DOPA après inhibition sélective de la décarboxylase extracérébrale. *Arch. int. Pharmacodyn.*, **180**, 423-437.
- GEORGE, R., HASLETT, W.L. & JENDEN, D.J. (1964). A cholinergic mechanism in the brain stem reticular formation: induction of paradoxical sleep. *Int. J. Pharmac.*, **3**, 451-552.
- GILLIN, J.C., MENDELSON, W.B., SITARAM, N. & WYATT, R.J. (1978). The neuropharmacology of sleep and wakefulness. *Ann. Rev. Pharmac. Tox.*, **18**, 563-579.
- GILLIN, J.C., SITARAM, N. & DUNCAN, W.C. (1979).

- Muscarinic supersensitivity: A possible model for the sleep disturbance of primary depression? *Psychiatry Res.*, **1**, 17-22.
- GITLER, C. (1976). On the nature of the lipid-protein interactions in biological membranes. In *The enzymes of biological membranes*. Physical and chemical techniques. Vol. 1, pp. 229-244. ed Matonosi, A. New York: John Wiley & Sons.
- GOLDSTEIN, J.C., ANDERSON, R.G.W. & BROWN, M.S. (1979). Coated pits, coated vesicles, and receptor-mediated endocytosis. *Nature (Lond.)*, **279**, 679-685.
- GORDON-WEEKS, P.R., JONES, D.H., GRAY, E.G. & BARRON, J. (1981). Trypsin separates synaptic junctions to reveal pre- and post-synaptic concanavalin A receptors. *Brain Res.*, **219**, 224-230.
- GOTHERT, M., HUTH, H. & SCHLICKE, E. (1981). Characterization of the receptor subtype involved in alpha-adrenoceptor-mediated modulation of serotonin release from rat brain cortex slices. *Naunyn-Schmiedeberg's Arch. Pharmacol.*, **317**, 199-203.
- GROVES, P.M. & WILSON, C.J. (1980a). Fine structure of rat locus coeruleus. *J. comp. Neurol.*, **193**, 841-852.
- GROVES, P.M. & WILSON, C.J. (1980b). Monoaminergic presynaptic axons and dendrites in rat locus coeruleus seen in reconstructions of serial sections. *J. comp. Neurol.*, **193**, 853-862.
- GRZANNA, R. & MOLLIVER, M.E. (1980). Cytoarchitecture and dendritic morphology of central noradrenergic neurons. In *The reticular formation revisited*, eds Hobson, J.A. & Brazier, M.A.B., pp. 83-97. New York: Raven Press.
- HAIGLER, H.J. & AGHAJANIAN, G.K. (1977). Serotonin receptors in the brain. *Fed. Proc.*, **36**, 2159-2164.
- HARTMANN, E., CHUNG, R., DRASKOCZY, P.R. & SCHILDKRAUT, J.J. (1971). Effects of 6-hydroxydopamine on sleep in the rat. *Nature (Lond.)*, **233**, 425-427.
- HARTMANN, E. & SCHILDKRAUT, J.J. (1973). Desynchronized sleep and MHPG excretion: an inverse correlation. *Brain Res.*, **61**, 412-416.
- HATA, F., TAKEYASU, K., UCHIDA, S. & YOSHIDA, H. (1980). Increase in response and number of alpha-adrenoceptors of rat vas deferens on brief pretreatment with an alpha-agonist. *Eur. J. Pharmacol.*, **67**, 193-199.
- HAZUM, E., CHANG, K.-J. & CUATRECASAS, P. (1981). Receptor redistribution induced by hormones and neurotransmitters: possible relationships to biological functions. *Neuropeptides*, **1**, 217-230.
- HENRIKSEN, S.J. & DEMENT, W.C. (1972). Effects of chronic intravenous administration of 1-alpha-methyltyrosine on sleep in the cat. In *Sleep Research*, Brain Information Service Brain Research Institute, eds Chase, M.H., Stern, W.C. & Walter, P.L. UCLA, Los Angeles, vol. I, 55 (Abstr.).
- HERON, D.S., SHINITZKY, M., HERSHKOWITZ, M. & SAMUEL, D. (1980). Lipid fluidity markedly modulates the binding of serotonin to mouse brain membranes. *Proc Nat. Acad. Sci. USA*, **77**, 7463-7467.
- HILAKIVI, I., MAKELA, J., LEPPAVUORI, A. & PUTKONEN, P.T.S. (1978). Effects of two adrenergic beta-receptor blockers on the sleep cycle of the cat. *Med. Biol.*, **56**, 138-143.
- HIRATA, F., TALLMAN, J.F. Jr, HENNEBERRY, R.C., MALLORGA, P., STRITTMATTER W.J. & AXELROD, J. (1980). Regulation of beta-adrenergic receptors by phospholipid methylation. In *Receptors for Neurotransmitters and Peptide Hormones*, eds Pepeu, G., Kuhar, M.J. & Enna, S.J., pp. 91-97. New York: Raven Press.
- HOBSON, J.A., McCARLEY, R.W. & McKENNA, T.M. (1976). Cellular evidence bearing on the pontine brainstem hypothesis of desynchronized sleep control. *Prog. Neurobiol.*, **6**, 280-376.
- HOBSON, J.A., McCARLEY, R.W., PIVIK, R.T. & FREEDMAN, R. (1974). Selective firing by cat pontine brain stem neurons in desynchronized sleep. *J. Neurophysiol.*, **37**, 497-511.
- HOBSON, J.A., McCARLEY, R.W. & WYSINSKY, P.W. (1975). Sleep cycle oscillation: Reciprocal discharge by two brainstem neuronal groups. *Science*, **189**, 55-58.
- HÖKFELT, T. (1967). On the ultrastructural localization of noradrenaline in the central nervous system of the rat. *Z. Zellforsch.*, **79**, 110-117.
- HOLTZMAN, E. & MERCURIO, A.M. (1980). Membrane circulation in neurons and photoreceptors: some unresolved issues. *Int. Rev. Cytol.*, **67**, 1-67.
- INSEL, P.A., NIRENBERG, P., TURNBULL, J. & SHATTIL, S.J. (1978). Relationships between membrane cholesterol, alpha-adrenergic receptors, and platelet function. *Biochemistry*, **17**, 5269-5274.
- ISKANDER, T.N. & KAEHLING, R. (1970). Catecholamines, a dream sleep model, and depression. *Am. J. Psychiat.*, **127**, 43-50.
- JOHANSSON, O., HÖKFELT, T., PERNOW, B., JEFFCOATE, S.L., WHITE, N., STEINBUSCH, H.W.M., VERHOFSTAD, A.A.J., EMSON, P.C. & SPINDEL, E. (1981). Immunohistochemical support for three putative transmitters in one neuron: coexistence of 5-hydroxytryptamine, substance P and thyrotropin releasing hormone-like immunoreactivity in medullary neurons projecting to the spinal cord. *Neuroscience*, **6**, 1857-1881.
- JONES, B.E. (1972). The respective involvement of noradrenaline and its deaminated metabolites in waking and paradoxical sleep: A neuropharmacological model. *Brain Res.*, **39**, 121-136.
- JONES, B.E. (1979). Elimination of paradoxical sleep by lesions of the pontine gigantocellular tegmental field in the cat. *Neurosci. Lett.*, **13**, 285-293.
- JONES, B., BOBILLIER, P. & JOUVET, M. (1969). Effets de la destruction des neurones contenant des catécholamines du mésencéphale sur le cycle veille-sommeil du chat. *CR Soc. Biol. (Paris)*, **163**, 176-180.
- JONES, B.E., HARPER, S.T. & HALARIS, A.E. (1977). Effects of locus coeruleus lesions upon cerebral monoamine content, sleep-wakefulness states and the response to amphetamine in the cat. *Brain Res.*, **124**, 473-496.
- JOUVET, M. (1962). Recherches sur les structures nerveuses et les mécanismes responsables des différentes phases du sommeil physiologique. *Arch. ital. Biol.*, **100**, 125-206.
- JOUVET, M. (1969). Biogenic amines and the states of sleep. *Science*, **163**, 32-41.
- JOUVET, M. (1972). The role of monoamines and acetylcholine containing neurons in the regulation of the sleep-waking cycle. *Ergebn. Physiol.*, **64**, 163-307.
- JUSTAFRE, J.-C. & GAILLARD, J.-M. (1981). Diminution du sommeil paradoxal chez le rat sous antidépresseur tricyclique: rôle de l'inhibition collatérale. *Rev. EEG Neurophysiol. clin.*, **11**, 228-235.

- KAFI, S., BOURAS, C., CONSTANTINIDIS, J. & GAILLARD, J.-M. (1977). Paradoxical sleep and brain catecholamines in the rat after single and repeated administration of alpha-methyl-paratyrosine. *Brain Res.*, **135**, 123–134.
- KAFI, S. & GAILLARD, J.-M. (1978). Biphasic effect of chlorpromazine on rat paradoxical sleep: a study of dose-related mechanisms. *Eur. J. Pharmac.*, **49**, 251–257.
- KAFI, S. & GAILLARD, J.-M. (1980). The effect of presynaptic alpha blocker yohimbine in rat paradoxical sleep. *Waking & Sleeping*, **4**, 131–137.
- KAFI, S. & GAILLARD, J.-M. (1981). Effect of alpha-stimulation by clonidine on the evolution of paradoxical sleep in the rat. *J. Psychiat. biol. Thé.*, **1**, 13–20.
- KAPLAN, J. (1981). Polypeptide-binding membrane receptors: analysis and classification. *Science*, **212**, 14–20.
- KEANE, P.E., CANDY, J.M. & BRADLEY, P.B. (1976). The role of endogenous catecholamines in the regulation of electrocortical activity in the 'encéphale isolé' cat. *Electroenceph. clin. Neurophysiol.*, **41**, 561–570.
- KING, C.D. & JEWETT, R.E. (1971). The effects of alpha-methyl-tyrosine on sleep and brain norepinephrine in cats. *J. Pharmac. exp. Ther.*, **177**, 188–195.
- KITAHAMA, M. & VALATX, J.L. (1975). Effets du chloramphénicol et du thiamphénicol sur le sommeil de la souris. *CR Soc. Biol. (Paris)*, **169**, 1522–1524.
- KLEINOGEL, H., SCHOLTYSIK, G. & SAYERS, A.C. (1975). Effects of clonidine and BS 100-141 on the EEG sleep pattern in rats. *Eur. J. Pharmac.*, **33**, 159–163.
- KOBINGER, W. & PICHLER, L. (1980). Investigation into different types of post- and presynaptic alpha-adrenoceptors at cardiovascular sites in rats. *Eur. J. Pharmac.*, **65**, 393–402.
- KOBINGER, W. & PICHLER, L. (1981). Alpha₁ and alpha₂-adrenoceptors subtypes: selectivity of various agonists and relative distribution of receptors as determined in rats. *Eur. J. Pharmac.*, **73**, 313–321.
- KOELLA, W.P., FELDSTEIN, A. & CZICMAN, J.S. (1968). The effect of parachlorophenylalanine on the sleep of cats. *Electroenceph. clin. Neurophysiol.*, **25**, 481–490.
- LACKIE, J.M. (1980). The structure and organization of the cell surface. In *Membrane structure and function 1*, ed. Bittar, E.E., pp. 73–102. New York: John Wiley & Sons.
- LAGUZZI, R., PEITJEAN, F., PUJOL, J.F. & JOUVET, M. (1972). Effets de l'injection intraventriculaire de 6-hydroxydopamine. II. Sur le cycle veille-sommeil du chat. *Brain Res.*, **48**, 295–310.
- LANGER, S.Z. (1977). Presynaptic receptors and their role in the regulation of transmitter release. *Br. J. Pharmac.*, **60**, 481–497.
- LAPIERRE, Y., BEAUDET, A., DEMIANCZUK, N. & DESCARRIES, L. (1973). Noradrenergic axon terminals in the cerebral cortex of rat. II. Quantitative data revealed by light and electron microscope radioautography on the frontal cortex. *Brain Res.*, **63**, 175–182.
- LEPPAVUORI, A. (1980). The effects of an alpha-adrenergic agonist or antagonist on sleep during blockade of catecholamine synthesis in the cat. *Brain Res.*, **193**, 117–128.
- LEPPAVUORI, A. & PUTKONEN, P.T.S. (1980). Alpha-adrenoceptive influences on the control of the sleep-waking cycle in the cat. *Brain Res.*, **193**, 95–115.
- LIDBRINK, P. (1974). The effect of lesions of ascending noradrenaline pathways on sleep and waking in the rat. *Brain Res.*, **174**, 19–40.
- LINDSTROM, J.M., SEYBOLD, M.E., LENNON, V.A., WITTINGHAM, S. & DRAKE, D.D. (1976). Antibody to acetylcholine receptor in myasthenia gravis: prevalence, clinical correlates, and diagnostic value. *Neurology (Minneapolis)*, **26**, 1054–1059.
- LOH, H.H. & LAW, P.Y. (1980). The role of membrane lipids in receptor mechanisms. *Ann. Rev. Pharmac. Tox.*, **20**, 201–234.
- LOREZ, H.P., PIERI, L. & RICHARDS, J.G. (1975). Disappearance of supra-ependymal 5-HT axons in the rat forebrain after electrolytic and 5,6-DHT-induced lesions of the medial forebrain bundle. *Brain Res.*, **100**, 1–12.
- MAEDA, T., ELDRIDGE, C., TOYAMA, S., OHNISHI, S.I., ELSON, E.L. & WEBB, W.W. (1979). Membrane receptor mobility changes by Sendai virus. *Exp. Cell Res.*, **123**, 333–343.
- MAGGI, A., U'PRICHARD, D.C. & ENNA, S.J. (1980a). Beta-adrenergic regulation of alpha₂-adrenergic receptors in the central nervous system. *Science*, **207**, 645–647.
- MAGGI, A., U'PRICHARD, D.C. & ENNA, S.J. (1980b). Differential effects of antidepressant treatment on brain monoaminergic receptors. *Eur. J. Pharmac.*, **61**, 91–98.
- MAJEWSKI, H. & RAND, M.J. (1981). An interaction between prejunctional alpha-adrenoceptors and prejunctional beta-adrenoceptors. *Eur. J. Pharmac.*, **69**, 493–498.
- MANDELL, A.J. & SPOONER, C.E. (1968). Psychochemical research studies in man. *Science*, **162**, 1442–1453.
- MARANTZ, R., LOVELL, R.A. & WHITEHEAD, P.K. (1968). Effect of alpha-methyltyrosine on the recovery from paradoxical sleep deprivation in the rat. *Commun. Behav. Biol.*, **2**, 161–164.
- MATSUYAMA, S., COINDET, J. & MOURET, J. (1973). 6-hydroxydopamine intracisternale et sommeil chez le rat. *Brain Res.*, **57**, 85–95.
- MATUS, A., DE PETRI, D. & RAFF, M.C. (1973). Mobility of concanavalin A receptors in myelin and synaptic membranes. *Nature New Biol.*, **244**, 278–279.
- MCCARLEY, R.W. & HOBSON, J.A. (1975). Neuronal excitability modulation over the sleep cycle: A structural and mathematical model. *Science*, **189**, 58–60.
- MEIBACH, R.C., MAAAYANI, S. & GREEN, J.P. (1980). Characterization and radioautography of H-LSD binding by rat brain slices *in vitro*: the effect of 5-hydroxytryptamine. *Eur. J. Pharmac.*, **67**, 371–382.
- MICHELL, R.H. & KIRK, C.J. (1981). Why is phosphatidylinositol degraded in response to stimulation of certain receptors? *Trends Pharmac. Sci.*, **2**, 86–89.
- MINNEMAN, K.P., PITTMAN, R.N. & MOLINOFF, P.B. (1981). Beta-adrenergic receptor subtypes: properties, distribution, and regulation. *Ann. Rev. Neurosci.*, **4**, 419–461.
- MISHRA, R., JANOWSKY, A. & SULSER, F. (1980). Action of mianserin and zimelidine on the norepinephrine receptor coupled adenylate cyclase system in brain: sub-sensitivity without reduction in beta-adrenergic receptor binding. *Neuropharmacology*, **19**, 983–987.
- MITLER, M.M. & DEMENT, W.C. (1974). Cataplectic-like behaviour in cats after micro-injections of carbachol in pontine reticular formation. *Brain Res.*, **68**, 335–343.
- MOURET, J., FROMENT, J.L., BOBILLIER, P. & JOUVET, M. (1967). Etude neuropharmacologique et biochimique

- des insomnies provoquées par la PCPA. *J. Physiol. (Paris)*, **59**, 463-464.
- NICOLSON, G.L. (1976). Transmembrane control of the receptors on normal and tumor cells. I. Cytoplasmic influence over cell surface components. *Biochim. biophys. Acta*, **457**, 57-108.
- OSWALD, I., THACORE, V.R., ADAM, K., BREZINOVA, V. & BURACK, R. (1975). Alpha-adrenergic receptor blockade increases human REM sleep. *Br. J. clin. Pharmacol.*, **2**, 107-110.
- PANKSEPP, J., JALOWIEC, J.E., MORGANE, P.J., ZOLOVICK, A.J. & STERN, W.C. (1973). Noradrenergic pathways and sleep-waking states in cats. *Exp. Neurol.*, **41**, 233-245.
- PAPAPHILIS, A. & DELICONSTANTINOS, G. (1980). Modulation of serotonergic receptors by exogenous cholesterol in the dog synaptosomal plasma membrane. *Biochem. Pharmacol.*, **29**, 3325-3327.
- PASSOUANT, P., CADILHAC, J. & RIBSTEIN, M. (1972). Les privations de sommeil avec mouvements oculaires par les anti-dépresseurs. *Rev. neurol.*, **127**, 173-192.
- PEARSE, B.M.F. (1976). Clathrin: a unique protein associated with intracellular transfer to membrane by coated vesicles. *Proc. Nat. Acad. Sci. USA*, **77**, 1255-1259.
- PESTRONK, A. & DRACHMAN, D.B. (1978). Motor nerve sprouting and acetylcholine receptors. *Science*, **199**, 1223-1225.
- PESTRONK, A., DRACHMAN, D.B., STANLEY, E.F., PRICE, D.L. & GRIFFIN, J.W. (1980). Cholinergic transmission regulates extrajunctional acetylcholine receptors. *Exp. Neurol.*, **70**, 690-696.
- PETITJEAN, J., BUDA, C., JANIN, M., DAVID, M. & JOUVET, M. (1979). Effets du chloramphénicol sur le sommeil du chat. Comparaison avec les thiamphénicol, l'érythromycine et l'oxytracycline. *Psychopharmacology*, **66**, 147-153.
- PETITJEAN, F. & JOUVET, M. (1970). Hypersomnie et augmentation de l'acide 5 hydroxy-indolacétique cérébral par lésion isthmique chez le chat. *CR Soc. Biol. (Paris)*, **164**, 2288.
- PETRE-QUADENS, O. (1967). Ontogenesis of paradoxical sleep in the human newborn. *J. neurol. Sci.*, **4**, 153-157.
- PIKE, L.J. & LEFKOWITZ, R.J. (1980). Use of cell fusion techniques to probe the mechanism of catecholamine-induced desensitization of adenylate cyclase in frog erythrocytes. *Biochim. biophys. Acta*, **632**, 354-365.
- POLLET, R.J. & LEVEY, G.S. (1980). Principles of membrane receptor physiology and their application to clinical medicine. *Ann. int. Med.*, **92**, 663-680.
- POO, M. (1979). Molecular movements of receptors on cell surface. *Biorheology*, **16**, 309-315.
- PUIZILLOUT, J.J., GAUDIN-CHAZAL, G., SAYADI, A. & VIGIER, D. (1981). Serotonergic mechanisms and sleep. *J. Physiol. Paris*, **77**, 415-424.
- PUTNEY, W.R. Jr (1977). Muscarinic, alpha-adrenergic and peptide regulate the same calcium influx sites in the parotid gland. *J. Physiol. (Lond.)*, **268**, 139-149.
- PUTNEY, W.R. Jr (1981). Recent hypotheses regarding the phosphatidylinositol effect. *Life Sci.*, **29**, 1183-1194.
- RADULOVACKI, M., WOJCIK, M.J., FORMAL, C. & MILETICH, R. (1980). Elimination of REM sleep rebound in rats by alpha-adrenoceptor blockers, phentolamine and phenoxybenzamine. *Pharmac. Biochem. Behav.*, **13**, 51-55.
- RADULOVACKI, M., WOJCIK, M.J., WALOVICH, R. & BRODIE, M. (1981). Phenoxybenzamine and bromocriptine attenuate need for REM sleep in rats. *Pharmac. Biochem. Behav.*, **14**, 371-375.
- RAMBOURG, A. & DROZ, B. (1980). Smooth endoplasmic reticulum and axonal transport. *J. Neurochem.*, **35**, 16-25.
- RAMM, P. (1979). The locus coeruleus, catecholamines, and REM sleep: a critical review. *Behav. Biol.*, **25**, 415-448.
- RICHARDS, J.G., LOREZ, H.P. & TRANZER, J.P. (1973). Indolealkylamine nerve terminals in cerebral ventricles: identification by electron microscopy and fluorescence histochemistry. *Brain Res.*, **57**, 277-288.
- RIMON, G., HANSKI, E., BRAUN, S. & LEVITZKI, A. (1978). Mode of coupling between hormone receptors and adenylate cyclase elucidated by modulation of membrane fluidity. *Nature*, **276**, 394-396.
- ROFFWARG, H.P., MUZIO, J.N. & DEMENT, W.C. (1966). Ontogenetic development of the human sleep-dream cycle. *Science*, **152**, 604-619.
- ROTHMAN, J.E., BURSZTYN-PETTIGREW, H. & FINE, R.E. (1980). Transport of the membrane glycoprotein of vesicular stomatitis virus to the cell surface in two stages by clathrin-coated vesicles. *J. cell. Biol.*, **86**, 162-171.
- ROTHMAN, J.E. & LODISH, H.F. (1977). Synchronized transmembrane insertion and glycosylation of a nascent membrane protein. *Nature (Lond.)*, **269**, 775-780.
- RYAN, G.B., BORYSENKO, J.Z. & KARNOVSKY, M.J. (1974). Factors affecting the redistribution of surface-bound concanavalin A on human polymorphonuclear leukocytes. *J. cell. Biol.*, **62**, 351-365.
- SAKAI, K. (1980). Some anatomical and physiological properties of ponto-mesencephalic tegmental neurons with special reference to the PGO waves and postural atonia during paradoxical sleep in the cat. In *The reticular formation revisited*, eds Hobson, J.A. & Brazier, M.A.B., pp. 427-447. New York: Raven Press.
- SALISBURY, J.L., CONDEELIS, J.S., MAIHLE, N.J. & SATIR, P. (1981). Calmodulin localization during capping and receptor-mediated endocytosis. *Nature*, **294**, 164-166.
- SALISBURY, J.L., CONDEELIS, J.S. & SATIR, P. (1980). Role of coated vesicles, microfilaments, and calmodulin in receptor-mediated endocytosis by cultured B Lymphoblastoid cells. *J. cell. Biol.*, **87**, 132-141.
- SALLANON, M. (1981). Mécanismes sérotoninergiques et non sérotoninergiques des états de sommeil. *Thèse No 1102*, présentée devant l'Université Claude-Bernard-LYON 1.
- SASTRE, J.-P., SAKAI, K. & JOUVET, M. (1981). Are the gigantocellular tegmental field neurons responsible for paradoxical sleep? *Brain Res.*, **229**, 147-161.
- SCHWARTZ, J.C., COSTENTIN, J., MARTRES, M.P., PROTAIS, P. & BAUDRY, M. (1978). Modulation of receptor mechanisms in the CNS: Hyper- and hypo-sensitivity to catecholamines. *Neuropharmacology*, **17**, 665-685.
- SHIMIZU, N. & IMAMOTO, K. (1970). Fine structure of the locus coeruleus in the rat. *Arch. Histol. jap.*, **31**, 229-246.
- SHINITZKY, M. & HENKART, P. (1979). Fluidity of cell membranes. Current concepts and trends. *Int. Rev. Cytol.*, **60**, 121-147.
- SHINITZKY, M. & RIVNAY, B. (1977). Degree of exposure of membrane proteins determined by fluorescence quenching. *Biochemistry*, **16**, 982-986.
- SIEVER, L.J., COHEN, R.M. & MURPHY, D.L. (1981). Anti-

- depressants and α_2 -adrenergic autoreceptor desensitization. *Ann. J. Psychiat.*, **138**, 681–682.
- SILBERMAN, E.K., VIVALDI, E., GARFIELD, J., MCCARLEY, R.W. & HOBSON, J.A. (1980). Carbachol triggering of desynchronized sleep phenomena: enhancement via small volume infusions. *Brain Res.*, **191**, 215–224.
- SINGER, S.J. (1974). The molecular organization of membranes. *Ann. Rev. Biochem.*, **43**, 805–833.
- SINGER, S.J. & NICOLSON, G.L. (1972). The fluid mosaic model of the structure of cell membranes. *Science*, **175**, 720–731.
- SITARAM, N., MOORE, A.M. & GILLIN, J.C. (1978). Experimental acceleration and slowing of REM sleep ultradian rhythm by cholinergic agonist and antagonist. *Nature*, **274**, 490–492.
- SLADEK, J.R. Jr & PARNAVELAS, J.G. (1975). Catecholamine-containing dendrites in primate brain. *Brain Res.*, **100**, 657–662.
- SMITH, C.B., GARCIA-SEVILLA, J.A. & HOLLINGSWORTH, P.J. (1981). α_2 -adrenoceptors in rat brain are decreased after long-term tricyclic antidepressant drug treatment. *Brain Res.*, **210**, 413–418.
- SMITH, A.P. & LOH, H.H. (1977). The organization of proteins in mouse synaptosomal plasma membrane. *Res. Comm. Chem. Path. Pharmac.*, **18**, 106–116.
- SMITH, A.P. & LOH, H.H. (1978). Mobile and immobile proteins of synaptosomal plasma membrane. *Biochemistry*, **17**, 4343–4347.
- SMITH, A.P. & LOH, H.H. (1979). The architecture of the nerve ending membrane. *Life Sci.*, **24**, 1–20.
- SPYRAKI, C. & FIBIGER, H.C. (1980). Functional evidence for subsensitivity of noradrenergic α_2 -receptors after chronic desipramine treatment. *Life Sci.*, **27**, 1863–1867.
- STARKE, K. & ALTMANN, K.P. (1973). Inhibition of adrenergic neurotransmission by clonidine: an action on pre-junctional α -receptors. *Neuropharmacology*, **12**, 339–347.
- STARKE, K. & MONTEL, H. (1973). Involvement of α -receptors in clonidine-induced inhibition of transmitter release from central monoamine neurones. *Neuropharmacology*, **12**, 1073–1080.
- STARKE, K., MONTEL, H., GAYK, W. & MERKER, R. (1974). Comparison of the effects of clonidine on pre- and postsynaptic adrenoceptors in the rabbit pulmonary artery. *Naunyn-Schmiedeberg's Arch. Pharmac.*, **285**, 133–150.
- STEIN, D., JOUVET, M. & PUJOL, J.-F. (1974). Effects of α -methyl-*p*-tryptosine upon cerebral amine metabolism and sleep states in the cat. *Brain Res.*, **72**, 360–365.
- STERIADE, M. & HOBSON, J.A. (1976). Neuronal activity during the sleep-waking cycle. In *Progress in Neurobiology*, eds Kerkut, G.A. & Phillips, J.W., Vol. 6, pp. 155–376. Oxford: Pergamon Press.
- STERN, W.C. & MORGANE, P.J. (1973). Effects of α -methyltyrosine on REM sleep and brain amine levels in the cat. *Biol. Psychiat.*, **6**, 301–306.
- STERN, W.C. & MORGANE, P.J. (1974). Theoretical view of REM sleep function: maintenance of catecholamine in the central nervous system. *Behav. Biol.*, **11**, 1–32.
- STRITTMATTER, W.J., HIRATA, F. & AXELROD, J. (1979). Phospholipid methylation unmasks cryptic β -adrenergic receptors in rat reticulocytes. *Science*, **204**, 1205–1207.
- SVENSSON, T.H., BUNNEY, B.S. & AGHAJANIAN, G.K. (1975). Inhibition of both noradrenergic and serotonergic neurons in brain by the α -adrenergic agonist clonidine. *Brain Res.*, **92**, 291–306.
- SWANSON, L.W. (1976). The locus coeruleus: A cytoarchitectonic, Golgi and immunohistochemical study in the albino rat. *Brain Res.*, **110**, 39–56.
- TANAKA, T. & STARKE, K. (1980). Antagonist/agonist-preferring α -adrenoceptors or α_1/α_2 -adrenoceptors? *Eur. J. Pharmac.*, **63**, 191–194.
- TAUBE, H.D., STARKE, K. & BOROWSKI, E. (1977). Presynaptic receptor systems on the noradrenergic neurones of rat brain. *Naunyn-Schmiedeberg's Arch. Pharmac.*, **299**, 123–141.
- TAYLOR, R.B., DUFFUS, W., RAFF, M. & PETRIS, S. (1971). Redistribution and pinocytosis of lymphocyte surface immunoglobulin molecules induced by anti-immunoglobulin antibody. *Nature New Biol.*, **233**, 225–229.
- TIMMERMANS, P.B.M.W.M. & VAN ZWEITEN, P.A. (1981). The postsynaptic α_2 -adrenoceptors. Minireview. *J. auton. Pharmac.*, **1**, 171–183.
- TISSOT, R. (1966). Essai méthodologique d'analyse du cycle du sommeil. *Encéphale*, **2**, 137–169.
- TITELER, M. & SEEMAN, P. (1980). Presynaptically acting catecholamines bind to α_2 -adrenoceptors labelled by ^3H -clonidine. *Eur. J. Pharmac.*, **67**, 187–192.
- TORDA, C. (1968). Effects of changes of norepinephrine content of sleep cycle in rat. *Brain Res.*, **10**, 221–235.
- TSUDA, M. & AKINO, T. (1981). Lipid-protein interaction in the photolysis of octopus rhodopsin. *Biochim. biophys. Acta*, **643**, 63–75.
- VENTER, J.C. (1979). High efficiency coupling between β -adrenergic receptors and cardiac contractility: Direct evidence for 'spare' β -adrenergic receptors. *Mol. Pharmac.*, **16**, 429–440.
- WANG, R.Y. & AGHAJANIAN, G.K. (1978). Collateral inhibition of serotonergic neurones in the rat dorsal raphe nucleus: Pharmacological evidence. *Neuropharmacology*, **17**, 819–825.
- WATABE, K. & SATOH, T. (1980). Cross-correlation analysis of neuronal activities in bilateral nuclei of locus coeruleus. *Brain Res.*, **199**, 231–234.
- WEISS, C.B. & LEVIN, R.M. (1978). Mechanism for selectively inhibiting the activation of cyclic nucleotide phosphodiesterase and adenylate cyclase. *Adv. cyclic nucleotide Res.*, **9**, 285–303.
- WEITZMAN, E.D., MCGREGOR, P., MOORE, C. & JACOBY, J. (1969). The effect of α -methyl-paratyrosine on sleep patterns of the monkey. *Life Sci.*, **8**, 751–757.
- WICKNER, W. (1980). Assembly of proteins into membranes. *Science*, **210**, 861–868.
- WIRZ-JUSTICE, A., KRAUCHII, K., LICHTSTEINER, M. & FEER, H. (1978). Is it possible to modify serotonin receptor sensitivity? *Life Sci.*, **23**, 1249–1254.
- WYATT, R.J. (1972). The serotonin-catecholamine-dream bicycle: A clinical study. *Biol. Psychiat.*, **5**, 33–64.
- WYATT, R.J., CHASE, T.N., KUPFER, D.J., SCOTT, J., SNYDER, F., SJOERDSMA, A. & ENGELMAN, K. (1971). Brain catecholamines and human sleep. *Nature*, **233**, 63–65.
- YAHARA, I. & EDELMAN, G.M. (1973). Modulation of lymphocyte receptor redistribution of concanavalin A, anti-mitotic agents and alterations of pH. *Nature*, **246**, 152–155.

Biochemische Pharmakologie des paradoxen Schlafes

J.-M. GAILLARD

1 Die Rolle der noradrenergen Zellen bei der Regulierung des paradoxen Schlafes ist immer noch strittig; Versuchsergebnisse führten zu widersprüchlichen Interpretationen gegeben.

2 Frühere Untersuchungen konzentrierten sich auf die chemische Neurotransmission. Der Ablauf der Informationsübertragung zwischen den Zellen schliesst jedoch eine Reihe von weiteren Faktoren ein und die Zelloberfläche ist ein wichtiger Ort zur Transduktion von Informationen in Veränderungen der postsynaptischen Zellaktivitäten.

3 α -adrenerge Rezeptoren spielen vermutlich bei der Kontrolle des Wachseins und des paradoxen Schlafes eine wesentliche Rolle. Experimentelle Befunde deuten darauf hin, dass eine physiologische Modulation der Rezeptorsensibilität, eventuell durch spezifische Neuromodulatoren, ein Schlüssel-

mechanismus bei der synaptischen Transmission darstellen könnte.

4 Bei der Untersuchung der bei der Regulierung des paradoxen Schlafes mitwirkenden Mechanismen, führten Läsionen des Locus coeruleus zu widersprüchlichen Ergebnissen. Die kollaterale Hemmung—vermutlich über α_2 -adrenerge Rezeptoren—scheint ein stark wirksamer Mechanismus zu sein. Der genaue zeitliche Zusammenhang zwischen der noradrenergen Zellaktivierung und der Entstehung von paradoxem Schlaf ist nicht gesichert; 5-HT scheint jedoch dabei eine Rolle zu spielen. Unterschiede zwischen paradoxem Schlaf und Wachsein können mit einer physiologischen Modulation der α_2 -adrenergen Rezeptorsensibilität zusammenhängen.

Pharmacologie biochimique du sommeil paradoxal

J.-M. GAILLARD

1 Le rôle joué par les cellules noradrénergiques dans la régulation du sommeil paradoxal est encore controversé, et les résultats expérimentaux ont donné lieu à des interprétations contradictoires.

2 Les premières recherches ont porté principalement sur la neurotransmission chimique. Cependant, le processus de transmission des informations entre les cellules implique de nombreux autres facteurs, et la surface cellulaire est un site important de la transduction de messages dans des modifications de l'activité des cellules postsynaptiques.

3 On pense que les récepteurs α -adrénergiques jouent un rôle important dans la commande de l'état vigile et du sommeil paradoxal. Il semble, d'après les résultats d'expériences, que la modulation physiologique de la sensibilité des récepteurs, peut-être par

des neuro-modulateurs spécifiques, soit un mécanisme clef de la transmission synaptique.

4 L'investigation des mécanismes impliqués dans la régulation du sommeil paradoxal n'a pas donné, en ce qui concerne les lésions du locus coeruleus, de résultats concluants. Il semble que l'inhibition collatérale, probablement atteinte par l'intermédiaire des récepteurs α_2 -adrénergiques, constitue un mécanisme puissant. La relation temporelle exacte entre l'activation de la cellule noradrénergique et la production du sommeil paradoxal n'a pu encore être établie, mais il semble que la sérotonine joue un rôle dans ce processus. Les différences entre le sommeil paradoxal et l'état de veille pourraient être liées à une modulation physiologique de la sensibilité des récepteurs α_2 -adrénergiques.

Farmacología bioquímica del sueño paradójal

J.-M. GAILLARD

1 El papel de las células noradrenérgicas en la regulación del sueño paradójal sigue siendo todavía objeto de controversia; datos experimentales han suscitado interpretaciones contradictorias.

2 Las primeras investigaciones se centraron principalmente en la neurotransmisión química. No obstante, el proceso de transmisión de información entre células afecta a muchos otros factores y la superficie celular constituye un importante sitio para

la transducción de mensajes en modificaciones de la actividad de células postsinápticas.

3 Se cree que los α -adrenoceptores juegan un importante papel en el control de la vigilia y del sueño paradójal. La evidencia experimental sugiere que la modulación fisiológica de la sensibilidad de los receptores, posiblemente por neuromoduladores específicos, puede constituir un mecanismo clave en la transmisión sináptica.

4 Al investigar los mecanismos involucrados en la regulación del sueño paradójico, lesiones del locus coeruleus han dado lugar a resultados equívocos. Parece ser que la inhibición colateral, mediada probablemente por α_2 -adrenoreceptores, representa un mecanismo potente. No se ha establecido la relación

temporal exacta entre activación celular noradrenérgica y producción de sueño paradójico, pero parece intervenir la serotonina. Diferencias entre sueño paradójico y vigilia pueden guardar relación con una modulación fisiológica de la sensibilidad de los α_2 -adrenoreceptores.