may be due to more rapid excretion and metabolism of proxiphylline since it is a more water soluble compound.

It is difficult to assign a clinical significance to these results since figures are not available for the relative potencies of proxiphylline and theophylline as bronchodilators. Considering the three theophylline preparations, if the degree of broncho-dilatation is related to the concentration of theophylline, then it would seem that the sustained release preparation should produce a steadier, more prolonged effect, especially during the night.

B.A. SIMS

Waveney Hospital, Ballymena, Northern Ireland

J.G. KELLY & R.G. SHANKS

Department of Therapeutics & Pharmacology, Queen's

University of Belfast, Institute of Clinical Sciences, Belfast BT12 6BJ, Northern Ireland

Received April 21, 1975

References

- JENNE, J.W., WYZE, M.S.E., READ, F.S. & MACDONALD, F.M. (1972). Pharmacokinetics of theophylline. *Clin. Pharmac. Ther.*, **13**, 329-359.
- MITENKO, P.A. & OGILVIE, R.I. (1973). Rational intravenous doses of theophylline. *New Eng. J. Med.*, **289**, 600-603.
- SCHACK, J.A. & WAXLER, S.H. (1949). An ultraviolet • spectrophotometric method for the determination of theophylline and theobromine in blood and tissues. *J. Pharmac. exp. Ther.*, **97**, 283-291.
- TURNER-WARWICK, M. (1957). Study of theophylline plasma levels after oral administration of new theophylline compounds. *Br. med. J.*, **2**, 67-69.
- WAXLER, S.H. & SCHACK, J.A. (1950). Administration of aminophylline. *J. Am. med. Ass.*, **142**, 736-739.

THE INFLUENCE OF pH ON THE BUCCAL ABSORPTION AND PLASMA AND RENAL ELIMINATION OF ACEBUTOLOL

Acebutolol, 1-(2-acetyl-4-*n*-butyramidophenoxy)-2-hydroxy-3-isopropyl-aminopropane, is a new β -adrenoceptor blocking drug effective in man when given orally (Lewis, Bakst, Kitchiner & Gotsman, 1973; Cuthbert & Collins, 1975). Although it has a similar structure to practolol, acebutolol has been shown to be less cardio-selective in man (Kumana, Kaye, Leighton, Turner & Hamer, 1975). Before detailed investigation of its pharmacokinetics could be undertaken it was necessary to establish whether elimination of acebutolol is dependent on urine pH. As buccal absorption is a useful model of renal tubular reabsorption, a preliminary study was made of the influence of pH on the buccal absorption of acebutolol in four subjects. The method used was that described by Beckett & Triggs (1967) and the amount of drug remaining after the buccal rinse was determined by extraction of the diluted saliva with ethyl acetate, and after back extraction into acid, measuring the amount of UV absorption at 320 nm. The mean ($\pm$ s.e. mean) results of the four subjects are shown in Figure 1, and clearly demonstrate that the buccal absorption of acebutolol is not pH dependent.

In order to determine the influence of urine pH on the plasma and renal elimination of acebutolol,

Figure 1 Influence of pH on the buccal absorption of acebutolol. The results are shown as mean $\pm$ s.e. mean.

four healthy male volunteers (three of whom had taken part in the previous study) each took 3 $\times$ 100 mg capsules of acebutolol (Sectral) on two separate occasions. On one occasion urine pH was maintained at around 5 by ammonium chloride ingestion, and on the other occasion at about 8 by sodium bicarbonate ingestion (Beckett & Rowland, 1965). Urine was collected for 24 h after each dose, and in two subjects (1 & 3) venous

Figure 2 Influence of urine pH on the renal excretion of acebutolol □ acidic urine; ▨ alkaline urine.

blood samples were obtained at 2, 3, 4, 5, 6, 8, 10 and 12 h after dosing. The acebutolol in urine and plasma was determined spectrophotometrically by a method based on that described by Cuthbert & Collins (1975). The 24 h urinary recovery of acebutolol as measured for the four subjects on each occasion is shown in Figure 2. In any individual urinary recovery was similar whether urine pH was acidic or alkaline, indicating that the renal excretion of acebutolol is independent of urine pH. As the assay method used also measures 1-(2-acetyl-4-acetamidophenoxy)-2-hydroxy-3-isopropylaminopropane, a known metabolite of acebutolol in man (George & Collins, personal communication), the urine samples were further examined by t.l.c. (using chloroform:methanol:glacial acetic acid 7:3:1 by volume as the solvent) to see whether the overall results were due to a variation in the amount of this metabolite excreted. It was found that the relative amounts of parent drug and metabolite did not appear to vary with different conditions of urine pH.

The plasma elimination of acebutolol was studied in two of the subjects, and the results for subject 1 are shown in Figure 3. Following acebutolol (300 mg), the plasma level of the drug quickly declined between 4 and 12 h, having a half-life of 2.3 h when urine pH was around 5 and 2.7 h when urine pH was about 8. The results with subject 3 were similar, the respective half-lives being 2.7 and 3.3 hours. Such small changes suggest that the plasma elimination of acebutolol is virtually independent of urine pH.

The present results are consistent with those obtained previously with other drugs. Acebutolol has a pKa of 9.40 and water solubility of 0.31% (Collins, personal communication) and these are similar to the corresponding values of 9.50 and

Figure 3 Influence of urine pH on the plasma elimination of acebutolol ○ acidic urine, $T_{1/2}$ 2.3 h; ● alkaline urine, $T_{1/2}$ 2.7 h.

0.25% for practolol, the buccal absorption and renal excretion of which are also independent of pH (Kaye, Robinson & Turner, 1973). Thus in subsequent studies with acebutolol urine pH need not be controlled.

We thank Professor P. Turner and Dr R.F. Collins for useful discussion, Dr C.R. Kumana for his co-operation the British Heart Foundation for financial support, and May & Baker Limited for supplying acebutolol (Sectral, M&B 17,803), its *N*-dealkylated derivative (M&B 17,127) and the *N*-acetyl derivative of the latter (M&B 16,942).

C.M. KAYE & A.D. LONG

Department of Clinical Pharmacology, St. Bartholomew's Hospital, London EC1A 7BE

Received July 17, 1975

References

- BECKETT, A.H. & ROWLAND, M. (1965). Urinary excretion kinetics of amphetamine in man. *J. Pharm. Pharmac.*, **17**, 628-639.
- BECKETT, A.H. & TRIGGS, E.J. (1967). Buccal absorption of basic drugs and its application as an *in vivo* model of passive drug transfer through lipid membranes. *J. Pharm. Pharmac.*, **19**, 31S-41S.
- CUTHBERT, M.F. & COLLINS, R.F. (1975). Plasma

levels and β -adrenoceptor blockade with acebutolol, practolol and propranolol in man. *Br. J. clin. Pharmac.*, **2**, 49-56.

KAYE, C.M., ROBINSON, D.G. & TURNER, P. (1973).

The influence of urine pH on the renal excretion of practolol and propranolol. *Br. J. Pharmac.*, **49**, 155P.

KUMANA, C.R., KAYE, C.M., LEIGHTON, M.,

TURNER, P. & HAMER, J. (1975). Cardiac and pulmonary effects of acebutolol in man. *Lancet*, **ii**, 89-93.

LEWIS, B.S., BAKST, A., KITCHINER, D. & GOTSMAN, M.S. (1973). Effectiveness of Sactal (M&B 17,803A) as a beta-blocking agent in man. *S. Afr. med. J.*, **47**, 1181-1184.

PRELIMINARY OBSERVATIONS ON THE ELIMINATION OF ACEBUTOLOL IN SEVERE CHRONIC RENAL FAILURE

Acebutolol is a new β -adrenoceptor blocking agent effective in man (Lewis, Bakst, Kitchiner & Gotsman, 1973a; Cuthbert & Collins, 1975); Kumana, Kaye, Leighton, Turner & Hamer, 1975), which may be useful in the treatment of hypertension, angina pectoris (Lewis, Bakst, Kitchiner & Gotsman, 1973b; Biron & Tremblay, 1975), and cardiac arrhythmias (Biron, Proulx, Lapointe, Nadeau & Tremblay, 1975). The use of β -adrenoceptor blocking drugs in the control of hypertension is now well established, and it is therefore likely that this drug may be prescribed for patients who have hypertension associated with severe renal impairment. This letter reports some preliminary findings obtained in the course of investigations to establish the effect if any of chronic renal failure on the elimination of acebutolol.

Six healthy male volunteers (mean $\pm$ s.d. age 30 ± 6 years, mean $\pm$ s.d. weight 73 ± 9 kg,

mean $\pm$ s.d. creatinine clearance 103 ± 7 ml/min) and six patients (five males and one female, mean $\pm$ s.d. age 42 ± 13 years, mean $\pm$ s.d. weight 67 ± 10 kg) with severe chronic renal failure (mean $\pm$ s.d. creatinine clearance 4 ± 2 ml/min) took part in the study. Other drug treatment in these patients continued unchanged and did not interfere with the assay. Each subject received a single oral dose of acebutolol (3×100 mg capsules Sactal) after which venous blood samples were obtained at 2, 4, 6, 8 and 10 h in the healthy volunteers and at 2, 4, 6, 8, 10 and 24 h in the patients (it being previously known that after such a dose the plasma level by 24 h in normal subjects is below the detection limit of the method). Urine was collected from each subject for 24 h following the dose. Plasma and urinary acebutolol were measured by a spectrophotometric method based on that described by Cuthbert & Collins (1975) which also measures an active metabolite of

Figure 1 Mean ($\pm$ s.d.) plasma levels (parent drug plus metabolite, see text) after the oral administration of acebutolol (300 mg) to six normal subjects ($\bullet$) and six patients ($\blacksquare$) with severe chronic renal failure.