- PRIME, F.J., KAMBUROFF, P.L. & GUNTNER, W. (1971). Comparison of the bronchodilatory action of two drugs. *Int. J. clin. Pharmac. Ther. Tox.*, 5, 293-295.
- SHENFIELD, GILLIAN M. & PATERSON, J.W. (1973). Clinical assessment of bronchodilator drugs delivered by aerosol. *Thorax*, 28, 124-128.
- SVEDMYR, N., MALMBERG, R. & THIRINGER, G. (1972). The effect of a new adrenergic β_2 -receptor-stimulating agent (rimeterol, R798) in patients with chronic obstructive lung disease. *Scand. J. resp. Dis.*, 53, 302-313.
- WARRELL, D.A., ROBERTSON, D.G., NEWTON HOWES, J., CONOLLY, M.E., PATERSON, J.W., BEILIN, L.J. & DOLLERY, C.T. (1970). Comparison of cardiorespiratory effects of isoprenaline and salbutamol in patients with bronchial asthma. *Br. med. J.*, 1, 65-70.

STUDIES OF THE EFFECT OF THYROID DYSFUNCTION ON THE ELIMINATION OF β -ADRENOCEPTOR BLOCKING DRUGS

It is now clearly recognised that drug metabolism may be altered by disease. Hence renal disease may effect drug clearance through its influence on elimination or on plasma binding. Thyroid dysfunction has also been shown to alter the metabolism of several drugs. The elimination half-life of antipyrine is shortened in hyperthyroid subjects and prolonged in hypothyroid subjects (Crooks, Headley, McNee & Stevenson, 1973; Vesell & Passananti, 1973; Eichelbaum, Bodem, Gugler, Schneider-Deters & Dengler, 1974): the plasma concentrations and serum half-lives of digoxin have been shown to be lower in hyperthyroid and higher in hypothyroid patients apparently related to changes in creatinine clearance (Croxon & Ibbertson, 1975). However, recently the apparent half-life and other pharmacokinetic parameters of propylthiouracil in ten hyperthyroid subjects were not found significantly different from corresponding values in a previous study in euthyroid subjects (Kampmann & Skovsted, 1974; Kampmann & Skovsted, 1975).

In the past few years, the use of β -adrenoceptor blocking drugs, particularly propranolol, has become established in various aspects of the management of hyperthyroidism (Turner, 1974; McDevitt, 1976). No information is currently available on the influence of thyroid disease on the

metabolism of β -adrenoceptor blocking drugs. We have, therefore, studied the elimination of propranolol and practolol, two drugs with different modes of clearance (Shand, 1974), in patients both with hyperthyroidism and hypothyroidism.

Eight patients with hyperthyroidism and four with hypothyroidism have been studied. Their age, sex and biochemical findings are shown in Table 1. Patients with heart disease, bronchial asthma and hepatic or renal disease were excluded. After an overnight fast, each patient was given propranolol (100 mg) orally at 09.00 h on Day 1 and practolol (400 mg) orally at 09.00 h on Day 2. The order of drug administration was not randomised due to the longer half-life of practolol and the need to have patients in hospital for as brief a time as possible. Blood samples were withdrawn through an indwelling catheter in a forearm vein. With propranolol, blood samples were withdrawn before and at 1, 2, 3, 4, 6, 9 and 12 h after the drug was given: samples were heparinised. Blood samples for practolol were withdrawn before and at 1, 2, 3, 5, 7, 9, 12, 24, 36 and 48 h after drug administration. Blood samples for practolol were oxalated. Plasma propranolol concentrations were subsequently measured by the standard spectrofluorometric method (Shand, Nuckolls & Oates, 1970) and blood practolol concentrations by a

Table 1 Mean $\pm$ s.e. mean observations on eight patients with hyperthyroidism and four patients with hypothyroidism

Thyroid status	Sex	Age (years)	T_3^* (% uptake)	T_4^* (mmol/l)	F.T.I.†
Hyperthyroid	7F, 1M	50.8 $\pm$ 4.8	85.9 $\pm$ 4.5	234.5 $\pm$ 18.3	280.1 $\pm$ 31.1
Hypothyroid	4F	48.0 $\pm$ 6.5	127.4 $\pm$ 1.1	21.1 $\pm$ 8.0	17.5 $\pm$ 5.8

* Normal range, 95-122% uptake; + Normal range, 54.1-144.1 mmol/l; † Normal range, 51.5-141.6

Figure 1 Mean plasma propranolol concentrations $\pm$ s.e. mean after the administration of propranolol (100 mg orally) to seven hyperthyroid patients ($\bullet$) and four hypothyroid patients ($\circ$). * = $P < 0.05$ for comparison between plasma concentrations in the two groups. + = mean of only 2 patients.

high-speed liquid chromatography technique (Kelly & Leahey, unpublished observations). Drug half-lives were estimated from the terminal portion of the plasma or blood curves by least squares regression analysis. Data from 3 to 12 h was used from the propranolol curves and from 7 to 48 h after practolol. The area under the plasma concentration/time curve (AUC) was estimated by the trapezoid rule after extrapolation to infinity. Statistical analysis was carried out using Student's unpaired *t*-test. Results are expressed as the mean $\pm$ s.e. mean.

Figure 2 Mean blood practolol concentrations $\pm$ s.e. mean after the administration of practolol (400 mg orally) to eight hyperthyroid ($\bullet$) and four hypothyroid patients ($\circ$). * = $P < 0.05$ for comparison between blood concentrations in two groups.

The results are shown in Figures 1 and 2 and in Table 2. After propranolol (100 mg) orally, plasma elimination curves were obtained in only seven hyperthyroid and in four hypothyroid patients. Propranolol appeared to be absorbed much more rapidly in hyperthyroid than hypothyroid subjects, and the plasma propranolol concentration after 1 h was significantly higher in the former group than that obtained in the latter

Table 2 Mean $\pm$ s.e. mean results for kinetic parameters in eight hyperthyroid and four hypothyroid patients after the administration of propranolol (100 mg) and practolol (400 mg) orally

Drug	Parameter	Hyperthyroid* (n = 8)	Hypothyroid (n = 4)	P (unpaired t-test)
Propranolol (100 mg orally)	$T_{1/2}$ (h)	2.9 $\pm$ 0.5	3.4 $\pm$ 0.5	NS
	AUC _{0-∞} (ng ml ⁻¹ h)	700.3 $\pm$ 193.1	469.0 $\pm$ 119.7	NS
	Peak plasma concentration (ng/ml)	123.4 $\pm$ 36.7	75.0 $\pm$ 14.4	NS
	Time to observed peak (h)	1.3 $\pm$ 0.2	3.5 $\pm$ 0.3	< 0.001
Practolol (400 mg orally)	$T_{1/2}$ (h)	9.8 $\pm$ 0.5	16.0 $\pm$ 3.2	< 0.02
	AUC _{0-∞} (µg ml ⁻¹ h)	53.3 $\pm$ 6.0	97.3 $\pm$ 17.1	< 0.02
	Peak blood concentration (µg/ml)	5.7 $\pm$ 1.1	7.2 $\pm$ 1.1	NS
	Time to observed peak (h)	1.8 $\pm$ 0.3	2.0 $\pm$ 0.4	NS

* Results for propranolol in hyperthyroid patients are the mean of only 7.

($P < 0.05$). In addition, the mean time from administration of drug to the observed peak plasma concentration was significantly shorter in the hyperthyroid compared to the hypothyroid patients ($P < 0.001$). However, the elimination half-lives, the $AUC_{0 \rightarrow \infty}$ and the peak plasma concentrations did not differ significantly between the two groups. Neither could any difference be shown between the slopes of the two elimination curves by analysis of covariance.

Following practolol (400 mg) orally, the pattern was quite different. One hour after administration the blood practolol concentrations were approximately the same in both groups of patients, but after 2 h mean practolol concentrations were higher in the hypothyroid patients ($P < 0.01$). However, significant differences in the peak blood concentrations and the time to observed peak blood practolol levels were not observed. The elimination half-life was significantly longer and the $AUC_{0 \rightarrow \infty}$ was significantly greater in hypothyroid patients than in those with hyperthyroidism. Comparison of the slopes of the elimination curves by analysis of covariance showed a significant difference between the two ($P < 0.001$).

Within the limitations of this study, thyroid status did not appear to affect the metabolism of propranolol materially. Apart from a slower rate of absorption in patients with hypothyroidism, evidenced by early differences in plasma propranolol concentrations and the time taken for peak concentrations to be achieved, no significant differences were observed between hyperthyroid and hypothyroid patients. If $AUC_{0 \rightarrow \infty}$ after oral administration be taken as a crude estimate of bioavailability, it did not differ significantly between the two groups. The elimination half-lives obtained for propranolol appear to be similar to those previously reported by Shand *et al.* (1970) in euthyroid subjects. Sevenfold variations in plasma propranolol concentrations between individuals were found in this study, again similar to those reported for normal subjects (Shand *et al.*, 1970).

In contrast, the elimination of practolol did appear to be dependent on thyroid function. Patients with hypothyroidism had a mean half-life of 16.0 ± 3.2 h compared with 9.8 ± 0.5 h in hyperthyroid subjects. In normal subjects, the elimination half-life of practolol is 10–11 h (Carruthers, Kelly, McDevitt & Shanks, 1974). It would appear, therefore, that elimination is prolonged in hypothyroid patients. As the clearance of practolol is principally renal (Bodem & Chidsey, 1973), this may be a further manifestation of alteration in creatinine clearance in patients with thyroid dysfunction which has

recently been implicated in disturbances of digoxin metabolism with these diseases (Croxon & Ibbertson, 1975). Bioavailability also appeared to be less in patients with hyperthyroidism than in those with hypothyroidism. As practolol is completely absorbed in normal subjects (Bodem & Chidsey, 1973; Carruthers *et al.*, 1974), this may indicate incomplete absorption in patients with hyperthyroidism perhaps related to rapid transit time. However, we have not ruled out the possibility that alterations in the apparent volume of distribution may occur in thyroid disease.

Thus the effect of thyroid dysfunction on β -adrenoceptor blocking drugs may depend on the metabolism of the drug. More detailed studies with propranolol are underway to try and clarify this point. In the meantime, considerations of currently available information might suggest the following hypothesis for the metabolism of drugs in thyroid disease—drugs dependent on renal clearance, like digoxin and practolol, will be affected by thyroid disease, where hyperthyroidism increases creatinine clearance and hypothyroidism decreases it; drugs dependent on hepatic clearance will be affected in proportion to their hepatic extraction ratios—the metabolism of drugs with a high extraction ratio, such as propranolol, is more dependent on alterations in blood flow than in enzyme activity, whilst the metabolism of drugs with a low extraction ratio, e.g. antipyrine, is affected by alterations in intrinsic (hepatic) metabolic clearance (Wilkinson & Shand, 1975). If thyroid dysfunction affects intrinsic metabolic clearance as well as creatinine clearance, it could explain why the metabolism of digoxin and practolol, as well as that of antipyrine, are affected on the one hand, whereas propranolol metabolism is little altered.

We would like to thank Mr W. Leahey for technical assistance.

Requests for reprints should be addressed to Dr D.G. McDevitt, Whitla Medical Building, 97 Lisburn Road, Belfast BT9 7BL.

J.M. BELL, C.J. RUSSELL & J.K. NELSON

The Ulster Hospital, Dundonald, Belfast, Northern Ireland

J.G. KELLY & D.G. McDEVITT

Department of Therapeutics and Pharmacology, The Queen's University of Belfast, Belfast, Northern Ireland

Received June 16, 1976

References

- BODEM, G. & CHIDSEY, C.A. (1973). Pharmacokinetic studies of practolol, a beta adrenergic antagonist, in man. *Clin. Pharmac. Ther.*, **14**, 26-29.
- CARRUTHERS, S.G., KELLY, J.G., McDEVITT, D.G. & SHANKS, R.G. (1974). Blood levels of practolol after oral and parenteral administration and their relationship to exercise heart rate. *Clin. Pharmac. Ther.*, **15**, 497-509.
- CROOKS, J., HEADLEY, A.J., McNEE, C. & STEVENSON, I.H. (1973). Changes in drug metabolising ability in thyroid disease. *Br. J. Pharmac.*, **49**, 156-157P.
- CROXSON, M.S. & IBBERTSON, H.K. (1975). Serum digoxin in patients with thyroid disease. *Br. med. J.*, **3**, 566-568.
- EICHELBAUM, M., BODEM, G., GUGLER, R., SCHNEIDER-DETERS, C. & DENGLE, H.J. (1974). Influence of thyroid status on plasma half-life of antipyrine in man. *New Engl. J. Med.*, **290**, 1040-1042.
- KAMPMANN, J. & SKOVSTED, L. (1974). The pharmacokinetics of propylthiouracil. *Acta pharmac. tox.*, **35**, 361-369.
- KAMPMANN, J. & SKOVSTED, L. (1975). The kinetics of propylthiouracil in hyperthyroidism. *Acta pharmac. tox.*, **37**, 201-210.
- McDEVITT, D.G. (1976). Propranolol in the treatment of thyrotoxicosis: a review. *Postgrad. med. J.*, **52** (suppl. 4), 156-160.
- SHAND, D.G. (1974). Pharmacokinetic properties of the β -adrenergic receptor blocking drugs. *Drugs*, **7**, 39-47.
- SHAND, D.G., NUCKOLLS, E.M. & OATES, J.A. (1970). Plasma propranolol levels in adults, with observations in four children. *Clin. Pharmac. Ther.*, **11**, 112-120.
- TURNER, P. (1974). β -adrenergic receptor blocking drugs in hyperthyroidism. *Drugs*, **7**, 48-54.
- VESELL, E.S. & PASSANANTI, G.T. (1973). Inhibition of drug metabolism in man. *Drug Metab. Disposition*, **1**, 402-410.
- WILKINSON, G.R. & SHAND, D.G. (1975). A physiological approach to hepatic drug clearance. *Clin. Pharmac. Ther.*, **18**, 377-390.

ALTERED PLASMA ALBUMIN IN THE NEWBORN INFANT

Qualitative differences in the plasma protein binding of certain drugs in newborn infants have been widely reported (Ganshorn & Kurz, 1968; Chignell, Vesell, Starkweather & Berlin, 1971; Ehrnebo, Agurell, Jalling, Yaffe & Boréus, 1971; Pruitt & Dayton, 1971; Rane, Lunde, Jalling, Yaffe & Sjöqvist, 1971; Krasner, Giacoia & Yaffe, 1973; Gorodischer, Krasner & Yaffe, 1974; Windorfer, Kuenzer & Urbanek, 1974). The presence of competing ligands such as bilirubin (Chignell *et al.*, 1971; Rane *et al.*, 1971), free fatty acids and steroids (Krasner *et al.*, 1973), in addition to hypoalbuminaemia, has been suggested as a contributory factor to the observed reduction in binding in neonates, and in a previous study (Wallace, 1976) this possibility was examined. It has also been postulated that altered plasma proteins might be present in neonates (Krasner *et al.* 1973) and that these might be involved in reduced binding. This study investigates the properties of albumin isolated from neonatal plasma.

Blood from neonates and adults was obtained as previously described (Wallace, 1976). Samples of albumin were prepared from adult and neonatal plasma using an affinity chromatography technique based on the methods of Böhme, Kopperschlager, Schultz & Hofmann (1972) and Travis & Pannell (1973). Phosphate buffer (0.05 M) with 2 M NaCl, pH 7.4, was used to elute the albumin

in place of 6 M urea as described by Travis & Pannell (1973).

Samples (100 μ g) of albumin were freeze-dried in vacuum hydrolysis tubes (Pierce) and 1 ml of constant-boiling HCl added to each tube. Samples were hydrolyzed under vacuum at 110°C for 24 h, freeze-dried once more, and analyzed on a Technicon amino acid analyzer.

Immunodiffusion studies of adult and neonatal albumin were carried out using agarose gel plates with trypan blue indicator (Miles). Rabbit anti-serum (20 μ l) to human serum albumin (Miles) was placed in the central well and allowed to diffuse for 30 min before 10 μ l of adult and neonatal albumin solutions of equal concentration were placed alternately into the six surrounding wells. The plate was incubated at 22°C for 24 h and examined for precipitin lines.

Isoelectric focussing of adult and neonatal albumin was carried out using the method detailed in Bio-Rad Bulletin 1030 (1975). The gels were stained in a solution of ethanol, water, acetic acid (50 : 45 : 5) containing 0.2% bromophenol blue. Destaining was carried out in the Canalco Electrophoretic Destainer, loaded with 10% isopropanol and 10% acetic acid, and the gels were then scanned using a Gilford spectrophotometer with Model 2520 gel scanning attachment.

Crystalline HSA was used as a reference stan-