CLINICAL STUDY OF CLOBAZAM IN ALCOHOL DETOXIFICATION

C. MÉRY

Laboratoires Diamant S.A., Direction Médicale, Cédex No. 3, 92080 Paris-la-Défense, France

J.-C. MARTIN

Médecin des Hôpitaux Psychiatriques, Hôpital Beauregard, 18000 Bourges, France

CLOBAZAM has been compared in a double-blind study with diazepam and placebo in the treatment of anxiety symptoms in chronic alcoholic patients undergoing withdrawal therapy (Martin, 1975).

Patients of either sex admitted to hospital for an alcohol detoxification programme were selected for the trial. On admission patients were randomly allocated to treatment with clobazam 15 mg daily, clobazam 30 mg daily, diazepam 15 mg daily or placebo, in divided doses, which was continued for 12 days. Clinical assessments were made on days 0, 4, 8 and 12, using the Hamilton Anxiety Scale (HAS). A five-point global assessment was completed by both the investigator and the patient at the completion of the

treatment period.

Forty patients entered the trial. The reduction in HAS mean scores revealed a significant difference in efficacy between the three active drug groups and the placebo group ($P < 0.05$) by day 8, though there was no significant difference between the active treatments.

The global assessment demonstrated a highly significant difference between both clobazam groups and placebo ($P < 0.001$); a significant difference in favour of clobazam, between clobazam 30 mg daily and diazepam 15 mg daily; and no significant difference in activity between clobazam 15 mg daily and diazepam 15 mg daily.

All treatments were well tolerated by the patients.

Reference

MARTIN, J.-C. (1975). Essai contrôlé d'un nouveau tranquillisant (le clobazam) chez des alcooliques chroniques en phase de sevrage. *J. Pharmac. Clin.* II, 21–27.