References

- DOLL, R. (1971). Unwanted effects of drugs. *Br. med. Bull.*, **27**, 25–31.
- FEINSTEIN, A.R. (1974). Clinical Biostatistics XXVIII: The biostatistical problems of pharmaceutical surveillance. *Clin. Pharmac. Ther.*, **16**, 110–123.
- GREENBLATT, D.J. & KOCH-WESER, J. (1973). Adverse reactions to β -adrenergic receptor blocking drugs. *Am. Heart J.*, **86**, 478–484.
- GREENBLATT, D.J. & KOCH-WESER, J. (1974). Adverse reactions to β -adrenergic receptor blocking drugs. *Drugs*, **7**, 118–129.
- KOCH-WESER, J. (1979). Detection of drug-induced disease.

- Br. J. clin. Pharmac.*, **8**, 83–84.
- LAWSON, D.H. (1979). Detection of drug-induced disease. *Br. J. clin. Pharmac.*, **7**, 13–18.
- PFEIFER, H.J., GREENBLATT, D.J. & KOCH-WESER, J. (1976a). Clinical use and toxicity of intravenous lidocaine. *Am. Heart J.*, **92**, 168–173.
- PFEIFER, H.J., GREENBLATT, D.J. & KOCH-WESER, J. (1976b). Clinical toxicity of reserpine in hospitalised patients. *Am. J. med. Sci.*, **271**, 269–276.
- PFEIFER, H.J., GREENBLATT, D.J. & KOCH-WESER, J. (1977). Adverse reactions to practolol in hospitalised patients. *Eur. J. clin. Pharmac.*, **12**, 167–170.

CHLORDIAZEPOXIDE IN ACUTE ALCOHOLISM

In a recent article Whiting, Lawrence, Skellern & Meier (1979) claimed that in patients acutely intoxicated with alcohol the elimination half-life of chlordiazepoxide was prolonged and its clearance from plasma was reduced. Unfortunately, the design of their study renders the data difficult to interpret. The effect of variable age on chlordiazepoxide disposition was not controlled in the paper of Whiting *et al.* (1979). The mean age of their control subjects was 31.2 years while for the alcoholic subjects it was 51.6 years. Recently, both Roberts, Wilkinson, Branch & Schenker (1978) and Shader, Greenblatt, Harmatz, Franke & Koch-Weser (1977) have reported a significant age-dependent alteration in chlordiazepoxide disposition. From the data of Roberts *et al.* (1978) the estimated half-life in normal

subjects at 31 years would be 5 h and at 52 years it would be 22 h; clearance of chlordiazepoxide would fall from 30 to 19 ml/min, respectively. These figures are similar to the actual changes observed by Whiting *et al.* (1979) and attributed solely to the effects of alcohol. Before that interpretation can be accepted as valid, carefully age-matched subjects must be compared for the single variable (i.e., alcohol) being evaluated.

PAUL V. DESMOND & STEVEN SCHENKER

Vanderbilt University School of Medicine & Nashville V. A. Medical Center, Nashville, Tennessee 37203, USA

Received March 13, 1979

References

- ROBERTS, R.K., WILKINSON, G.R., BRANCH, R.A. & SCHENKER, S. (1978). Effect of age and parenchymal liver disease on the disposition and elimination of chlordiazepoxide (Librium). *Gastroenterology*, **75**, 479–485.
- SHADER, R.I., GREENBLATT, D.J., HARMATZ, J.S., FRANKE, K. & KOCH-WESER, J. (1977). Absorption and

- disposition of chlordiazepoxide in young and elderly male volunteers. *J. clin. Pharmac.*, **17**, 709–718.
- WHITING, B., LAWRENCE, J.R., SKELLERN, G.G. & MEIER, J. (1979). Effect of acute alcohol intoxication on the metabolism and plasma kinetics of chlordiazepoxide. *Br. J. clin. Pharmac.*, **7**, 95–100.

CHLORDIAZEPOXIDE IN ACUTE ALCOHOLISM

We would like to acknowledge the points made in Desmond & Schenker's (1979) letter about the age-dependency of chlordiazepoxide disposition. Unfortunately, our data was insufficient to show this but we did emphasize that the increased variability associated with the alcoholic data may have been partly due to age. Our inability to show any

significant correlation between age and chlordiazepoxide half-life or clearance in the controls or in the alcoholics, or in the group as a whole, prompted us to analyse the data in terms of alcohol as the single most important variable. Recently, Sellers, Greenblatt, Zilm & Degani (1978) have also shown that alcohol causes alterations in chlordiazepoxide pharmaco-