possible that changes in sweat gland responsiveness in anxiety states are different in these two skin regions.

SUWANTI ISKANDAR, C.M. BRADSHAW & E. SZABADI

Department of Psychiatry, University of Manchester, Stopford Building, Oxford Road, Manchester M13 9PT

Received March 18, 1980

References

- BENJAMIN, F.B. (1953). Sweating responses to local heat application. *J. appl. Physiol.*, **5**, 594–598.
- CLUBLEY, M., BYE, C.E., HENSON, T., PECK, A.W. & RIDDINGTON, C. (1978). A technique for studying the effects of drugs on human sweat gland activity. *Eur. J. clin. Pharmac.*, **16**, 221–226.
- DOBSON, R.L. & SATO, K. (1972). The stimulation of eccrine sweating by pharmacologic agents. In *Pharmacology and the Skin*, eds. Montagna, W., Stoughton, R.B. & van Scott, J., pp. 447–475. New York: Appleton-Century-Crofts.
- LADER, M.H. (1977). *The Psychophysiology of Mental Illness*. London: Routledge and Kegan Paul.
- LADER, M.H. & WING, L. (1966). *Physiological Measures, Sedative Drugs, and Morbid Anxiety*. London: Oxford University Press.
- LEWIS, D. (1960). *Quantitative Methods in Psychology*. New York: McGraw-Hill.
- OGATA, K. & ISHIHASHI, T. (1935). Sweat reflexes due to changes in the posture of the human body. *J. Orient. Med.*, **3**, 95–97.
- SILVERMAN, C.J. & POWELL, V.E. (1944). Studies on palmar sweating. Palmar sweating in a general hospital. *Psychosomat. Med.*, **6**, 243–249.
- SNAITH, R.P., BRIDGE, G.W.K. & HAMILTON, M. (1976). The Leeds scales for the self-assessment of anxiety and depression. *Br. J. Psychiat.*, **128**, 156–165.
- SZABADI, E., GASZNER, P. & BRADSHAW, C.M. (1980). Comparison of the effects of adrenoceptor and cholinergic agonists on sweat gland activity in man. *Br. J. clin. Pharmac.*, in press.
- WILKINSON, G.N. (1961). Statistical estimations in enzyme kinetics. *Biochem. J.*, **80**, 324–332.

DECREASED CONJUGATED TYRAMINE OUTPUT IN DEPRESSION: THE EFFECT OF ORAL L-CYSTEINE

We have previously shown that orally administered anhydrous sodium sulphate (300 mg) has no significant effect on the excretion of conjugated tyramine (tyramine-*O*-sulphate), after an oral tyramine load (Bonham Carter, Sandler, Sepping & Bridges, 1978a). This was true both for a group of normal subjects and a group of patients suffering from severe depression who exhibited an apparent conjugation defect as previously reported (Sandler, Bonham Carter, Cuthbert & Pare, 1975; Bonham Carter, Sandler, Goodwin, Sepping & Bridges, 1978b), despite claims by Smith & Mitchell (1974) of an increase in tyramine-*O*-sulphate output after oral sodium sulphate administration. However, this finding was hardly surprising since the gut absorbs inorganic sulphate with difficulty (Fingl, 1965) and it does not rule out the possibility that the depressive subjects in whom we observe the apparent conjugation defect are suffering from chronic sulphate depletion associated with persistent administration of drugs which form sulphate conjugates (Levy & Matsuzawa, 1967; Bennett, Blackwell & Davies, 1975). The source or inorganic sulphate in the body is thought to be the sulphur containing amino acids such as cysteine (Sörbo, 1965) and an inadequate intake of cysteine

and other sulphur containing amino acids in the diet could be another possible explanation for the apparent conjugation defect observed in depressive illness. We set out to test this possibility by giving a dietary supplement of L-cysteine to normal subjects and depressed patients using a similar design to Levy & Matsuzawa (1967) and examining the effect on the conjugation of orally administered tyramine.

The experiment described was carried out on a group of six normal volunteers (three men and three women) and six severely depressed patients (all women) whose illnesses were resistant to all standard forms of treatment and who were about to undergo modified leucotomy (Knight, 1965; Göktepe, Young & Bridges, 1975; Bridges & Bartlett, 1977). The subjects were given 100 mg tyramine (125 mg tyramine hydrochloride) orally on one day and two successive 3 h urine collections were made as previously described (Sandler *et al.*, 1975). The specimens were acidified, diluted if necessary and an aliquot stored deep frozen until analysis (Bonham Carter *et al.*, 1978b). After a time interval of at least 48 h the tyramine loading was repeated with additional L-cysteine. L-Cysteine (1 g, as three capsules of 333 mg) was given orally every hour from

$|F| > 0.05, |F| < 0.05$

3 h before the tyramine loading until 1 h before the second urine collection was completed, 9 g L-cysteine in all. Two successive 3 h urine collections were made after tyramine loading and stored as before. The second part of the experiment was much more difficult to carry out correctly and had to be repeated by some subjects.

Free and total tyramine levels were measured in all the urine samples as previously described (Sandler *et al.*, 1975; Bonham Carter *et al.*, 1978a, 1978b). Conjugated tyramine levels were then obtained by difference. Total *p*-hydroxyphenylacetic acid (*p*-HPAA) levels were also measured in these samples by a gas chromatographic procedure (Bonham Carter *et al.*, 1978b). *p*-HPAA is the major urinary metabolite of orally administered tyramine and is formed by oxidative deamination of tyramine by the enzyme monoamine oxidase.

Cysteine administration brought about a significant increase in the output of conjugated tyramine after tyramine ingestion both in the normal subjects (Table 1; $P < 0.001$ for 0–3 h period, $P < 0.05$ for 3–6 h period, $P < 0.005$ for combined 0–6 h period, paired *t*-test) and in the depressed patients (Table 1; $P < 0.01$ for 0–3 h period, $P < 0.001$ for combined 0–6 h period, paired *t*-test). The previously demonstrated low output of conjugated tyramine by depressed patients compared with normal subjects (Sandler *et al.*, 1975; Bonham Carter *et al.*, 1978a, 1978b) was confirmed in this group of patients with depressive illness (Table 1; $P < 0.01$ for 0–3 h period after tyramine and $P < 0.05$ for 0–3 h period and for combined 0–6 h period after tyramine plus L-cysteine, Student's *t*-test). Despite this decreased excretion by the depressed patients the percentage increase in conjugated tyramine output after cysteine was not significantly greater in the patients than in the normal subjects (Table 1).

Cysteine administration had no significant effect on the output of free tyramine after tyramine ingestion in either group of subjects (Table 1). However, free tyramine excretion was significantly lower in the depressed patients, compared with normal subjects, in the first 3 h after tyramine

loading (Table 1; $P < 0.05$, Student's *t*-test) as previously demonstrated (Bonham Carter *et al.*, 1978b).

Although ingestion of cysteine had no statistically significant effect on the excretion of *p*-HPAA in either group of subjects, the output of the acid was lower in both groups after cysteine (Table 2). The tendency for depressed patients to excrete less *p*-HPAA than normal subjects after tyramine loading (Bonham Carter *et al.*, 1978b) was demonstrated and was statistically significant for the 0–3 h collection period (Table 2; $P < 0.02$, Student's *t*-test) but not for the combined 0–6 h period. However, this finding may just reflect a delay in the excretion of the acid or absorption of tyramine since the depressed patients excreted more *p*-HPAA than normal subjects in the 3–6 h period, though not significantly so.

These findings demonstrate that neither the normal subjects nor the depressed patients are saturated with sulphate since both groups excreted increased amounts of conjugated tyramine after the cysteine supplement. The output of conjugated tyramine was increased by a similar proportion in both groups; the large dietary excess of cysteine did not boost the levels of the conjugate excreted by the depressed patients up to those excreted by normal subjects which indicates that our consistent findings of low conjugated tyramine output in depressive illness (Sandler *et al.*, 1975; Bonham Carter *et al.*, 1978a, 1978b) is not due to chronic depletion of sulphate by drugs which form sulphate conjugates or to an inadequate intake of cysteine and other sulphur containing amino acids in the diet.

The enzyme which conjugates tyramine with sulphate in man is phenolsulphotransferase (PST) (Hart, Renskers, Nelson & Roth, 1979). This enzyme transfers sulphate from phosphoadenosine-5'-phosphosulphate (PAPS), 'active sulphate', to a phenolic acceptor such as tyramine, so that ingested cysteine must be converted to PAPS before it can be used for conjugation. It may be that patients with depressive illness are deficient in their ability to convert cysteine to PAPS, or in the enzyme, PST, itself. Either of these possibilities could explain the

Table 2 Excretion of *p*-HPAA (mg/3 h, mean $\pm$ s.e. mean) by control subjects and depressed patients after administration of tyramine (100 mg) and tyramine (100 mg) plus L-cysteine (9 g; 1 g/h for 9 h)

Period (h)	Control subjects		Depressed patients	
	Tyramine	Tyramine + cysteine	Tyramine	Tyramine + cysteine
0–3	61.0 $\pm$ 6.0	49.4 $\pm$ 5.7	36.8 $\pm$ 4.7†	30.7 $\pm$ 2.5
3–6	16.9 $\pm$ 2.7	17.8 $\pm$ 2.0	30.9 $\pm$ 7.1	22.5 $\pm$ 2.1
Combined 0–6	77.9 $\pm$ 5.5	67.2 $\pm$ 6.4	67.8 $\pm$ 7.2	53.2 $\pm$ 3.9

Significant results: Control subjects *v* depressed patients (Student's *t*-test) † $P < 0.02$.

apparent conjugation defect we observe in these patients. Other possible explanations for our findings are a deficit in the absorption or transport of tyramine (Bonham Carter *et al.*, 1978b) or poor absorption of dietary cysteine and other sulphur containing amino acids. We are planning to investigate some of these possibilities in the near future.

SUSAN M. BONHAM CARTER, B.L. GOODWIN & M. SANDLER

Bernhard Baron Memorial Research Laboratories and Institute of Obstetrics and Gynaecology, Queen Charlotte's Maternity Hospital, Goldhawk Road, London W6 0XG

P.K. GILLMAN & P.K. BRIDGES

The Geoffrey Knight Psychosurgical Unit, Brook General Hospital, London SE18 4LW

Received May 19, 1980

References

- BENNETT, P.N., BLACKWELL, E. & DAVIES, D.S. (1975). Competition for sulphate during detoxification in the gut wall. *Nature, Lond.*, **258**, 247–248.
- BONHAM CARTER, S., SANDLER, M., SEPPING, P. & BRIDGES, P.K. (1978a). Decreased conjugated tyramine output in depression: gastrointestinal factors. *Br. J. clin. Pharmac.*, **5**, 269–272.
- BONHAM CARTER, S., SANDLER, M., GOODWIN, B.L., SEPPING, P. & BRIDGES, P.K. (1978b). Decreased urinary output of tyramine and its metabolites in depression. *Br. J. Psychiat.*, **132**, 125–132.
- BRIDGES, P.K. & BARTLETT, J.R. (1977). Psychosurgery: yesterday and today. *Br. J. Psychiat.*, **131**, 249–260.
- FINGL, E. (1975). Laxatives and cathartics. In *The Pharmacological Basis of Therapeutics*, 5th Edn., ed. Goodman, L.S. & Gilman, A., pp. 976–986. New York: Macmillan.
- GÖKTEPE, E.O., YOUNG, L.B. & BRIDGES, P.K. (1975). A further review of the results of stereotactic subcaudate tractotomy. *Br. J. Psychiat.*, **126**, 270–280.
- HART, R.F., RENSKE, K.J., NELSON, E.B. & ROTH, J.A. (1979). Localization and characterization of phenol sulfotransferase in human platelets. *Life Sci.*, **24**, 125–130.
- KNIGHT, G. (1965). Stereotactic tractotomy in the surgical treatment of mental illness. *J. Neurol. Neurosurg. Psychiat.*, **28**, 304–310.
- LEVY, G. & MATSUZAWA, T. (1967). Pharmacokinetics of salicylamide elimination in man. *J. Pharmac. exp. Ther.*, **156**, 285–293.
- SANDLER, M., BONHAM CARTER, S., CUTHBERT, M.F. & PARE, C.M.B. (1975). Is there an increase in monoamine oxidase activity in depressive illness? *Lancet*, **i**, 1045–1048.
- SMITH, I. & MITCHELL, P.D. (1974). The effect of oral inorganic sulphate on the metabolism of 4-hydroxyphenylethylamine (tyramine) in man: tyramine-O-sulphate measurement in human urine. *Biochem. J.*, **142**, 189–191.
- SÖRBO, B. (1965). Sulfur metabolism and its clinical chemistry: sulfonic acids and inorganic sulfur compounds. *Scand. J. Lab. clin. Invest.*, **17**, 21–31.

FATTY ACIDS MODULATION OF MEAL-INDUCED VARIATIONS IN DIAZEPAM FREE FRACTION

Diazepam is highly bound to human serum albumin. Only $1.33 \pm 0.21\%$ (mean $\pm$ s.d.) of diazepam in plasma is in the unbound form over the range of concentrations normally encountered in therapy (Abel, Sellers, Naranjo, Shaw, Kadar & Romach, 1979). Since drug elimination is dependent on free fraction (Klotz, Antonin & Bieck, 1976) changes in diazepam free fraction may, therefore, cause important variations in disposition and effect. Meals induce a significant decrease in diazepam free fraction associated with reciprocal changes in total plasma

diazepam concentrations (Naranjo, Sellers, Giles & Abel, 1980). Meals decrease free fatty acids (FFA) (Barter, Carroll & Nestel, 1971; Schlierf & Dorow, 1973) and FFA modify the binding of many drugs (Birkett, Myer & Hagedorn, 1979). Therefore, FFA could be the modulators of meal-induced variations in diazepam free fraction. We studied the effect of standard breakfast on diazepam free fraction and the correlation of these changes with concomitant variations in FFA.

Eight healthy subjects (7 males, 1 female, aged