rooted is it that they would be at a loss to live without the indulgence."

¹ Allbutt TC, Dixon WE. In: Allbutt TC, Rolleston HD, eds. *System of medicine*. London: 1906.

² Michel de Crèvecoeur. *Letters from an American farmer*. 1782.

Patient information booklets

Dr P J M SLOAN (Department of Medicine, Dudley Road Hospital, Birmingham B18 7QH) writes: I was glad to see the interest aroused by my article on patient information booklets (24 March, p 915). A list of booklets by nearly 400 voluntary organisations exists.¹ No similar list of pharmaceutical companies' publications was available before my article. Patient associations produce many excellent booklets, but these usually cost money. One advantage to the doctor of pharmaceutical companies' literature is that it is free.

Any list of current items is out of date as soon as it is published. I was dependent on the cooperation of individual companies in the preparation of the list, and the items referred to by Dr H J N Bethell (19 May, p 1540) could not be included as the firm concerned did not reply to my letter.

¹ Edsall B. *Health education index*. London: B Edsall and Co Ltd, 1983.

Major disaster planning

Mr I M R LOWDON (John Radcliffe Hospital, Oxford OX3 9DU) writes: Mr A R Bliss correctly recommends that if wounds cannot be adequately treated within eight hours they should be left open (12 May, p 1433). He then makes a plea for early primary closure on the grounds that delayed closure increases the risk of infection. Delayed closure is intended to prevent infection, and if the wounds are adequately dressed in theatre and re-inspected properly there should be no risk of late contamination. The other disadvantages of delayed closure to which he refers (a further anaesthetic and delayed mobilisation) are of little consequence compared with a wound infection possibly in communication with joint or bone. The lessons of war surgery and surgical teaching are quite clear: if in doubt about the advisability of suture—don't.^{1 2}

¹ Jackson DS, Batty CG, Ryan JM, Macgregor WSP. The Falklands war: army field surgical experience. *Ann R Coll Surg Engl* 1983;**65**:281-5.

² Rains AJH, Ritchie HD, eds. *Bailey and Love. Short practice of surgery*. London: Lewis, 1981:3-6.

Poor management of unconscious rugby player watched by millions

Dr J ADNO (Parktown, Johannesburg) writes: Mr N Goddard highlighted the poor management of an unconscious player during a televised rugby match (21 April, p 1229). I have reported two similar incidents both of which occurred during televised internationals.¹ Obviously the training of those involved in such events needs to be improved.

¹ Adno J. Rugby injuries. *S Afr Med J* 1983;**64**:306

Salicylate intoxication in the elderly due to benorylate

Dr J S MCCracken (Nottingham NG5 5AP) writes: Dr T R O Beringer identifies a problem inherent in prescribing for those patients least able to identify the side effects of treatment (5 May, p 1344). Such groups include the elderly, the very young, the mentally handicapped, and also the hospital inpatient who must comply with treatment dispensed at the routine drug round.

In general practice the doctor soon learns from his articulate patients that they often have good reasons for not taking their drugs. I have for some years now reduced the prescribed dose of benorylate

suspension to a maximum 10 ml (4 g) or 5 ml (2 g) twice daily depending on whether nocturnal or diurnal relief of arthritic pain is preferred. Previously patients complained of tinnitus and dizziness. I agree with Dr Beringer that the maximum dose of benorylate in the elderly should be 4 g daily but do not think that serum salicylate concentrations really need to be monitored.

When the bowels turn to water

Dr E W VAN HEYNINGEN (North Hinksey Village, Oxford OX2 0NA) writes: I am amazed that Professor B McNichol should think that Dr Seal and I are unaware of William Brooke O'Shaughnessy's pioneer work on cholera (19 May, p 1540). If he looks at our book he will find him referred to on pages 2, 12, 20-23, 24, 49, 67, 68, 81, and 93.¹ He will also find five of O'Shaughnessy's publications listed in the references.

¹ *Cholera: the American scientific experience 1947-1980*. Heyningen WEV, Seal JR. Westview Press. 1983.

Loss of vision

Dr D McLEOD (Moorfields Eye Hospital, London EC1V 2PD) writes: Dr M Jamieson's presentation of the role of the pupillary light reflex is naive (19 May, p 1523). The "swinging flashlight test" of pupil activity is the simplest and single most useful test of visual function.¹ A relative afferent pupillary defect should always be looked for.

¹ Thompson HS, Corbett JJ, Cox TA. How to measure the relative afferent pupil defect. *Surv Ophthalmol* 1981;**26**:39-42.

Medicine in the workplace

Dr A SEMMENCE and Mrs DOROTHY RADWANSKI (Civil Service Medical Advisory Service, London SW1H 9EU) write: Professor W R Lee's article is timely (5 May, p 1326). As a House of Lords select committee noted, the problems of occupational health provision in the civil service mirror those of the United Kingdom as a whole.¹ Our service relies heavily on general practitioners to provide local advice with the support centrally of specialist occupational health physicians. If general practitioners are going to undertake this work then their nursing colleagues must be taken into account. It is exceptional for an occupational health nurse to be employed directly by a group practice. An extension of the health visitors' work into occupational health might provide the nursing element in the health centres and group practices which the select committee suggested should be used.

¹ House of Lords Select Committee on Science and Technology. *Occupational health and hygiene services*. London: HMSO, 1983.

Making an appointment system work

Dr M J WHITE (Royal Gwent Hospital, Newport, Gwent NP2 2UB) writes: Dr D N H Greig's article (12 May, p 1423) reminded me of a project which I undertook as a trainee in general practice. The rate at which surgeries were booked for an "average" week in December 1982 was analysed during the previous four weeks. Eight principals and two trainees saw 1119 patients. Of these 519 attended morning surgeries and 600 attended evening surgeries. A total of 32% of appointments for morning surgeries and 52% of appointments for evening surgeries were made on the day itself, and 28% of morning appointments and 23% of evening appointments were made on the previous working day. Only 7% of appointments were made more than one week in advance. Any patient making an appointment more than 24 hours in advance could see their first choice of doctor. The longer estab-

lished doctors tended to become booked more quickly.

Several factors affect the booking rate. Predictable factors include the number of doctors, their list sizes, their annual leave, and bank holidays. Less predictable ones include epidemics, doctors' sick leave, patients' annual leave, and temporary residents. Smaller list sizes and advance planning can improve doctors' availability and the service to patients.

Kielland's forceps delivery

Dr CONOR CARR (Portiuncula Hospital, Ballinasloe, Co Galway) writes: I feel every sympathy with Dr Sheerin (10 March, p 769) but her emotive description should not pass without comment. The major fault in her management was the use of Kielland's forceps under pudendal block anaesthesia. No rotational deliveries should be attempted without epidural or caudal block, or failing these general anaesthesia. If proper anaesthesia is used the risk of neurological sequelae becomes minimal. Following Dr Sheerin's paper I intend to carry out a retrospective survey among mothers whom I delivered with these forceps.

Alcohol withdrawal and epilepsy

Dr MARGARET A PATTERSON (London SW17) writes: Dr E B Ritson emphasises alcohol withdrawal as a possible cause of epileptic fits (28 April, p 1311). If any of our patients being withdrawn from alcohol have an epileptic fit or show preconvulsive signs, an immediate rough estimate of blood sugar concentration is made by plastic strip. Often a low value is found. An intravenous glucose injection causes immediate recovery, and anti-convulsants are not required.

Death certification in general practice

Dr CHARLES B FREER (Primary Medical Care, Faculty of Medicine, Aldermoor Health Centre, Southampton SO1 6ST) writes: The review by Dr D A Black and Dr S J Jachuck emphasises the importance of accurate information on death certificates (14 April, p 1127). Concern for such accuracy was a central feature of two previous general practice studies not discussed in this paper.^{1 2} In my study a 10% random sample of Scottish general practitioners was invited to complete a questionnaire providing details of the next death for which they were required to provide a certificate. In 18 of the 54 certificates where ischaemic heart disease was given as the cause of death the general practitioner had no supporting evidence for the diagnosis, thus supporting Walford's view that general practitioners introduce "into the coronary mortality statistics of this country a considerable though unquantifiable error."

The knowledge that ischaemic heart disease is the commonest cause of death may influence doctors faced with certifying sudden and unexpected deaths. This may lead to some valid assumptions, but since an accurate pathological diagnosis cannot always be made even with a necropsy, we shall remain unsure of the importance of this source of inaccuracy to studies based on mortality data.

¹ Walford PA. Sudden death in coronary thrombosis. *J R Coll Gen Pract* 1971;**21**:654-6.

² Freer CB. Ischaemic heart death certification in general practice. *Update* 1980;**20**:629-35.

Learning to live with cash limits

Dr D P MURRAY (Derbyshire Royal Infirmary, Derby DE1 2QY) writes: Mr G Greenshields makes the common mistake of confusing managers with administrators (5 May, p 1393). An administrator ensures that a system works, whereas a manager gets the best out of the people within an organisation. Management is to do with people not systems.