AUTHOR'S CORRECTION

Specific Inhibition of Elm1 Kinase Activity Reveals Functions Required for Early G₁ Events

Aparna Sreenivasan, Anthony C. Bishop, Kevan M. Shokat, and Douglas R. Kellogg

Sinsheimer Laboratories, Department of Molecular, Cellular, and Developmental Biology, University of California, Santa Cruz, California 95064, and Department of Cellular and Molecular Pharmacology, University of California, San Francisco, California 94143-0450

Volume 23, no. 17, p. 6327–6337, 2003. We have been unable to reproduce a control experiment reported in this paper. Specifically, we found that 25 μ M 1-naphthol-methyl PP1 (1NM-PP1) causes a G₁ delay in wild-type cells that is similar to the delay observed in *elm1-as* cells. While attempting to reproduce this control, we found that addition of supplemental adenine to yeast extract-peptone-dextrose (YPD) growth medium significantly reduces the effects of 1NM-PP1 on wild-type cells. Since our strains are *ade*[–] and grow slowly on YPD medium, we often supplement the media with additional adenine. The addition of adenine to 100 mg/liter largely eliminates the G₁ delay observed in wild-type cells in the presence of 25 μ M 1NM-PP1. We also found that the defects in septin organization that we observed before bud emergence were seen only at high concentrations of 1NM-PP1 (25 μ M). They may therefore be the combined result of full inhibition of *elm1-as* and partial inhibition of other kinases. Alternatively, high concentrations of 1NM-PP1 may be necessary to achieve full inhibition of *elm1-as*. The other results of the paper are unaffected.