Published in final edited form as:

Med Vet Entomol. 2006 March ; 20(1): 138–149.

Microgeographic breeding structure of the tsetse fly, *Glossina pallidipes* in southwestern Kenya

J. O. OUMA^{1,2}, J. G. MARQUEZ¹, and E. S. KRAFSUR¹

¹Department of Entomology, Iowa State University, Ames, Iowa 50011-3222, USA

Abstract

The origins of extant *G. pallidipes* Austen (Diptera: Glossinidae) populations in the ecologically well studied Lambwe and Nguruman valleys in Kenya are controversial because populations have recovered after seemingly effective attempts to achieve high levels of control. We investigated microgeographic breeding structure of the tsetse fly, *Glossina pallidipes* (Diptera: Glossinidae) by analyzing spatial and temporal variation at eight microsatellite loci to test hypotheses about endemism and immigration. Samples were obtained at seasonal intervals from trap sites separated by 200 m to 14 km and arranged into blocks. *G. pallidipes* populations nearest to Lambwe and Nguruman also were sampled. Spatial analysis indicated genetic differentiation by genetic drift was much less among trapping sites within Lambwe and Nguruman ($F_{ST} \leq 0.049$) than between them ($F_{ST} = 0.232$). F_{ST} between Serengeti and Nguruman was 0.16 and F_{ST} between Koder Forest and Lambwe was 0.15. The genetic variance in *G. pallidipes* explained by dry and wet seasons (0.33%) was about one-fifth the variance among collection dates (1.6%) thereby indicating reasonable temporal stability of genetic variation. Gene frequencies in Koder and Serengeti differed greatly from Lambwe and Nguruman thereby falsifying the hypothesis that Lambwe and Nguruman were repopulated by immigrants. Harmonic mean effective (= breeding) population sizes were 180 in Lambwe and 551 in Nguruman. The genetic data suggest *G. pallidipes* in Lambwe and Nguruman have been endemic for long intervals.

Keywords

Glossina pallidipes; microsatellites; genetic differentiation; genetic drift; gene flow; effective population size

Introduction

Tsetse flies, *Glossina* spp (Diptera: Glossinidae) are medically and economically important insects confined to sub-Saharan Africa and the southern tip of the Arabian peninsula. They are exclusively blood-feeding and are the only vectors of African trypanosomiasis. Mark-release-recapture studies indicate that tsetse flies in the *morsitans* group have a great capacity for dispersal and colonization of suitable habitats. Rogers (1977) observed that *G. morsitans morsitans* Westwood dispersed over two dimensions a mean 230 m daily and Vale *et al.* 1984 considered *G. pallidipes* Austen to be the most vagile tsetse fly. Furthermore, Williams *et al.* (1992) estimated *G. pallidipes* root mean square displacement rates of 360 to 1,100 m day⁻¹. The displacement rate varied with season. In principle, such rapid dispersals would homogenize allele frequencies among populations and thus prevent genetic differentiation by drift.

Correspondence: Prof. E. S. Krafur, Entomology, Iowa State University, Ames, Iowa 50011-3222, U.S.A., E-mail:ekrafur@iastate.edu.

²Current address: Trypanosomiasis Research Center, Kenya Agricultural Research Institute, P.O. Box 362, Kikuyu, Kenya

Despite the abundant evidence that indicates *morsitans* group tsetse flies have a great capacity for dispersal, genetic data from nuclear and mitochondrial DNA markers seem to indicate otherwise. Krafur *et al.* 1997 examined breeding structure of *G. pallidipes* at allozyme loci, and found surprisingly high levels of genetic differentiation among 11 populations. F_{ST} measured by the method of Weir & Cockerham (1984) was 0.24 ± 0.05 . This translates into an equivalent exchange rate of only 0.8 reproducing flies per population per generation. Genetic studies using single-strand conformation polymorphism in mitochondrial DNA indicated about the same level of gene flow among eighteen *G. pallidipes* populations (Krafur & Wohlford, 1999). More recently, comparative analyses of microsatellite, allozyme, and mitochondrial loci have provided further evidence that gene flow among *G. pallidipes* populations is restricted (Krafur 2002a; Ouma *et al.*, in press). The apparent inconsistency between ecological and genetic data calls for further investigation. Theoretically, gene flow declines with geographic distance (Wright, 1943; Rousset, 1997). Therefore, comparison of gene flow estimates with direct measures of dispersal should be made where studies are done over similar spatial scales. Krafur (2003) suggested that direct comparisons of tsetse fly genetic data with ecological data obtained at discrepant geographic scales may be misleading. Most published ecological data pertain to distances less than ten to tens of kilometers whereas genetic sampling involves distances ranging from tens to thousands of kilometers (Krafur, 2002b). Hence, it is necessary to examine local patterns of gene frequencies. The Lambwe and Nguruman regions of Kenya lend themselves to microgeographic studies because the ecology of *G. pallidipes* and trypanosomiasis has been studied extensively there (Turner, 1981, 1986; Turner and Brightwell, 1986; Welde, 1989; Brightwell *et al.*, 1992, 1997). Additionally, the Kenya Trypanosomiasis Research Institute (KETRI) has on-going tsetse monitoring programs in both regions.

G. pallidipes populations in Nguruman undergo seasonal expansions and contractions of great magnitude (Brightwell *et al.*, 1992, 1997). Trapping data suggested to the foregoing authors that *G. pallidipes* invaded Nguruman in massive numbers each wet season from an escarpment that borders Tanzania. Lesser numbers were said to invade during dry seasons. Such massive immigration predicts genetic homogeneity between the source and host populations. This suggests the hypothesis that allele frequencies would be homogeneous across the escarpment, the Nguruman valley, and adjacent lowlands.

Ecological research suggests additional genetic hypotheses. Another trapping study in Nguruman showed *G. pallidipes* to be highly aggregated even at scales of 4 km (Odulaja *et al.*, 2001), thereby suggesting the hypothesis that *G. pallidipes* populations maintain a high degree of local sub-structuring. Does the spatial aggregation predict detectable genetic differentiation?

There are additional issues that may be addressed by applying genetic methods. *G. pallidipes* populations in Nguruman and Lambwe Valley were subjected to human interference in two major ways: destruction of vegetation that forms the flies' habitat; and attempts to eradicate tsetse by using insecticides (Turner, 1986; Brightwell *et al.*, 2001) and traps (Dransfield *et al.*, 1990; Dransfield and Brightwell, 2001). Habitat destruction can lead to environmental heterogeneity which in turn could create subdivided populations adapted to local conditions (Mopper and Strauss, 1998). Also, effective control efforts cause drastic declines in population numbers. But as recently as 2003, Lambwe was heavily infested with *G. pallidipes* (Ouma and Krafur, personal observation) raising the question, did undetected, small numbers of *G. pallidipes* survive in treated areas, or did flies migrate from neighboring areas to re-infest Lambwe? To obtain information on the effects of season and control efforts on *G. pallidipes* populations, it is helpful to estimate effective population sizes (N_e).

Microsatellite markers were used to analyze genetic variation and estimate rates of gene flow among and within *G. pallidipes* trapping sites separated by distances ranging from 200 m to 14 km. Microsatellites are simple nucleotide repeats (e.g. CA-CA-CA) scattered throughout the nuclear genome (Queller *et al.*, 1993). They vary in the number of repeats (and may also demonstrate single nucleotide polymorphisms) and hence in molecular weight. The research had three broad objectives: (1) to determine the degree of isolation of Lambwe Valley *G. pallidipes* from other conspecific populations by examining genetic distances and rates of gene flow among resident populations; (2) to examine effects of season and immigration in shaping the breeding structure of *G. pallidipes* in the Nguruman, and (3) to determine if earlier control operations had an impact on the effective population sizes of *G. pallidipes*. Specifically, the following hypotheses were tested: (i) allele frequencies are homogeneous among sites and blocks incorporating Lambwe and Koderia, (ii) allele frequencies are homogeneous across a transect incorporating Serengeti, the Rift Valley escarpment bordering Tanzania, Nguruman, and adjacent lowlands, (iii) allele frequencies are temporally homogeneous.

Materials and methods

Study sites

G. pallidipes was obtained from Lambwe Valley and Nguruman in western and south-western Kenya, respectively (Fig. 1). There were 15 trapping sites in Lambwe (0°41.29'–0°35.14'S, 34°13.36'–34°18.11'E) and 27 sites in Nguruman (1°57.04'–1°48.02'S, 36°07.52'–36°03.08'E). Detailed descriptions of the geography of Lambwe Valley are available (Turner, 1986; Welde, 1989). The two regions are about 240 km apart. The Lambwe Valley encompasses 324 km², and lies between the Kanyamwa escarpment to the east and the Gwasi and Gembe Hills to the west. Lake Victoria lies 10 km to the northwest. The climate is warm and humid, with little fluctuation in mean monthly temperatures and saturation deficits (Turner, 1984). Average annual rainfall is approximately 1200 mm, with wet seasons in March-May and October-December. The southern end of the valley hosts a variety of game animals, and is fenced in a 120-km² National Park, the Ruma. There are human settlements and livestock on the park's periphery. We subdivided Ruma National Park into three sampling units: blocks A, B, and C that represent dense scrub thickets separated by grassland. Although *G. pallidipes* previously inhabited areas adjacent to Ruma such as the Kanyamwa Escarpment and Ruri Hills (Groenendijk *et al.*, 1993), its distribution seems to have shrunk due to control operations by KETRI, the Kenya Wildlife Services, and the International Centre for Insect Physiology and Ecology. Our attempts to catch flies in these Ruma neighborhoods were unsuccessful.

Nguruman is located in Kajiado district, south-west Kenya. Its altitude is about 700 m, with a mean annual rainfall of 500-700 mm seasonally distributed as in Lambwe. The Nguruman escarpment, with its steep wall rising to 1500 m, forms the western boundary (Mwangi *et al.*, 1998). Tsetse habitat in Nguruman is approximately 600 km². The vegetation consists of patches of woodland and thicket infested with two tsetse species, *G. pallidipes* and *G. longipennis* with the former being the predominant species. The lowland woodland patches are surrounded by open savannah, through which tsetse disperse during wet seasons (Brightwell *et al.*, 1997). There is abundant game in Nguruman (Tarimo-Nesbitt *et al.*, 1999). Nguruman sites were grouped into six blocks, Campsite, Kalema, Lengobei, Lengongu, Mukinyo, and Sampu.

The nearest known population tsetse centres were sampled to test hypotheses about origins of populations in Lambwe and Nguruman. Koderia is a planted coniferous forest ~ 50 km west of Lambwe. Serengeti lies in Tanzania between the Ngorongoro highlands and the Kenya-Tanzania border and is the nearest known *G. pallidipes* belt from Nguruman, about 170 km distant (Fig. 1).

Tsetse sampling

Tsetse flies were caught by using georeferenced biconical traps (Challier & Laveissiere, 1973) baited with cow urine (Baylis & Nambiro, 1993) and acetone. Traps were deployed along established transects, located in thicket and *Acacia* woodland, known habitats for *G. pallidipes*. Catches were emptied daily and the flies sexed and stored in aqueous 85% ethanol. Flies from each trap were stored separately. Geographical distances between sampling sites were determined by using a distance calculator (<http://www.nau.edu/~cvm/latlongdist.html>).

DNA extraction, PCR and microsatellite genotyping

DNA was extracted by using the cetyltrimethyl ammonium bromide (CTAB) protocol of Boyce *et al.* (1989) as modified by Shahjahan *et al.* (1995). Most flies were females. We amplified eight autosomal microsatellite loci via PCR. Primer sequences and PCR conditions were set forth in Ouma *et al.* (2003, GenBank accession numbers AY220498-AY220504) and Baker & Krafur (2001, GenBank accession number AY033512). Genotyping flies was performed by using GeneScan® and Genotyper® as described (Krafur, 2002a; Ouma *et al.*, 2003).

Data analysis

Statistical analyses were performed by using GENEPOP version 3.4 (Raymond & Rousset, 1995a, <http://wbiomed.curtin.edu.au/genepop/>), SAS (SAS, 2001), ARLEQUIN version 2.0 (Schneider *et al.*, 2000), and FSTAT version 2.9.3 (Goudet, 1995, 2001).

For each locus and sample, we estimated allele frequencies and observed and expected heterozygosities. Homogeneity of allele frequencies among trap sites and blocks was determined by using contingency chi-square tests. We also estimated probabilities of departure from Hardy-Weinberg equilibrium (HWE) within each sample at each locus. A Markov chain approximation (Guo & Thompson, 1992) was used to estimate the exact *P*-values. Significance of multiple *P*-values was tested by using Fisher's method and *P*-values adjusted by applying the sequential Bonferroni correction (Holm, 1979; Rice, 1989). Genotypic linkage disequilibrium was tested by estimating the probability of independence of genotypes for each pair of loci within each sample by using a Markov Chain approximation.

Genetic differentiation between regions, blocks, and trap sites was assessed by exact tests of homogeneity of allele distribution (Raymond & Rousset, 1995b). Significance of pairwise comparisons was evaluated after applying the Bonferroni correction. Spatial structure was analyzed by means of Wright's (1951) *F*-statistics. F_{ST} is the standardized variance in allele frequencies among populations. It also measures the average correlation of alleles within populations relative to the total. F_{IT} , the overall inbreeding coefficient, measures the correlation of alleles within individuals among all populations. F_{IS} is the nonrandom component of F_{IT} , and describes the correlation of alleles within individuals in a population. The three statistics are interrelated, thus: $(1 - F_{IT}) = (1 - F_{ST})(1 - F_{IS})$. The extent of population differentiation was quantified by computing Weir & Cockerham's (1984) estimators of *F*-statistics. Genetic structure was analyzed using an analysis of variance framework (AMOVA, Weir & Cockerham, 1984; Excoffier *et al.* 1992; Weir, 1996).

To test the hypothesis that $F_{ST} = 0$ at a locus, $X^2 = 2NF_{ST}(k - 1)$, with $(r - 1)(k - 1)$ degrees of freedom and *N* is the sample size, *k* is the number of alleles at the locus, and *r* is the number of subpopulations (Workman and Niswander, 1970). Rates of gene flow (N_m) among populations were estimated by rearranging Wright's (1969) formula, $F_{ST} = 1/(1 + 4N_m)$. Thus, $N_m = (1 - F_{ST})/4F_{ST}$, assuming equilibrium between migration and drift. N_m represents the effective number of reproducing migrants between populations per generation (Slatkin, 1993). N_m and F_{ST} were also estimated by the private allele method that can provide less biased estimates of F_{ST} where allelic diversities are large (Slatkin & Barton, 1989).

Hierarchical analysis of diversity was performed according to Nei (1987). Of particular interest is G_{ST} , analogous to Wright's F_{ST} . G_{ST} is a linear measure of genetic distance while F_{ST} is a geometric distance measure. Kruskal-Wallis one-way ANOVA was used to test hypotheses of homogeneity of allele counts and diversity estimates among populations.

Effective population size (N_e) is the size of an ideal population subject to the same effect of random genetic drift as the population under study (Wright, 1931). Effective harmonic mean population sizes (N_e) were estimated based on temporal variation in allele frequencies for samples collected at different times between May 2000 and December 2003. The standardized variance of allele frequencies (F_k) was calculated according to Pollak (1983) and discussed by Waples (1989). A variance statistic for K alleles is $F_k = 1/(K-1) \sum (x_i - y_i)^2 / (x_i + y_i)/2$, where x_i and y_i are allele i frequencies at the first and second samplings, respectively. Calculations were done by using NeEstimator version 1.3 (Peel *et al.*, 2004). Estimates of N_e assumed random sampling and negligible effects of mutation, selection, and migration (Waples, 1989). The number of generations per year for tsetse flies was taken to be eight.

It is important to note that genetic data based on microsatellite loci underestimate genetic differentiation because F_{ST} , G_{ST} , etc. cannot take values greater than the level of homozygosity (Hedrick, 1999). Homozygosities varied from 0.32 to 0.36 so the maximum differentiation we may detect is only ~ 0.36 . Microsatellites are also subject to homoplasy (Estoup *et al.*, 1995; Balloux *et al.*, 2000), whereby two or more alleles are identical in state by convergence from different ancestors. Such alleles cannot be distinguished on gels because they have the same molecular weights. Clearly, microsatellites underestimate genetic differentiation thereby overestimating magnitudes of gene flow (Nagylaki, 1998)

Results

Distribution and correlation of allele frequencies

A total of 1388 flies were genotyped. One hundred and forty-one alleles were recorded in the two regions, 35 of which were private and 23 were singular. Shared alleles are set forth in Table 1. Of the private alleles, 16 were restricted to Lambwe and 19 were unique to Nguruman. Each locus showed two or three alleles shared between the regions. Generally, the predominant alleles were the same in both regions (Fig. 2). Alleles unique to either Lambwe or Nguruman occurred in low frequencies ($\leq 5\%$).

Considering only Lambwe and Koderia, 116 alleles were detected. Thirty-eight of these (33%) were private. Nine of the 38 private alleles were restricted to Koderia. Fifty private alleles were found in Nguruman and Serengeti, 27 of which were restricted to Serengeti.

Table 1 shows correlations of allele frequencies pooled over sites within blocks. Correlation coefficients between allele frequencies in the same region were strongly positive and highly significant. Very low correlations were observed between allele frequencies in Lambwe and Nguruman indicating historically little gene flow between them. The mean correlation between Lambwe and Koderia was 28%; that between Nguruman and Serengeti was 42%. Allele frequencies were heterogeneous among blocks (Table 2).

Hardy Weinberg and linkage disequilibrium

When the Bonferroni correction was applied, 1.3% and 1.6% of the tests for heterozygote deficiency (positive F_{IS} , $P < 0.05$) were significant in Lambwe and Nguruman, respectively. The deficiency of heterozygotes occurred randomly at different sampling locations at *GpA19a*, *GpB20b*, and *GpC5b*, most likely indicating the presence of null alleles.

Among the Lambwe samples, exact tests for genotypic linkage disequilibrium within sites showed 13 significant values of 420 comparisons. Twenty-one significant values are expected at the 5% level by chance alone. In Nguruman, twenty-six of 756 comparisons were significant. When the Bonferroni correction was applied only one significant value was observed in each region. No pair of loci was in linkage disequilibrium in more than one site, indicating the studied loci were independent markers.

Microsatellite diversities

Numbers of alleles and diversities observed among blocks or sites were homogeneous in Lambwe and in Nguruman (data not shown). Diversities and numbers of alleles observed among samples became heterogeneous when Serengeti was included in the analysis of Nguruman and when Koderia was included with Lambwe ($X^2_{[7]} \geq 14.7$, $P \sim 0.04$).

Genetic differentiation and gene flow

F estimates departures from random mating. Estimates of F_{IS} averaged over loci indicated that matings within and among sites were random. Mean F_{IS} values did not differ significantly from zero (Table 3). The paucity of heterozygotes in Lambwe or Nguruman at *GpA23b*, *GpB6b*, *GpC5b* or *GpCI0b* can be attributed to null alleles. Average F_{ST} values estimated over 15 sites within Lambwe and 27 sites within Nguruman were 0.049 and 0.04, respectively. F_{ST} estimated by the private allele method were 0.064 and 0.032 among Lambwe and Nguruman populations, respectively. When sites were pooled by region, $F_{IS} = 0.065$ and $F_{ST} = 0.232$ between Lambwe and Nguruman (Table 3).

Because gene frequencies at trap site were homogeneous, they were pooled into blocks. Pairwise F_{ST} values between pooled sites are presented in Table 4. The highest F_{ST} s were observed between Lambwe and Koderia, Nguruman and Serengeti and between Lambwe and Nguruman. Differentiation between blocks within regions was also significant ($F_{ST} \leq 0.051$, $P < 0.001$, Table 5), thereby indicating a moderate level of genetic differentiation.

AMOVAs showed that most variation ($\geq 94.9\%$) was contained within trap sites (Table 5). Approximately 3 - 4% of the variance lay among sites within blocks and $\leq 1 - 2.3\%$ of the variation was attributed to differences among blocks. When alleles were pooled by region, 22.9% of the variance in allele frequencies lay between Lambwe and Nguruman, whereas 73.8% was within blocks. Only 3.3% of the variance was attributed to blocks within regions. The corresponding F -statistics showed significant differentiation between Lambwe and the Nguruman regions ($F_{RT} = 0.262$, $P < 0.0001$; Table 5). In this case, F_{RT} estimates the chance that two randomly chosen alleles in a region are identical by descent relative to the pooled total. Differentiation among blocks within regions was significantly different from zero ($F_{SR} = 0.043$, $P < 0.0001$).

Nei's hierarchical partition of microsatellite diversity is presented in Table 6. Unshared diversity between blocks (D_{BT}) in Lambwe and Koderia was 0.39, and diversity of sites within blocks, $D_{SB} = 0.59$. The corresponding differentiation between blocks, $G_{BT} = 0.25$, and differentiation among sites within blocks, $G_{SB} = 0.37$. The greater differentiation among sites within than among blocks may indicate lack of hierarchical genetic structure within Lambwe or a sampling artifact. When Koderia was excluded from the analysis differentiation between blocks became 0.05 and that among sites within blocks became 0.10. Total diversity H_T for Nguruman and Serengeti was 0.85 ± 0.05 , most of it being within sites ($H_S = 0.77$). Unshared diversity between blocks ($D_{BT} = 0.06$) was twice as great as that among sites within blocks ($H_{SB} = 0.03$). Differentiation between blocks ($G_{BT} = 0.07$) was much greater than among sites within blocks ($G_{SB} = 0.02$). These values translate into exchange of 3.4 and 11.8 reproducing individuals per generation, between blocks and among sites, respectively. To verify the

influence of Serengeti on the overall diversity and differentiation, analysis was performed without Serengeti. Unshared diversity and differentiation between blocks became 0.03 and 0.04, respectively (Table 6).

Seasonal variation in allele frequencies

No significant seasonal differences were detected in allelic or genotypic diversities. Dry season populations were more differentiated genetically than wet season populations, but not significantly so. No substantial genetic subdivision was found among collection dates. Only <1% and 1.6% of the total variance was attributed to differences in allele frequencies among seasons and sampling dates, respectively (data not shown).

Estimates of effective population sizes

Estimates for Lambwe varied from 102 to 522 reproducing flies (Table 7). Harmonic mean population sizes from Nguruman varied from 150 to 1026. N_e estimates in Nguruman were about twice as large as those in Lambwe.

Discussion

Allelic frequency distribution and genetic diversity

Microsatellite allelic frequencies were homogenous among sites within blocks in Lambwe and Nguruman. However, allele frequencies differed significantly among blocks, indicating that gene flow was insufficient to neutralize genetic drift.

Hardy-Weinberg Equilibrium

Thirteen of 64 possible R_{ma} tests and twenty-eight of 162 possible site-locus combinations in Nguruman showed deviation from HWE. All deviations were due to heterozygote deficits leading to positive F_{IS} values. Heterozygote deficiencies can, in principle, be attributed to non-random matings. Much more likely, however, the heterozygote deficits were caused by null alleles. Null alleles occur due to mutations at primer annealing sites and are a common problem with microsatellites (Lehmann *et al.*, 1996; Krafur & Endsley, 2002).

Genetic differentiation and gene flow

In Lambwe, we found no evidence of genetic differentiation among sites within blocks. There was, however, significant differentiation among blocks. The same genetic pattern was detected in Nguruman. F_{ST} s among sites in Lambwe and the Nguruman were 0.048 and 0.038, respectively. It seems, therefore, that tsetse tend to remain close to their neighbourhoods even when such neighbourhoods are within a kilometer or two of each other. There may be adaptive significance to 'staying at home' viz. conservation of water and energy reserves, that is, if mammalian hosts shelter in the same thickets.

It might be expected that a tendency to remain near a 'home' thicket would be stronger during dry seasons than wet seasons, but a statistically significant effect of season was not detected even though dry season F_{ST} s were greater than wet season indices. Analysis of Nguruman trap catches indicated a high degree of spatial aggregation and substructure (Odulaja *et al.*, 2001).

Lambwe and Koderia flies were highly differentiated ($F_{ST} \geq 0.150$) suggesting a low rate of gene flow between the two populations, equivalent to 1.2 or fewer reproducing flies every generation. This can be attributed to land-use changes. Most likely Lambwe and Koderia were previously part of a contiguous fly-belt, but became separated by habitat fragmentation caused by agriculture. Indeed, a recent land-use study in Lambwe showed a three-fold increase in cultivation over the last 50 years, and a significant decrease in woody vegetation cover and

grasslands (Muriuki *et al.*, 2005). The foregoing authors concluded that land-use changes were encouraged by numerous tsetse control programmes that caused a decline in tsetse populations and a great diminution of sleeping sickness from the valley. Tsetse populations in fragmented habitats are expected to become divergent due to founder effects, genetic drift, and reduced gene flow (Templeton *et al.*, 1990).

Serengeti flies were significantly different from Nguruman flies ($F_{ST} \geq 0.158$), which predicts an exchange rate of ~ 1.3 reproducing flies per generation. Published literature suggests that *G. pallidipes* are highly invasive and re-invade Nguruman every year from the neighboring Tanzania escarpment (Dransfield *et al.*, 1990; Williams *et al.*, 1992; Brightwell *et al.*, 1997). Indeed, it was estimated that about 100,000 females per month invaded Nguruman during the long rains and 5,000 monthly in the short rains, to which may be added a further 500 females monthly during the dry seasons (Brightwell *et al.*, 1997). If this were so, we should expect allele frequencies to be statistically homogeneous between Serengeti and Nguruman if we can assume that Serengeti is the nearest fly belt to Nguruman. Our data indicate, however, that genetic drift between Serengeti and Nguruman is very much stronger than gene flow.

At the macro level, little gene flow can be inferred between Lambwe and Nguruman: Lambwe and Nguruman are separated by highlands that are too cold for tsetse survival. Tsetse flies do not inhabit highland or grassland areas where woody vegetation is lacking (Glasgow, 1963). A shared gene pool may be explained by common ancestry and a tenuous gene flow presumably via a stepping stone model.

Effective population sizes

Effective population size is the harmonic mean number of reproducing individuals in an ideal population that shows the same allele frequencies as the sampled populations. Lambwe estimates of N_e in *G. pallidipes* were less than those in Nguruman for which we offer two reasons. Although tsetse control operations were undertaken in both areas, they were more intense in Lambwe, leading to claims of near eradication (Opiyo *et al.*, 1990). Moreover, *G. pallidipes* in Lambwe occupy a smaller and more isolated geographic habitat than Nguruman. In Lambwe, flies are mostly restricted to the Ruma National Park where bushbuck, bush-pig and buffalo inhabiting the park provide blood meals (Turner, 1986). On the other hand, Nguruman is a larger habitat than the Ruma at least by a factor of two.

To conclude, the genetic data suggest that *G. pallidipes* in Lambwe and Nguruman are geographically isolated from known nearby population centres leading to strong measures of genetic differentiation. Large effective population sizes and high genetic diversities also support the contention that *G. pallidipes* have long standing endemities in Lambwe and Nguruman. The locations support plentiful game (Turner, 1986; Tarimo-Nesbitt *et al.*, 1999) and dense thickets of vegetation that provide refugia during hot, dry intervals in which, we suggest, both tsetse and their hosts can survive virtually undetected by normal sampling procedures.

Acknowledgements

We thank Messrs. Paul Thandi and Jacob Lukango for their excellent assistance with sampling. Thanks to Dr. Japheth Kiragu, for help in identifying sampling locations and with logistics and Dr. J.M. Ndung'u, for allowing us to use KETRI facilities. Serengeti samples were provided by Dr. I. Malele of Tsetse and Trypanosomiasis Research Institute, Tanzania. This work was funded by grant AI 52456 from the USPHS NIH to ESK, and an International Atomic Energy Agency Fellowship to JOO.

References

- Baker MD, Krafur ES. Identification and properties of microsatellite markers in tsetse flies *Glossina morsitans sensu lato* (Diptera: Glossinidae). *Molecular Ecology Notes* 2001;1:234–236. [PubMed: 16479272]
- Balloux F, Brunner H, Lugon-Moulin N, Hausser J, Goudet J. Microsatellites can be misleading: an empirical and simulation study. *Evolution* 2000;54:1414–1422. [PubMed: 11005307]
- Baylis M, Nambiro CO. The effect of cattle infection by *Trypanosoma congolense* on attraction, and feeding success, of the tsetse fly *Glossina pallidipes*. *Parasitology* 1993;106:357–361. [PubMed: 8316432]
- Boyce TM, Zwick ME, Aquadro CF. Mitochondrial DNA in bark weevils: size, structure, and heteroplasmy. *Genetics* 1989;123:825–836. [PubMed: 2612897]
- Brightwell R, Dransfield RD, Williams BG. Factors affecting seasonal dispersal of the tsetse flies, *Glossina pallidipes* and *G. longipennis* (Diptera: Glossinidae) at Nguruman, south-west Kenya. *Bulletin of Entomological Research* 1992;82:167–182.
- Brightwell R, Dransfield RD, Stevenson P, Williams BG. Changes over twelve years in populations of *Glossina pallidipes* and *Glossina longipennis* (Diptera: Glossinidae) subject to varying trapping pressure at Nguruman, south-west Kenya. *Bulletin of Entomological Research* 1997;87:349–370.
- Brightwell R, Dransfield RD, Maudlin I, Stevenson P, Shaw A. Reality vs. rhetoric – a survey and evaluation of tsetse control in East Africa. *Agricultural Human Values* 2001;18:219–233.
- Challier A, Laveissière C. Un nouveau piège pour la capture des glossines (*Glossina*: Diptera, Muscidae): description et essais sur le terrain. [A new trap for catching glossina (*Glossina*: Diptera, Muscidae): description and field trials.] *Cahiers d'ORSTOM, entomologie médicale et parasitologie* 1973;11:251–262.
- Dransfield RD, Brightwell R, Kyorku C, Williams B. Control of tsetse fly (Diptera: Glossinidae) populations using traps at Nguruman, south-west Kenya. *Bulletin of Entomological Research* 1990;80:265–276.
- Dransfield RD, Brightwell R. Trap efficiency for *Glossina pallidipes* (Diptera: Glossinidae) at Nguruman, south-west Kenya. *Bulletin of Entomological Research* 2001;91:429–444. [PubMed: 11818038]
- Estoup A, Tailliez C, Cornuet JM, Solignac M. Size homoplasy and mutational processes of interrupted microsatellites in two bee species, *Apis mellifera* and *Bombus terrestris* (Apidae). *Molecular Biology and Evolution* 1995;12:1074–1084. [PubMed: 8524041]
- Excoffier L, Smouse PE, Quattro JM. Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* 1992;131:479–491. [PubMed: 1644282]
- Glasgow, J.P. (1963) *The Distribution and Abundance of Tsetse*. Pergamon Press. Oxford.
- Goudet J. Fstat, Version 1.2: a computer program to calculate *F* statistics. *Journal of Heredity* 1995;86:485–486.
- Goudet, J. (2001) *FSTAT, a Program to Estimate and Test Gene Diversities and Fixation Indices* Version 2.9.3. <http://www.unil.ch/izea/software/fstat.html> Updated from Goudet (1995)
- Groenendijk, et al. A survey of the distribution of *Glossina pallidipes* Austen around the Lambwe Valley, South Nyanza District, Kenya. *Insect Science and its Application* 1993;14:225–228.
- Guo SW, Thompson EA. Performing the exact test of Hardy-Weinberg proportion for multiple alleles. *Biometrics* 1992;48:361–372. [PubMed: 1637966]
- Hedrick PW. Highly variable loci and their interpretation in evolution and conservation. *Evolution* 1999;53:313–318.
- Holm S. A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics* 1979;6:65–70.
- Krafur ES, Griffiths N, Brockhouse CL, Brady J. Breeding structure of *Glossina pallidipes* (Diptera: Glossinidae) populations in East and southern Africa. *Bulletin of Entomological Research* 1997;87:67–73.
- Krafur ES, Wohlford DL. Breeding structure of *Glossina pallidipes* populations evaluated by mitochondrial variation. *Journal of Heredity* 1999;90:635–642. [PubMed: 10589514]

- Krafsur ES. Population structure of the tsetse fly *Glossina pallidipes* estimated by allozyme, microsatellite and mitochondrial diversities. *Insect Molecular Biology* 2002a;11:37–45. [PubMed: 11841501]
- Krafsur ES. Genetic diversity and gene flow in *morsitans* group tsetse flies. *Tsetse Trypanosomiasis Information Quarterly* 2002b;25:141–146.
- Krafsur ES. Tsetse fly population genetics: an indirect approach to dispersal. *Trends in Parasitology* 2003;19:162–166. [PubMed: 12689645]
- Krafsur ES, Endsley MA. Microsatellite diversities and gene flow in the tsetse fly, *Glossina morsitans* s.l. *Medical and Veterinary Entomology* 2002;16:292–300. [PubMed: 12243230]
- Lehmann T, Hawley WA, Collins FH. An evaluation of constraints on microsatellite loci using null alleles. *Genetics* 1996;144:1155–1163. [PubMed: 8913757]
- Mopper, S., Strauss, S., 1998. Genetic structure and local adaptation in natural insect populations: effects of ecology, life history, and behavior. Chapman & Hall, New York, NY, USA.
- Muriuki GW, Njoka TJ, Reid RS, Nyariki DM. Tsetse control and land-use change in Lambwe Valley, south-western Kenya. *Agriculture, Ecosystems And Environment* 2005;106:99–107.
- Mwangi EK, Stevenson P, Gettinby G, Reid SWJ, Murray M. Susceptibility to trypanosomosis of three *Bos indicus* cattle breeds in areas of differing tsetse fly challenge. *Veterinary Parasitology* 1998;79:1–17. [PubMed: 9777722]
- Nagylaki T. Fixation indices in subdivided populations. *Genetics* 1998;148:1325–1332. [PubMed: 9539445]
- Nei, M. (1987) Molecular evolutionary genetics. New York: Columbia University Press.
- Odulaja A, Baumgärtner J, Mihok S, Abu-zinid IM. Spatial and temporal distribution of tsetse fly trap catches at Nguruman, southwest Kenya. *Bulletin of Entomological Research* 2001;91:213–220. [PubMed: 11415476]
- Opiyo EA, Njogu AR, Omuse JK. Use of insecticide impregnated targets for control of *Glossina pallidipes* in Kenya. *Insect Science and its Application* 1990;11:417–425.
- Ouma JO, Cummings MA, Jones KC, Krafsur ES. Characterization of microsatellite markers in the tsetse fly, *Glossina pallidipes* (Diptera: Glossinidae). *Molecular Ecology Notes* 2003;3:450–453. [PubMed: 16718306]
- Ouma, J.O., Marquez, J.G. & Krafsur, E.S. (2005) Macrogeographic population structure and genetic diversity in *Glossina pallidipes*. *Bulletin of Entomological Research* (In press).
- Peel, D., Ovenden, J.R. & Peel, S.L. (2004) NeEstimator: software for estimating effective population size, Version 1.3. Queensland Government, Department of Primary Industries and Fisheries.
- Pollak E. A new method for estimating the effective population size from allele frequency changes. *Genetics* 1983;104:531–548. [PubMed: 17246147]
- Queller DC, Strassmann JE, Hughes CR. Microsatellites and kinship. *Trends in Ecology and Evolution* 1993;8:285–288.
- Raymond M, Rousset F. GENEPOP version 1.2, a population genetic software for exact tests and ecumenicism. *Journal of Heredity* 1995a;86:248–249.
- Raymond M, Rousset F. An exact test for population differentiation. *Evolution* 1995b;49:1280–1283.
- Rice WR. Analyzing tables of statistical tests. *Evolution* 1989;43:223–225.
- Rogers DJ. Study of a natural population of tsetse flies and a model for fly movement. *Journal of Animal Ecology* 1977;46:309–330.
- Rousset F. Genetic differentiation and estimation of gene flow from F-statistics under isolation by distance. *Genetics* 1997;145:11219–1228.
- SAS (2001). *SAS Language Reference*, Version 8.2. SAS Institute Inc, Cary, North Carolina.
- Schneider, S., Roessli, D. & Excoffier, L. (2000). Arlequin: a software for population genetics data analysis, version 2.000. Geneva, Switzerland: Genetics and Biometry Lab, University of Geneva.
- Shahjahan RM, Hughes KJ, Leopold RA, DeVault JD. Lower incubation temperature increases yield of insect genomic DNA isolated by CTAB method. *Biotechniques* 1995;19:333–334.
- Slatkin M. Isolation by distance in equilibrium and non-equilibrium populations. *Evolution* 1993;47:264–279.
- Slatkin M, Barton NH. A comparison of three indirect methods for estimating average levels of gene flow. *Evolution* 1989;43:1349–1368.

- Tarimo-Nesbitt RA, Golder TK, Dransfield RD, Chaudhury MF, Brightwell R. Trypanosome infection rate in cattle at Nguruman, Kenya. *Veterinary Parasitology* 1999;81:107–117. [PubMed: 10030753]
- Templeton AR, Shaw K, Routman E, Davis SK. The genetic consequences of habitat fragmentation. *Annals of the Missouri Botanical Garden* 1990;77:13–27.
- Turner DA. The colonization by the tsetse, *Glossina pallidipes* Austen, of a unique habitat—exotic coniferous plantation—with special reference to the Lambwe Valley, Kenya. *Insect Science and its Application* 1981;1:243–248.
- Turner DA. A preliminary assessment of some immediate and long-term effects of aerial spraying of endosulfan on *Glossina pallidipes* (Austen) in the Lambwe Valley, Kenya. *Insect Science and its Application* 1984;5:425–429.
- Turner DA. Tsetse and trypanosomiasis in the Lambwe Valley, Kenya. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1986;80:592–95. [PubMed: 3810794]
- Turner DA, Brightwell R. An evaluation of a sequential aerial spraying operation against *Glossina pallidipes* Austen (Diptera: Glossinidae) in the Lambwe Valley of Kenya: Aspects of the post-spray recovery and evidence of natural population regulation. *Bulletin of Entomological Research* 1986;76:331–349.
- Vale GA, Harshey BS, Hargrove JW. The use of small plots to study populations of tsetse (Diptera: Glossinidae). *Insect Science and its Application* 1984;5:403–410.
- Waples RS. A generalized approach for estimating effective population size from temporal changes in allele frequency. *Genetics* 1989;121:379–391. [PubMed: 2731727]
- Weir BS, Cockerham CC. Estimating F-statistics for the analysis of population structure. *Evolution* 1984;38:1358–1370.
- Weir, B.S. (1996) *Genetic data analysis II. Methods for Discrete Population Genetic Data* Sinauer Associates, Sunderland, Massachusetts.
- Wellde BT. Trypanosomiasis in the Lambwe Valley, Kenya. *Annals of Tropical Medicine and Parasitology* 1989;83 (Suppl):1–220. [PubMed: 2694984]
- Williams B, Dransfield R, Brightwell R. The control of tsetse flies in relation to fly movement and trapping efficiency. *Journal of Applied Ecology* 1992;29:163–179.
- Workman PL, Niswander JD. Population studies on southwestern Indian tribes. 2. Local genetic differentiation in the Papago. *American Journal of Human Genetics* 1970;22:24–49. [PubMed: 5411647]
- Wright S. Evolution in Mendelian populations. *Genetics* 1931;16:97–159. [PubMed: 17246615]
- Wright S. Isolation by distance. *Genetics* 1943;28:114–138. [PubMed: 17247074]
- Wright S. The genetic structure of populations. *Annals of Eugenics* 1951;15:323–354.
- Wright, S. (1969) *Evolution and the genetics of populations*, Vol. 2. The theory of gene frequencies. Chicago, University of Chicago Press.

Fig. 1. Kenya and Tanzania. The study region includes Kibera, Lambwe, Serengeti, and Nguruman.

Fig. 2.
Allele frequency distribution in Lambwe (black bars) and Nguruman (grey).

Table 1

Number of alleles scored (diagonally in *bold italics*), shared (above diagonal), and Kendall *Tau b* correlation coefficients (below diagonal) of allele frequencies among *G. pallidipes* in Lambwe and Nguruman.

Sample	Lambwe			Kodera	Nguruman				
	A	B	C		CampSite	Kalema	Lengobei	Lengongu	Mukinyo
A	81	72	57	29	39	29	44	39	39
B	0.74	91	60	32	41	20	45	37	38
C	0.68	0.66	64	22	33	29	37	31	32
Kodera	0.31	0.31	0.23	42	24	15	28	21	23
CampSite	0.38	0.37	0.36	0.29	56	33	48	41	40
Kalema	0.42	0.40	0.49	0.22	0.71	34	34	32	31
Lengobei	0.40	0.36	0.38	0.33	0.76	0.71	63	43	44
Lengongu	0.45	0.39	0.38	0.28	0.77	0.75	0.77	48	42
Mukinyo	0.43	0.37	0.39	0.32	0.73	0.71	0.76	0.83	50
Oloibortoto	0.31	0.28	0.33	0.29	0.70	0.68	0.76	0.69	0.66
Sampu	0.37	0.32	0.36	0.21	0.73	0.68	0.76	0.77	0.70
Serengeti	0.37	0.28	0.45	0.18	0.38	0.52	0.40	0.41	0.46

* All correlations were significant at $P < 0.001$

Table 2

G. pallidipes genetic variability and contingency chi-square analysis of homogeneity of allele frequencies among blocks.

Locus	h_e^a	Lambwe (3 blocks)		X^2	h_e^a	Nguruman (7 blocks)		X^2
		No.alleles	df			No.alleles	df	
<i>GpA19a</i>	0.582	7	12	68.5***	0.609	9	48	74.9**
<i>GpA23b</i>	0.743	16	30	49.7***	0.221	8	42	44.9***
<i>GpB6b</i>	0.655	11	20	66.9***	0.509	10	54	127.7***
<i>GpB20b</i>	0.831	27	52	497.3***	0.770	18	102	355.1***
<i>GpC5b</i>	0.680	12	22	33.5***	0.686	12	66	110.3***
<i>GpC10b</i>	0.712	16	30	310.4***	0.793	25	144	263.2***
<i>GpC26b</i>	0.687	11	20	27.2	0.717	14	78	135.9***
<i>GpCAG133</i>	0.016	6	10	16.9***	0.554	6	30	40.5***
Total	0.613	106	196	1067.4***	0.607	102	564	1160.6***

^a Expected unbiased heterozygosities (Nei, 1987) over blocks

** $P < 0.001$,

*** $P < 0.0001$

Table 3
Single locus F -statistics (Weir and Cockerham 1984) for *Glossina pallidipes*

Locus	Within regions				Between regions ^a	
	Lambwe (15 sites)		Nguruman (27 sites)		Lambwe vs. Nguruman	
	F_{IS}	F_{ST}	F_{IS}	F_{ST}	F_{IS}	F_{ST}
<i>GpA19a</i>	-0.007	0.036***	-0.029	0.003	0.028	0.081**
<i>GpA23b</i>	0.014	0.011	0.163	0.004	0.028	0.105***
<i>GpB6b</i>	0.179	0.028***	-0.270	0.061**	-0.019	0.185***
<i>GpB20b</i>	0.071	0.069***	0.078	0.129***	0.148	0.139***
<i>GpC5b</i>	0.365	0.003	0.312	0.020**	0.321	0.125***
<i>GpC10b</i>	0.105	0.158***	-0.120	0.029**	0.022	0.116***
<i>GpC26b</i>	-0.067	0.010	-0.068	0.012	-0.077	0.113***
<i>GpCAG133</i>	-0.147	0.125***	0.004	-0.002	0.001	0.460***
Means ^b	0.080	0.049	-0.016	0.040	0.065	0.232
S.E.	0.054	0.022	0.066	0.020	0.049	0.074

^aTrapping sites within a region were pooled

^bJackknife means and standard errors for F -statistics.

**
P<0.001,

P<0.0001

Pairwise genetic distance F_{ST} lower diagonal and gene flow ($N_e m$, upper diagonal) for *G. pallidipes* populations. Outgroups Kodera and Serengeti are indicated in bold.

Table 4

Population	Block A	Block B	Block C	Kodera	Campsite	Oloibortoto	Sampu	Kalena	Lengongu	Lengobei	Mukinyo	Serengeti
Block A	—											
Block B	0.025	9.75	8.37	1.64	0.77	0.77	0.89	0.73	0.75	0.88	0.80	2.02
Block C	0.029	—	83.08	1.27	0.74	0.69	0.84	0.73	0.75	0.80	0.78	2.02
Kodera	0.132	0.003	—	1.36	0.68	0.72	0.85	0.75	0.66	0.80	0.73	2.40
Campsite	0.245	0.164	0.155	—	0.58	0.61	0.66	0.53	0.57	0.66	0.61	1.07
Oloibortoto	0.246	0.253	0.270	0.300	—	15.34	17.61	4.38	124.75	35.46	83.08	1.20
Sampu	0.219	0.265	0.259	0.291	0.016	—	22.48	8.68	20.58	41.42	11.65	1.29
Kalena	0.254	0.229	0.228	0.274	0.014	0.011	—	14.45	27.53	49.75	35.46	1.51
Lengongu	0.251	0.255	0.251	0.322	0.054	0.028	0.017	—	4.96	6.33	6.00	1.47
Lengobei	0.222	0.264	0.274	0.306	0.002	0.012	0.009	0.048	—	62.25	41.42	1.15
Mukinyo	0.239	0.237	0.237	0.274	0.007	0.006	0.005	0.038	0.004	—	41.42	1.46
Serengeti	0.110	0.110	0.095	0.190	0.172	0.162	0.142	0.145	0.179	0.146	0.161	—

Table 5
AMOVA with a nested design of allele frequencies in *G. pallidipes* from Lambwe and Nguruman regions.

Source	d.f.	Variance	Percentage of variation	F-statistics
<u>Lambwe</u>				
Among blocks	2	0.05606	2.25	$F_{GT} = 0.023$
Among sites (blocks)	11	0.07189	2.89	$F_{SG} = 0.030^{***}$
Within sites	1044	2.36100	94.86	$F_{ST} = 0.051^{***}$
<u>Lambwe and Koderia</u>				
Between groups	1	0.34074	12.17	$F_{GT} = 0.122^{***}$
Among sites (groups)	12	0.10947	3.91	$F_{SG} = 0.044^{***}$
Within sites	1091	2.35063	83.93	$F_{ST} = 0.161^{***}$
Total	1105	2.80084		
<u>Nguruman</u>				
Among blocks	6	-0.00102	-0.04	$F_{GT} = 0.000$
Among sites (blocks)	20	0.09334	3.86	$F_{SG} = 0.039^{***}$
Within sites	1547	2.32477	96.18	$F_{ST} = 0.038^{***}$
<u>Nguruman and Serengeti</u>				
Between groups	1	0.40854	14.30	$F_{GT} = 0.143^{***}$
Among sites (groups)	26	0.09194	3.22	$F_{SG} = 0.038^{*}$
Within sites	1642	2.35583	82.48	$F_{ST} = 0.175^{***}$
Total	1669	2.85631		
<u>Lambwe and Nguruman</u>				
Among regions	1	0.72467	22.86	$F_{RT} = 0.262^{***}$
Among sites (regions)	40	0.10514	3.32	$F_{SR} = 0.043^{***}$
Within sites	2638	2.33972	73.82	$F_{ST} = 0.229^{***}$
Total	2679	3.16953		

* $P < 0.04$,

*** $P < 0.0001$

Table 6
Hierarchical microsatellite diversities $\pm$ SE within and among *G. pallidipes* populations (Nei 1987)

	<u>Lambwe & Koderia</u>	<u>Lambwe</u>
Diversity		
Within sites H_S	0.59 ± 0.03	0.59 ± 0.04
Sites (blocks) D_{SB}	0.59 ± 0.01	0.02 ± 0.00
Between blocks D_{BT}	0.39 ± 0.03	0.05 ± 0.00
Total H_T	1.57 ± 0.15	0.66 ± 0.43
Genetic differentiation		
Sites (blocks) G_{SB}	0.37 ± 0.06	0.03 ± 0.00
Between blocks G_{BT}	0.25 ± 0.02	0.07 ± 0.00
Gene flow rate		
Sites (blocks)	0.4	8.5
Between blocks	0.8	3.2
	<u>Nguruman & Serengeti</u>	<u>Nguruman</u>
Diversity		
Within sites H_S	0.77 ± 0.03	0.77 ± 0.03
Sites (blocks) D_{SB}	0.02 ± 0.00	0.02 ± 0.00
Between blocks D_{BT}	0.06 ± 0.00	0.03 ± 0.00
Total H_T	0.85 ± 0.05	0.82 ± 0.05
Genetic differentiation		
Sites (blocks) G_{SB}	0.02 ± 0.00	0.02 ± 0.00
Between blocks G_{BT}	0.07 ± 0.00	0.04 ± 0.00
Gene flow rate		
Sites (blocks)	11.8	9.9
Between blocks	3.4	6.5

Table 7
Effective population sizes estimated by temporal change in allele frequencies

	No. of generations	Variance (F_k)	N_e	95% C.I.
<u>Nguruman</u>				
May 2000 – Nov 2001	12	0.180	333	192 – 597
May 2000 – May 2003	24	0.289	415	246 – 686
May 2000 – July 2003	26	0.192	679	393 – 1190
May 2000 – Dec 2003	29	0.142	1026	572 – 1871
Nov 2001 – Jan 2003	14	0.283	247	150 – 401
Nov 2001 – Dec 2003	17	0.341	250	157 – 382
<u>Lambwe</u>				
June 2000 – Dec 2002	20	0.496	202	122 – 328
Jun 2000 – Dec 2003	28	0.268	522	302 – 931
Dec 2002 – Dec 2003	8	0.391	102	62 – 168