Journal of Zhejiang University SCIENCE B
ISSN 1673-1581 (Print); ISSN 1862-1783 (Online)
www.zju.edu.cn/jzus; www.springerlink.com
E-mail: jzus@zju.edu.cn

Science Letters:

A convenient and stereoselective synthesis of (Z)-allyl selenides via Sm/TMSCl system-promoted coupling of Baylis-Hillman adducts with diselenides*

LIU Yun-kui^{†1}, XU Dan-qian¹, XU Zhen-yuan^{†‡1}, ZHANG Yong-min²

(¹State Key Laboratory Breeding Base of Green Chemistry-Synthesis Technology, Zhejiang University of Technology, Hangzhou 310014, China)

(²Department of Chemistry, Zhejiang University, Hangzhou 310028, China)

[†]E-mail: ykuiliu@zjut.edu.cn; CHRC@zjut.edu.cn

Received Jan. 18, 2006; revision accepted Mar. 22, 2006

Abstract: A simple and convenient procedure for stereoselective synthesis of (Z)-allyl selenides has been developed by a one-pot reaction of diselenides with Baylis-Hillman adducts in the presence of samarium metal-trimethylsilyl chloride under mild conditions. Presumably, the diselenides are cleaved by Sm/TMSCl system to form selenide anions, which then undergo S_N2' substitution of Baylis-Hillman adducts to produce the (Z)-allyl selenides.

Key words: Stereoselective synthesis, (Z)-allyl selenides, Diselenides, Baylis-Hillman adducts, Sm/TMSCl system

doi:10.1631/jzus.2006.B0393

Document code: A

CLC number: O622.6

INTRODUCTION

Allyl selenides have drawn much attention as a class of important intermediates in organic synthesis. For example, they are useful synthetic precursors for the [1,3]- or [2,3]-sigmatropic rearrangements (Sharpless and Lauer, 1972; Fujita *et al.*, 1998; Carter and Bourland, 2000), allylic homocoupling reactions (Masugama *et al.*, 1989), and various other chemical transformations (Shea *et al.*, 1986; Back and Moussa, 2002; 2003). Generally, allyl selenides are synthesized from allylic halides with selenide anions, which could be generated by reduction of diselenides with sodium borohydride (Hori and Sharpless, 1979), samarium diiodide (Zhang *et al.*, 1993), indium (Munbunjong *et al.*, 2005) and lithium aluminium hydride (Suzuki *et al.*, 1985), etc. Other methods include reactions of organometallic reagents with organoselenium compounds (Yu *et al.*, 1997; Huang

and Chen, 2000), conversion from α -phenylseleno ketones (Nishiyama *et al.*, 1982), etc. In view of their significance in organic synthesis, it is still necessary to extend the scope of allyl selenides family and develop new synthetic routes.

The Baylis-Hillman reaction is well known as one of useful carbon-carbon bond-forming reaction in the last decades (Basavaiah *et al.*, 1996; 2003). The adducts of the reaction, 3-hydroxy-2-methylene-alkanoates (derived from acrylate esters), have been widely utilized as important precursors for stereoselective synthesis of trisubstituted alkenes (Basavaiah *et al.*, 1996; 2003). To date, a series of heteroatom-substituted allylic derivatives transferred from Baylis-Hillman adducts have been reported, among which include allyl halides (Basavaiah *et al.*, 1995; Chavan *et al.*, 1997; Yadav *et al.*, 2001), allyl sulfides (Calò *et al.*, 1988), allyl amine (Das *et al.*, 2005a), allyl ethers (Roy *et al.*, 2000), allyl phosphonates (Janecki and Bodalski, 1990), and allyl borates (Karbalka *et al.*, 2004), etc. However, to the best of our knowledge, no literature on the synthesis of Baylis-Hillman adduct-derived allyl selenides has been

[†] Corresponding author

* Project (No. 2004C21032) supported by the Key Technologies R & D Program of Zhejiang Province, China

reported. As a continuation of our interest in Baylis-Hillman reaction (Liu *et al.*, 2005; 2006) and application of samarium reagents in organic synthesis (Zhou and Zhang, 1999; Lu and Zhang, 1999), we report here a simple and convenient procedure for the stereoselective synthesis of (*Z*)-allyl selenides via one-pot reaction of diselenides with Baylis-Hillman adducts promoted by samarium-trimethylsilyl chloride system under mild conditions.

METHODS

A typical experimental procedure is as follows: under a nitrogen atmosphere, a solution of diphenyl diselenide (0.5 mmol) in THF (10 ml) was treated with samarium (1.0 mmol) and TMSCl (0.6 ml) for 2 h at room temperature until the powdered samarium was almost consumed and the yellow solution became almost colorless. Then to the mixture was added methyl 3-hydroxy-2-methylene-3-phenylpropanoate (**2a**) with the mixture being continuously stirred at room temperature for 4 h. After usual work-up, (*Z*)-methyl 2-phenylselenomethyl-3-phenylacrylate (**3a**) was obtained in 90% yield (Fig.1). It was shown in NOESY (nuclear overhauser effect spectroscopy) experiment that there is no NOE (nuclear overhauser effect) correlation between the signals of the internal olefin proton and the allylic methylene protons. Besides, according to literature (Das *et al.*, 2005b; Basavaiah *et al.*, 1994; Larson *et al.*, 1984), in the ¹H NMR spectrum of a trisubstituted alkene the β-vinylic proton, *cis*- and *trans*- to the ester group are known to resonate at δ 7.5 and δ 6.5, respectively, when alkene is substituted by an aryl group; while the same proton

cis- and *trans*- to an ester group appears at δ 6.8 and δ 5.7, respectively, when substituted by an alkyl one. All these data suggest the products adopt *Z*-stereoconfiguration.

Fig.1 Synthesis of (*Z*)-allyl selenides from Baylis-Hillman adducts and diselenides promoted by Sm/TMSCl system

RESULTS AND DISCUSSION

The results are summarized in Table 1. A variety of Baylis-Hillman adducts either possessing aryl groups (with donating as well as withdrawing functionalities) or alkyl ones were used in this reaction in order to establish the generality, with all reactions proceeding smoothly and giving desirable yield of products under similar conditions. As for diselenides, those dialkyl diselenides needed longer time to be cleaved by Sm/TMSCl system comprised of those diaryl diselenides. Besides good yields, the present process also exhibited excellent stereoselectivity. Only *Z*-stereoconfiguration of the products was obtained while no *E*-isomer was observed from the spectra.

Although the exact mechanism is not very clear to us at this moment, it is speculated that the reaction may involve a silyl enolate intermediate **2A**, which is a key intermediate for facilitating the S_N2' substitution of selenide anions to Baylis-Hillman adducts (Fig.2) (Ranu and Das, 2005).

Table 1 Synthesis of (*Z*)-allyl selenides from Baylis-Hillman adducts and diselenides promoted by Sm/TMSCl system^a

Entry	R ¹	R ²	Time (h)	Product ^b	Yield (%) ^c
1	C ₆ H ₅	C ₆ H ₅	6.0	3a	90
2	C ₆ H ₅	4-CH ₃ C ₆ H ₄	6.0	3b	92
3	C ₆ H ₅	4-CH ₃ OC ₆ H ₄	6.0	3c	88
4	C ₆ H ₅	4-ClC ₆ H ₄	6.0	3d	91
5	C ₆ H ₅	3-NO ₂ C ₆ H ₄	8.0	3e	86
6	C ₆ H ₅	2-furyl	6.0	3f	89
7	C ₆ H ₅	C ₆ H ₅ CH ₂ CH ₂	6.0	3g	92
8	C ₆ H ₅	<i>n</i> -C ₇ H ₁₅	6.0	3h	87
9	4-CH ₃ C ₆ H ₄	C ₆ H ₅	6.0	3i	89
10	C ₆ H ₅ CH ₂	4-CH ₃ C ₆ H ₄	8.0	3j	85

^a Reagents and conditions: **1** (0.5 mmol), **2** (1.0 mmol), Sm (1.0 mmol), TMSCl (0.6 ml), THF (10 ml), r.t., 6.0–8.0 h; ^b All products were characterized by ¹H NMR, MS and IR; ^c Isolated yields

Fig.2 Presumed mechanism for synthesis of (Z)-allyl selenides

CONCLUSION AND REPRESENTATIVE ANALYTICAL DATA

In conclusion, the Sm/TMSCl system-promoted one-pot coupling of diselenides with Baylis-Hillman adducts provides an efficient methodology for synthesizing (Z)-allyl selenides. The remarkable advantages of this process are its simple operation, mild reaction conditions, high yields and excellent stereoselectivity.

Representative analytical data for (Z)-allyl selenides. (Z)-methyl 2-phenylselenomethyl-3-phenylacrylate (**3a**): oil; ^1H NMR (CDCl_3 , 400 MHz) δ : 3.79 (s, 3H, OCH_3), 4.05 (s, 2H, methylene-H), 7.20~7.35 (m, 8H, ArH), 7.49~7.53 (m, 2H, ArH), 7.67 (s, 1H, ArCH=); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 25.22, 52.50, 127.78, 128.82, 128.98, 129.24, 129.63, 129.89, 130.24, 134.43, 135.11, 140.21, 168.06; IR (film) ν_{max} : 3057, 3025, 1715, 1625, 1577, 1266 cm^{-1} ; MS m/z (%): 332 (M^+). Anal. calcd. for $\text{C}_{17}\text{H}_{16}\text{O}_2\text{Se}$: C 61.64, H 4.87; Found: C 61.92, H 4.77.

References

- Back, T.G., Moussa, Z., 2002. Remarkable activity of a novel cyclic seleninate ester as a glutathione peroxidase mimetic and its facile in situ generation from allyl 3-hydroxypropyl selenide. *J. Am. Chem. Soc.*, **124**(41): 12104-12105. [doi:10.1021/ja028030k]
- Back, T.G., Moussa, Z., 2003. Diselenides and allyl selenides as glutathione peroxidase mimetics. Remarkable activity of cyclic seleninates produced in situ by the oxidation of allyl ω -hydroxyalkyl selenides. *J. Am. Chem. Soc.*, **125**(44):13455-13460. [doi:10.1021/ja0357588]
- Basavaiah, D., Sarma, P.K.S., Bhavani, A.K.D., 1994. Applications of the Baylis-Hillman reaction 2: a simple stereoselective synthesis of (E)- and (Z)-trisubstituted alkenes. *J. Chem. Soc., Chem. Commun.*, (9):1091-1092. [doi:10.1039/c39940001091]
- Basavaiah, D., Bhavani, A.K.D., Pandiaraju, S., Sarma, P.K.S., 1995. Baylis-Hillman reaction: magnesium bromide as a stereoselective reagent for the synthesis of [E]- and [Z]-allyl bromides. *Synlett.*, (3):243-244. [doi:10.1055/s-1995-4929]
- Basavaiah, D., Rao, P.D., Hyma, R.S., 1996. The Baylis-Hillman reaction: a novel carbon-carbon bond forming reaction. *Tetrahedron*, **52**(24):8001-8062. [doi:10.1016/0040-4020(96)00154-8]
- Basavaiah, D., Rao, A.J., Satyanarayana, T., 2003. Recent advances in the Baylis-Hillman reaction and applications. *Chem. Rev.*, **103**(3):811-891. [doi:10.1021/cr010043d]
- Calò, V., Lopez, L., Pesce, G., 1988. A simple three-step synthesis of β,β -disubstituted acrylates. *J. Organomet. Chem.*, **353**(3):405-409. [doi:10.1016/0022-328X(88)80328-0]
- Carter, R.G., Bourland, T.C., 2000. The first vanadium-catalyzed oxidation of aryl allylic selenides with in situ [2,3] sigmatropic rearrangement. *Chem. Commun.*, (20):2031-2032. [doi:10.1039/b006278m]
- Chavan, S.P., Ethiraj, K.S., Kamat, S.K., 1997. Facile synthesis of 2E-2-chloromethyl aryl-2-enoates. *Tetrahedron Lett.*, **38**(42):7415-7416. [doi:10.1016/S0040-4039(97)01744-9]
- Das, B., Mahender, G., Chowdhury, N., Banerjee, J., 2005a. Single-step stereoselective synthesis of (E)- and (Z)-allylamines from acetyl derivatives of Baylis-Hillman adducts. *Synlett.*, (6):1000-1002. [doi:10.1055/s-2005-864822]
- Das, B., Maji, A., Banerjee, J., Chowdhury, N., Venkateswarlu, K., 2005b. A highly efficient stereoselective synthesis of (Z)- and (E)-allyl iodides from Baylis-Hillman adducts. *Tetrahedron Lett.*, **46**(46):7913-7915. [doi:10.1016/j.tetlet.2005.09.097]
- Fujita, K., Kanakubo, M., Ushijima, H., Oishi, A., Ikeda, Y., Taguchi, Y., 1998. A symmetric [2,3] sigmatropic rearrangement of optically active allylic selenides. *Synlett.*, (9):987-988.
- Hori, T., Sharpless, K.B., 1979. Conversion of allylic phenylselenides to the rearranged allylic chlorides by N-chlorosuccinimide. Mechanism of selenium-catalyzed allylic chlorination of beta-pinene. *J. Org. Chem.*, **44**(23):4208. [doi:10.1021/jo01337a047]
- Huang, Y., Chen, R.Y., 2000. A novel synthesis of allyl selenides by reaction of an organosamarium reagent with alkyl selenocyanates. *Synth. Commun.*, **30**(2):377-381.
- Janecki, T., Bodalski, R., 1990. A convenient method for the synthesis of substituted 2-methoxycarbonyl- and 2-cyano-allylphosphonates. The allyl phosphite-allylphosphonate rearrangement. *Synthesis*, (9):799-801. [doi:10.1055/s-1990-27019]
- Kabalka, G.W., Venkataiah, B., Dong, G., 2004. Pd-catalyzed cross-coupling of Baylis-Hillman acetate adducts with bis(pinacolato)diboron: an efficient route to functionalized allyl borates. *J. Org. Chem.*, **69**(17):5807-5809. [doi:10.1021/jo0492618]

- Larson, G.L., de Kaifer, C.F., Seda, R., Terres, L.E., Ramirez, J.R., 1984. A stereoselective, two-step preparation of α -alkyl- α , β -unsaturated esters. *J. Org. Chem.*, **49**(18):3385-3386. [doi:10.1021/jo00192a028]
- Liu, Y.K., Li, J., Zheng, H., Xu, D.Q., Xu, Z.Y., 2005. An atom-economical and environmentally benign preparation of unsymmetrical bis-allyl ethers via dimerization of Baylis-Hillman adducts catalyzed by cesium hydroxide monohydrate. *Synlett.*, (19):2999-3001.
- Liu, Y.K., Xu, X.S., Zheng, H., Xu, D.Q., Xu, Z.Y., 2006. A facile and stereoselective synthesis of unsymmetrical dialkylsulfides via indium-promoted one-pot reaction of Baylis-Hillman acetates, sodium thiosulfate, and allyl bromide. *Synlett.*, (4):571-574.
- Lu, G.L., Zhang, Y.M., 1999. Facile formation of samarium(III) selenoates from potassium iodide catalyzed reaction of samarium metal and selenides, and their application in synthesis of selenoesters. *Synth. Commun.*, **29**(2):219-225.
- Masugama, Y., Yamada, K., Shimizu, S., 1989. Hexacarbonylbenzene(0)-induced dechalcogenization of allylic sulfides, sulfones, and selenides: nucleophilic substitution and reductive dechalcogenization. *Bull. Chem. Soc. Jpn.*, **62**(9):2913-2918.
- Munbunjong, W., Lee, E.H., Chavasiri, W., Jang, D.O., 2005. Indium-mediated mild and efficient one-pot synthesis of alkyl phenyl selenides. *Tetrahedron Lett.*, **46**(50):8769-8771. [doi:10.1016/j.tetlet.2005.10.025]
- Nishiyama, H., Itagaki, K., Osaka, N., Itoh, K., 1982. Transformation of α -phenylseleno ketones to allylic selenides via migration of the phenylseleno group. *Tetrahedron Lett.*, **23**(40):4103-4106. [doi:10.1016/S0040-4039(00)88359-8]
- Ranu, B.C., Das, A., 2005. A convenient synthesis of β -phenylselenocarbonyl compounds by in-TMSCl promoted cleavage of diphenyl diselenide and subsequent michael addition. *Adv. Synth. Catal.*, **347**(5):712-714. [doi:10.1002/adsc.200404355]
- Roy, O., Riahi, A., Henin, F., Muzart, J., 2000. Synergy or competition between palladium-catalysis and KF/alumina-mediation for the allylic substitution of the acetates of Baylis-Hillman adducts by phenols. *Tetrahedron*, **56**(41):8133. [doi:10.1016/S0040-4020(00)00727-4]
- Sharpless, K.B., Lauer, R.F., 1972. Facile thermal rearrangements of allyl selenides and diselenides [1,3] and [2,3] shifts. *J. Org. Chem.*, **37**(24):3973-3974. [doi:10.1021/jo00797a058]
- Shea, R.G., Fitzner, J., Fankhauser, J.E., Spaltenstein, A., Carpino, P.A., Peevey, R.M., Pratt, D.V., Tenge, B.J., Hopkins, P.B., 1986. Allylic selenides in organic synthesis: new methods for the synthesis of allyl amines. *J. Org. Chem.*, **51**(26):5243-5252. [doi:10.1021/jo00376a037]
- Suzuki, H., Yoshinaga, M., Takaoka, K., Hiroi, Y., 1985. A simple synthesis of aryl difluoromethyl selenides and tellurides. *Synthesis*, (5):497-499. [doi:10.1055/s-1985-31245]
- Yadav, J.S., Reddy, B.V.S., Madan, C., 2001. Montmorillonite clay-catalyzed stereoselective syntheses of aryl-substituted (*E*)- and (*Z*)-allyl iodides and bromides. *New J. Chem.*, **25**(9):1114-1117. [doi:10.1039/b103850h]
- Yu, M.X., Zhang, Y.M., Bao, W.L., 1997. The synthesis of allyl selenides by organosamarium reagent. *Synth. Commun.*, **27**(4):609-613.
- Zhang, Y.M., Yu, Y.P., Lin, R.H., 1993. Cleavage of selenium-selenium and tellurium-tellurium bond by samarium diiodide: synthesis of selenoesters, telluroesters, unsymmetrical alkyl phenyl selenides and tellurides. *Synth. Commun.*, **23**(2):189-193.
- Zhou, L.H., Zhang, Y.M., 1999. Low-valent titanium induced reductive coupling of diaryl diselenides with acid chlorides or acid anhydrides: facile synthesis of selenoesters. *J. Chem. Res.*, (1):28-29. [doi:10.1039/a803235a]

Editors-in-Chief: Pan Yun-he & Peter H. Byers
ISSN 1673-1581 (Print); ISSN 1862-1783 (Online), monthly

Journal of Zhejiang University
SCIENCE B

www.zju.edu.cn/jzus; www.springerlink.com
jzus@zju.edu.cn

JZUS-B focuses on "Biomedicine, Biochemistry & Biotechnology"

JZUS-B online in PMC: <http://www.pubmedcentral.nih.gov/tocrender.fcgi?journal=371&action=archive>