Journal of Zhejiang University SCIENCE B
 ISSN 1673-1581 (Print); ISSN 1862-1783 (Online)
 www.zju.edu.cn/jzus; www.springerlink.com
 E-mail: jzus@zju.edu.cn

Science Letters:

Functional analysis of helicase and three tandem HRDC domains of RecQ in *Deinococcus radiodurans*^{*}

HUANG Li-fen^{1,2}, HUA Xiao-ting¹, LU Hui-ming¹, GAO Guan-jun¹,
 TIAN Bing¹, SHEN Bing-hu³, HUA Yue-jin^{†‡1}

(¹Institute of Nuclear-Agricultural Science, Zhejiang University, Hangzhou 310029, China)

(²Jiangsu Province Key Lab of Crop Genetics and Physiology, Yangzhou University, Yangzhou 225009, China)

(³Division of Molecular Biology, City of Hope National Medical Center and Beckman Research Institute, Duarte, CA 91010-3000, USA)

[†]E-mail: yjhua@zju.edu.cn

Received Mar. 27, 2006; revision accepted Mar. 31, 2006

Abstract: RecQ is a highly conserved helicase necessary for maintaining genome stability in all organisms. Genome comparison showed that a homologue of RecQ in *Deinococcus radiodurans* designated as DR1289 is a member of RecQ family with unusual domain arrangement: a helicase domain, an RecQ C-terminal domain, and surprisingly three HRDC domain repeats, whose function, however, remains obscure currently. Using an insertion deletion, we discovered that the DRRecQ mutation causes an increase in gamma radiation, hydroxyurea and mitomycin C and UV sensitivity. Using the shuttle plasmid pRADK, we complemented various domains of the *D. radiodurans* RecQ (DRRecQ) to the mutant in vivo. Results suggested that both the helicase and helicase-and-RNase-D-C-terminal (HRDC) domains are essential for complementing several phenotypes. The complementation and biochemical function of DRRecQ variants with different domains truncated in vitro suggested that both the helicase and three HRDC domains are necessary for RecQ functions in *D. radiodurans*, while three HRDC domains have a synergistic effect on the whole function. Our finding leads to the hypothesis that the RecF recombination pathway is likely a primary path of double strand break repair in this well-known radioresistant organism.

Key words: RecQ, *Deinococcus radiodurans*, Helicase, HRDC
doi:10.1631/jzus.2006.B0373

Document code: A

CLC number: Q3

INTRODUCTION

Deinococcus radiodurans exhibits extreme resistance to ionizing radiation, UV-ray, desiccation and a variety of DNA damaging agents without any

mutation (Minton, 1994). With the release of the genome sequence of *D. radiodurans* R1 sequenced and annotated (White *et al.*, 1999), many functional genes involved in DNA repair, such as *recA*, *ssb* and so on have been studied (Egginton *et al.*, 2004; Daly *et al.*, 1994). However, the molecular mechanisms underlying its radiation resistance remain unclear.

As a recombination-specific DNA helicase from the SF2 family, RecQ helicases are highly conserved in evolution and are required for maintaining genome stability in all organisms (Khakhar *et al.*, 2003). Loss of RecQ helicase leads to a breakdown in the maintenance of genome integrity, in particular hyper-recombination. RecQ has been the focus of considerable attention recently because mutations in three *recQ* genes are responsible for human disorders

[‡] Corresponding author

^{*} Project supported by the National Basic Research Program (973) of China (No. 2004CB19604), Distinguished Young Scientist of China (No. 30425038), and the National Natural Science Foundation of China (No. 30330020)

Note: This is a part of our work submitted to DNA Repair on November 30, 2005. We were informed by the editor for revision on January 9, 2006. During revision, a similar work entitled "Three HRDC domains differentially modulate *Deinococcus radiodurans* RecQ helicase biochemical activity" by Micheal P. Killoran and James L. Keck was submitted to J. Biol. Chem. on January 3, 2006 and online on March 9, 2006

associated with cancer predisposition.

RecQ is a multifunctional protein and likely plays roles in many facets of DNA metabolism, such as replication, recombination, telomere maintenance and DNA damage response. In vitro, RecQ helicase initiates strand exchange and unwinds a wide variety of DNA substrates homologous pairing intermediates, including G4 DNA, three- or four-strand DNA structure including D-loops, forked structure and Holliday junction (Huber *et al.*, 2002). It is also suggested that RecQ generates an initiating signal that can recruit RecA for SOS induction and recombination at stalled replication forks (Hishida *et al.*, 2004).

The RecQ family can be distinguished from others by its three conserved domains referred to as the helicase, RQC and HRDC domains (Morozov *et al.*, 1997) (Fig.1a). In *D. radiodurans*, one component of the recombinational repair system that has unusual domain architecture is the RecQ helicase, the function of which remains obscure currently. It contains three tandem copies of HRDC domain instead of single copy present in all other bacteria except *Neisseria* that similarly possesses three copies (Makarova *et al.*, 2001). The three HRDC domains share many identical residues with other HRDC domains (Fig.1b), but whether they could all contribute the extreme resistance ability in the bacterium, or whether there are similarities and differences between the DRRecQ and other RecQ family members remain unknown. Our present study was undertaken to investigate the biochemical function of the helicase domain and the three tandem HRDC domains of the *D. radiodurans* RecQ protein in vivo and in vitro.

PHENOTYPE AND COMPLEMENTATION IN THE MQ MUTANT

We constructed an *recQ* disruption strain (designated MQ) in which the entire coding region of the DR1289 gene has been replaced with a kanamycin resistance cassette under the control of a constitutively expressed *D. radiodurans* promoter *groEL*. Cell survival after damage of the MQ strain showed that the DRRecQ mutation was sensitive to mitomycin C, and hydroxyurea and UV. Interestingly, we also found that when the dosage of gamma radiation up to 2 kilograys, it decreased remarkably compared to the wild type R1

(data not shown). Then, we complemented different HRDC domains truncated variants into the MQ mutant (Fig.1c) with shuttle vector pRADK. Western blot results showed that all the RecQ variants proteins could be detected against RecQ antiserum, with the molecular weight of each band corresponding to the weight of the various proteins. Therefore, we concluded that all the various RecQ proteins could be

Fig.1 (a) Representation of selected member of the RecQ helicase family. Proteins are aligned with their conserved helicase domain, RQC domain and HRDC domain. Nonconserved regions are shown in blank; (b) Alignment of the HRDC domain of DRRecQ, BLM, Sgs1, RecQ. Identical or similar residues are shaded in gray; (c) Schematic map of the complementary strain

over-expressed in MQ in vivo. It was previously proposed that the RECQ/QE helicase activity is important for cellular function and that the C-terminal domain has a special function in the absence of Top3 (Nakayama *et al.*, 2004). While our data suggested that all the three HRDC domains and the helicase domain of *D. radiodurans* are all necessary for complementing the phenotype of the MQ, indicating that the three tandem HRDC domains have a special function in *D. radiodurans*, whose function may distinguished from other HRDC domain in the RecQ family.

BIOCHEMICAL CHARACTERIZATION OF THE DRRECQ PROTEIN

Four variants of the DRRecQ protein along with the full length enzyme were constructed and purified (Fig.2). Helicase-mediated unwinding of the purified DRRecQ variants was monitored by using a continuous fluorescent dye-displacement assay. Hoechst 33258 functions as suitable reporter molecule and the dsDNA substrate used here was pUC19 digested by *Hind*III with nearly blunt dsDNA ends. Higher concentrations of DRRecQ cause an increase in the proportion of DNA molecules (Fig.3). Moreover, similar to the *E. coli* RecQ, the fluorescence signal was sensitive to the concentration of magnesium and sodium chloride in the reaction. When all the variants tested showed similar activity, with more amount of HRDC domains truncated, DRRecQ Δ 1D, DRRecQ Δ 2D, DRRecQ Δ 3D had lower helicase, ATPase, and binding affinity in a synergistic manner. In contrast, DRRecQ Δ H displays no detectable helicase and ATPase activity but displays all the binding properties of almost the full length protein. Thus, we proposed that the helicase domain is indispensable for the unwinding activity of the DRRecQ protein, while the three HRDC domains may elevate the helicase function as synergistic effectors. *D. radiodurans* RecQ is a special member of the RecF pathway of homologous recombination, its genetic importance in vivo and in vitro has been uncovered. These finding leads to the hypothesis that the RecF recombination pathway is likely a primary path of double strand break repair in this well-known radioresistant organism.

Fig.2 Identification of the recombinant DRRecQ variant protein. Ten micrograms of each protein were separated on a 12% SDS-PAGE gel and stained with Coomassie Brilliant Blue

Lane 1: DRRecQ; Lane 2: DRRecQ Δ 1D; Lane 3: DRRecQ Δ 2D; Lane 4: DRRecQ Δ 3D; Lane 5: DRRecQ Δ H; M: Marker

Fig.3 Rate of *Hind*III-cut pUC19 unwinding of fluorescent dye-displacement helicase assay with various concentrations of DRRecQ protein

The propagation of the HRDC domain in *D. radiodurans* was proposed to contribute to the repair phenotype, given the interactions of RecQ with RecA in recombination. The work presented here has uncovered the biochemical functions and phenotype of *D. radiodurans* RecQ and provides important clues for understanding the complex interplay between DNA replication, recombination and DNA repair.

References

- Daly, M.J., Ouyang, L., Fuchs, P., Minton, K.W., 1994. In vivo damage and *recA*-dependent repair of plasmid and chromosomal DNA in the radiation-resistant bacterium *Deinococcus radiodurans*. *J. Bacteriol.*, **176**(12): 3508-3517.
- Eggington, J.M., Haruta, N., Wood, E.A., Cox, M.M., 2004. The single-stranded DNA-binding protein of *Deinococcus radiodurans*. *BMC Microbiol.*, **4**(1):2. [doi:10.1186/1471-2180-4-2]

- Hishida, T., Han, Y.W., Shibata, T., Kubota, Y., Ishino, Y., Iwasaki, H., Shinagawa, H., 2004. Role of the *Escherichia coli* RecQ DNA helicase in SOS signaling and genome stabilization at stalled replication forks. *Genes Dev.*, **18**(15):1886-1897. [doi:10.1101/gad.1223804]
- Huber, M.D., Lee, D.C., Maizels, N., 2002. G4 DNA unwinding by BLM and Sgs1p: substrate specificity and substrate-specific inhibition. *Nucleic Acids Res.*, **30**(18):3954-3961. [doi:10.1093/nar/gkf530]
- Khakhar, R.R., Cobb, J.A., Bjergbaek, L., Hickson, I.D., Gasser, S.M., 2003. RecQ helicases: multiple roles in genome maintenance. *Trends. Cell Biol.*, **13**(9):493-501. [doi:10.1016/S0962-8924(03)00171-5]
- Makarova, K.S., Aravind, L., Wolf, Y.I., Tatusov, R.L., Minton, K.W., Koonin, E.V., Daly, M.J., 2001. Genome of the extremely radiation-resistant bacterium *Deinococcus radiodurans* viewed from the perspective of comparative genomics. *Microbiol. Mol. Biol. Rev.*, **65**(1):44-79. [doi:10.1128/MMBR.65.1.44-79.2001]
- Minton, K.W., 1994. DNA repair in the extremely radioresistant bacterium *Deinococcus radiodurans*. *Mol. Microbiol.*, **13**(1):9-15.
- Morozov, V., Mushegian, A.R., Koonin, E.V., Bork, P., 1997. A putative nucleic acid-binding domain in Bloom's and Werner's syndrome helicases. *Trends Biochem. Sci.*, **22**(11):417-418. [doi:10.1016/S0968-0004(97)01128-6]
- Nakayama, M., Kawasaki, K., Matsumoto, K., Shibata, T., 2004. The possible roles of the DNA helicase and C-terminal domains in RECQ5/QE: complementation study in yeast. *DNA Repair*, **3**(4):369-378. [doi:10.1016/j.dnarep.2003.12.001]
- White, O., Eisen, J.A., Heidelberg, J.F., Hickey, E.K., Peterson, J.D., Dodson, R.J., Haft, D.H., Gwinn, M.L., Nelson, W.C., Richardson, D.L., et al., 1999. Genome sequence of the radioresistant bacterium *Deinococcus radiodurans* R1. *Science*, **286**(5444):1571-1577. [doi:10.1126/science.286.5444.1571]

Editors-in-Chief: Pan Yun-he & Peter H. Byers
ISSN 1673-1581 (Print); ISSN 1862-1783 (Online), monthly

Journal of Zhejiang University **SCIENCE B**

www.zju.edu.cn/jzus; www.springerlink.com

jzus@zju.edu.cn

JZUS-B focuses on "Biomedicine, Biochemistry & Biotechnology"

JZUS-B online in PMC: <http://www.pubmedcentral.nih.gov/tocrender.fcgi?journal=371&action=archive>

Welcome Contributions to JZUS-B

Journal of Zhejiang University SCIENCE B warmly and sincerely welcome scientists all over the world to contribute Reviews, Articles and Science Letters focused on **Biomedicine, Biochemistry and Biotechnology**. Especially, Science Letters (3~4 pages) would be published as soon as about 30 days (Note: detailed research articles can still be published in the professional journals in the future after Science Letters is published by *JZUS-B*).