Published in final edited form as:

Am J Trop Med Hyg. 2006 May ; 74(5): 786–794.

PHENOTYPIC PLASTICITY AND GEOGRAPHIC VARIATION IN THERMAL TOLERANCE AND WATER LOSS OF THE TSETSE *GLOSSINA PALLIDIPE* (DIPTERA: GLOSSINIDAE): IMPLICATIONS FOR DISTRIBUTION MODELLING

JOHN S. TERBLANCHE*, C. JACO KLOK, ELLIOT S. KRAFSUR, and STEVEN L. CHOWN

Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University,
Stellenbosch, South Africa, Department of Entomology, Iowa State University, Ames, Iowa 50011

Abstract

Using the tsetse, *Glossina pallidipes*, we demonstrate that physiological plasticity (resulting from temperature acclimation) accounts for among-population variation in thermal tolerance and water loss rates. Critical thermal minimum (CT_{Min}) was highly variable among populations, seasons and acclimation treatments, and the full range of variation was $9.3^{\circ}C$ (maximum value = $3.1 \times$ minimum). Water loss rate showed similar variation (max = $3.7 \times$ min). By contrast, critical thermal maxima (CT_{Max}) varied least among populations, seasons and acclimation treatments, and the full range of variation was only $c. 1^{\circ}C$. Most of the variation amongst the four field populations could be accounted for by phenotypic plasticity, which in the case of CT_{Min} develops within five days of temperature exposure, and is lost rapidly on return to the original conditions. Limited variation in CT_{Max} supports bioclimatic models that suggest tsetse are likely to show range contraction with warming from climate change.

INTRODUCTION

Climate change and its interaction with human land use patterns are amongst the most significant threats facing biodiversity and society.¹ Therefore, forecasts of the interactive effects of climate change and land use are of considerable importance for predicting the future distribution and abundance of species, and for enabling society to mitigate and cope with the effects of these changes.¹ Typically, predictions of climate change effects on species ranges are based on bioclimatic models,^{2,3} although there is substantial variety in the way in which this modelling is done. These models are often able to explain much of the variation in species distributions, and abundances.^{4,5,6,7} They are therefore considered one of the most effective ways to obtain a rapid estimate of likely alterations in species' distributions.² Nonetheless, they have been criticized on a variety of grounds.⁸

From a physiological perspective, the bioclimatic modelling approach raises concerns from three principal perspectives. Spatial variation in population responses to the environment is often not considered;⁹ the rapid alterations to phenotypes that might take place via phenotypic plasticity in the form of developmental plasticity, acclimation, and hardening are typically ignored;¹⁰ and the likely outcome of covariation among abiotic variables is often not adequately assessed.¹¹ In consequence, it has been proposed that physiological investigations

JST, SLC:jst@sun.ac.za; slchown@sun.ac.za; Tel: +27 21 808 2403, Fax: +27 21 808 2405. Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Private Bag X1, Matieland, 7602, South Africa.
CJK:cjklok@asu.edu; School of Life Sciences, Box 874501, Arizona State University, Tempe, AZ, 85287-4501, U.S.A.
ESK:ekrafsur@iastate.edu; Department of Entomology, Iowa State University, Ames, Iowa 50011, U.S.A.

and biophysical modelling should be used in concert with large scale bioclimatic investigations of species responses to understand what the future might hold for various taxa under a changing climate.¹⁰

These concerns are equally, if not more, pertinent to understanding changes in the distributions of arthropod-borne diseases.¹² Nonetheless, it has been demonstrated that climate change is likely to alter, or already has altered, the geographic distribution of at least some of these diseases,^{12,13} although in other cases the effects are less straightforward to discern.^{11,14} Given that the distributions of vector species are important in influencing changing patterns of disease transmission,^{12,13} considerable attention has been given to the likely effects of environmental change on vectors. For Africa, much of the attention has focussed on malaria and its mosquito vectors.¹⁴ However, tsetse (*Glossina* spp. Diptera: Glossinidae), which are long-lived, haematophagous flies, with low reproductive rates, have also been the subject of consideration because they carry trypanosomes that have serious implications for human and animal health, and are considered a major limitation to the socio-economic development of the continent.^{7,15}

Several studies have predicted that the ranges of African trypanosomiases and their tsetse vectors are likely to change under forecast climate scenarios.^{16,17,18} These predictions arise from the fact that abiotic variables that will be affected by climate change, such as land surface temperature and saturation deficit, and other integrated environmental variables (e.g. the Normalized Difference Vegetation Index (NDVI)), are excellent correlates of tsetse distributions and abundances,^{4,5,6,7} explaining much of their variation. In some cases the mechanisms underlying these relationships are likely to be indirect.^{7,4,19} However, the direct role of abiotic environmental variables in affecting tsetse life history parameters is well established.²⁰ Temperature has major effects on birth rates, development, and mortality, and especially strong relationships exist between adult mortality and ambient temperature, which in some species are markedly non-linear.^{20,21} Water availability is also known to affect mortality rates; puparia and teneral adults are most sensitive to dry conditions, whilst mature adults are most sensitive to temperature.²⁰

Despite the foregoing information on how responses to climate might vary among different tsetse species and subspecies, based both on remote sensing work,^{5,6} and laboratory-based physiological studies,²² inter-population physiological variability and the rate of development and extent of phenotypic plasticity in tsetse are poorly understood (although some behavioural variation has been described²³). Nonetheless, it is clear that if the responses of these species to climate change are to be fully comprehended, then such information, combined with estimates of gene flow^{24,25} and dispersal rates^{20,25,26} is essential. Here, we therefore undertake such a study by investigating between-population variability in seasonal acclimatization, and the extent and time course of phenotypic plasticity of thermal tolerance and desiccation resistance in *Glossina pallidipes* Austen.

MATERIALS AND METHODS

Field sampling and study populations

Adult flies were sampled from four locations representing different thermal habitats in Kenya, East Africa (Table 1). To determine seasonal variation in physiology, the Nguruman District (mid-altitude) and Lambwe District (high-altitude) populations were sampled twice within the same year in mid-July to mid-August 2003 (the end of the long rainy season) and in mid-November to mid-December 2003 (the end of the short rainy season). The Narok (highest altitude) and Kwale Districts (low-altitude) were sampled in November-December 2003. For each of the field experiments, flies were collected from odour-baited, biconical traps¹⁵ set out in the respective localities (key attractive components: 4-methyl-phenol, 3-*n*-propynol, and

acetone). Following removal from the traps, the flies were transported in an insulated container, lined with moist paper towel, to field laboratories. These were located a maximum of two hours drive away from each of the sampling sites of a given district. At the laboratories, the flies were separated into groups for each of the experimental procedures, after which experiments commenced immediately. Depending on the time of the experiment, flies were collected either in the morning or in the late afternoon, usually from 08h00 to 10h00 and from 17h00 to 18h00. Thermal limit assessments were performed during the day using flies sampled in the morning, while desiccation experiments took place during the night using afternoon-collected flies.

Critical thermal limits

An insulated system of eleven double-jacketed isolation chambers was connected to a programmable water bath (LTD 20 with PZ1 programmer, Grant Instruments, Cambridge, UK) which regulated water flow around the chambers. A single fly was placed into each chamber and a 40 SWG type-T thermocouple was inserted into the control chamber to measure chamber temperature. The flies were allowed to equilibrate for ten minutes at either 16 or 35 °C before either minimum or maximum critical thermal limit assessments commenced, respectively.²⁷ After equilibration, chamber temperature was lowered or raised at 0.25 °C.min⁻¹. Preliminary experiments confirmed that body temperature did not lag behind or exceed chamber temperature at this rate of temperature change. Critical thermal minimum (CT_{Min}) was defined as the loss of coordinated muscle function, and critical thermal maximum (CT_{Max}) was defined as the onset of muscle spasms. The temperature at which either of these reactions occurred was recorded for each individual. These endpoints are readily identifiable for any species once an observer is practiced,²⁷ variance about the endpoints is typically low, and here a single observer (CJK) undertook all of this work. Moreover, this experimental procedure has been verified using thermolimit respirometry,²⁸ is widely used to assess thermal limits,²⁹ and the observer typically was not informed which acclimation treatment was being assessed. The procedure was repeated three times for both CT_{Min} and CT_{Max} to give a total n = 30 per population and trait. Preliminary, replicated experiments found no effect of gender, age or feeding status on critical thermal limits (unpublished data, C.J. Klok and J.S. Terblanche). Critical thermal maxima were lethal while minima were not. However, even in the case of CT_{Min} all experimental animals were discarded following a trial.

Water loss rates

Male flies (n = 16), held individually in 5 ml cuvettes were subjected to desiccation in flowing air (< 2.5% relative humidity, R.H.) for ten hours at 24.0 ± 1.0 °C (PTC-1 Peltier-controlled temperature cabinet, Sable Systems, Las Vegas, Nevada).³⁰ Air was pumped through a scrubbing column, containing silica gel and Drierite to remove residual water, and then into a mass flow controller (MFC). The MFC outlet was connected to a Sable Systems MF8 airflow manifold. Each outflow channel of the manifold was further split into two streams so that two cuvettes, each containing a fly, were attached to a single manifold channel. The air flow rate through each cuvette, tested with a second MFC, was regulated to produce a rate of 100 ml.min⁻¹. Experiments took place during the night (21h00–07h00), a period of minimal activity in tsetse, and were repeated twice per location (n = 16 per experiment), each time on individuals collected in the field on that particular day, to give a total n = 32. Each fly was weighed before and after a trial (Avery Berkel FA 304T, EU, Electronic microbalance, 0.1 mg resolution), and water loss rate expressed as mg H₂O.h⁻¹. Excretion usually takes place after a bloodmeal and here experiments took place well after such events. In addition, even in field-collected flies where the time since feeding could not be accurately determined, no excretion in the cuvettes was observed. Therefore, mass loss was considered representative of water loss even though a small amount of mass change took place as a consequence of metabolism.²² After the trials, flies were dried to constant mass (~50–60 °C for ~72 hrs) and re-weighed to determine water content. Solvent-based lipid extractions were used to determine lipid mass. Flies used in

desiccation trials were dismembered and soaked in 2 ml chloroform:methanol (2:1) solution for approximately 48 h to dissolve internal (body) lipids. Lipid content was estimated by determining dry body mass before and after lipid extraction.³¹

Activity levels

Several authors have suggested that differences in desiccation resistance among taxa might simply reflect divergence in activity levels.^{22,32} Because little is known about inter-population variation in tsetse activity levels, particularly under stressful conditions, the proportion of time individuals were active was estimated over a 30 min period for each population. Time active at 20, 24, 28 and 32 °C was determined using a single electronic activity detector (AD-1, Sable Systems) which exported voltage data to a computer via Sable Systems DATACAN V software. These experiments were performed during the same field sampling periods reported above. Voltage data were sliced from the raw Datacan V (Sable Systems) data files, copied to MS-Excel, converted to absolute values and sorted from lowest to highest. Based on preliminary observations, we considered activity as any signal > 0.01 Volt measured by the AD-1 (electronic spikes of 5 Volt were ignored and can easily be distinguished from true activity readings). The total time spent above this level of activity was expressed as a proportion of the total recording time. Whilst it is well known that activity in tsetse is sensitive to feeding state,³³ the objective here was to determine whether differences in activity level were a likely cause of any differences in water loss rates between populations, not whether the flies activity overall differs between populations.

Within-generation physiological plasticity

Logistic constraints precluded the use of field populations for investigations of phenotypic plasticity. Therefore, *G. pallidipes* from a laboratory colony maintained at the Entomology Unit, FAO/IAEA Agriculture and Biotechnology Laboratory, Seibersdorf, International Atomic Energy Agency, Vienna, Austria was used. The IAEA laboratory colony used in this study was established in 1975 from a natural population in Tororo, Uganda, close to the Kenyan border (and less than 200 km from our Lambwe field site). Gene diversities over two mitochondrial loci were within the range of 18 field populations from Ethiopia, East and southern Africa.³⁴ Therefore, the laboratory-reared flies may be considered genetically representative of flies from field populations. Puparia were immediately placed inside plastic containers and transferred to a climate chamber set to 24 °C (24.6 ± 2.5 °C, 12:12 L:D) and 76% relative humidity as described previously.³⁵ Feeding of the newly emerged adults took place using a membrane-tray system³⁶ every alternative day and at this time container locations were randomized within the climate chamber. Care was taken to ensure that all treatment groups were handled for the same duration in transferral from the climate chamber to the feeding area, and spent a similar amount of time outside the climate chamber whilst feeding (~25 min on each feeding day). Flies were held at 19, 24 and 29 °C for ten days in climate chambers with a synchronized photoperiod and constant 76% relative humidity levels. The 24 and 29 °C groups were stored in temperature-regulated climate chambers ((Labcon, South Africa) mean $\pm$ SD: 24.0 ± 2.3 [n = 1029]; 28.9 ± 1.7 °C [n = 937]) and the 19 °C group was stored in a Peltier-controlled temperature cabinet (mean $\pm$ SD: 19.4 ± 0.4 °C [n = 905]). Acclimation temperatures were chosen to represent the mean annual temperatures of the respective geographic locations sampled (reported MAT: warm-temperate midlands (< 1850 m): 18–20 °C; warm lowlands (900–1200 m) 22–24 °C; hot lowlands (0–900 m): 24–30 °C).³⁷ All experimental and rearing facilities were contained within a quarantine approved, air-conditioned laboratory. The same system and methods as those used in the field experiments for determining geographic variation were used in the laboratory determination of critical thermal limits and desiccation rate.

A separate batch of colony-reared flies was used to determine whether the temperature acclimation effect observed in the CT_{Min} of *G. pallidipes* is time- and/or temperature-dependent, and whether these changes in CT_{Min} are reversible. Flies were allowed to emerge in our laboratory as described above, and were fed twice within four days following eclosion. However, a small number of flies appeared only to feed during the second feeding round. Thus, typically after two bloodmeals, flies were randomly assigned to four temperature acclimation groups (24, 21, 19, and 16 °C, $n = 180 \sim 220$ per group, 5–6 cages per group). Humidity and photoperiod were regulated as described above, and randomization of cage locations within chambers followed the protocol adopted previously. Therefore, handling times were kept constant across all groups during feeding sessions and experimental treatments. The CT_{Min} of ten randomly selected flies from each treatment group was then determined on acclimation days 1, 3, 5, 7 and 9. After nine days of acclimation, the remaining flies were all returned to 24 °C for an additional nine-day period, after which another sample of flies was assessed on the tenth day. In this experiment, once a fly had been used for determination of CT_{Min} , it was discarded by bottling in 90% ethanol (i.e. no animals were re-used). Virgin females, separated from males into cages immediately after eclosion, were used to investigate effects of temperature acclimation on body mass. Feeding took place using the same methods described above only on days between experimental investigations (i.e. flies were always in a fasted state during experiments).

Statistical analyses

Data were assessed for normality using a Shapiro-Wilk's test prior to analyses. As critical thermal limit data were normally distributed, analyses of variance (ANOVA) using the raw data were used to compare critical thermal limits for both field and laboratory studies. Analyses of covariance (ANCOVA) were used to test for differences in rates of water loss and lipid content between populations and seasons (i.e. mass-independent water loss rates), using mean experimental or dry mass as the covariate. Pearson product-moment correlations were used to assess the relationships between time spent active and temperature. Data are presented as means $\pm$ standard error of the mean (S.E.) unless otherwise stated. Significance was set at $P = 0.05$. In no instances for any trait were experimental assessments performed on the same individual on more than one occasion (i.e. animals were not re-used).

Results

Seasonal variation

In the only populations which were sampled on two occasions, season had a significant effect on the majority of the variables examined (Table 2). In the Lambwe and Nguruman populations, CT_{Min} increased and water loss rate declined after the short rains. Although lipid mass also declined in the Lambwe population, the change was not significant in flies from Nguruman. Seasonal changes in body mass and CT_{Max} were opposite in sign in the two populations, though the seasonal difference in CT_{Max} was only 0.4 °C in both populations. Body mass increased in the Lambwe population, but declined at Nguruman, leading to an overall difference in mass between the two populations after the short rains (Table 2; $F_{1, 91} = 20.35$; $P < 0.0001$), but not after the long rains (Table 2; $F_{1, 126} = 0.414$; $P = 0.52$).

Geographic variation

Using data collected within the same season after the short-rains, CT_{Max} and CT_{Min} differed among the four populations (Table 3). The mid-altitude and low-altitude populations differed only marginally, whereas both CT_{Min} and CT_{Max} declined with a further increase in elevation. The slope of the two altitude-critical limit relationships also differed significantly (ANCOVA $F_{1, 234} = 59.42$; $P < 0.0001$), with CT_{Min} showing a steeper slope and stronger relationship

with altitude than CT_{Max} . Overall CT_{Min} varied by 3.3 °C among the four populations, whereas CT_{Max} differed by only 0.4 °C.

Water loss rate also varied significantly among the populations (Fig. 1). It seems unlikely that differences in activity levels among the populations could account for the variation in water loss rate. Total activity time was positively, though in one case weakly, correlated with temperature across the range of 20–32 °C in all four populations (Narok: $r = 0.64$, $P < 0.01$; Lambwe: $r = 0.47$, $P = 0.068$; Nguruman: $r = 0.64$, $P < 0.01$; Kwale: $r = 0.50$, $P < 0.05$). Because there were no sex-related differences in activity time in any of the populations, sexes were pooled for the inter-population comparisons. There were no among-population differences for time spent active (ANCOVA (covariate: temperature): $F_{3, 59} = 1.38$; $P = 0.26$). Therefore, differences in activity probably did not account for differences in water loss rates.

Temperature acclimation of laboratory-bred flies

Ten days of laboratory acclimation at three different temperatures resulted in significant among-treatment variation in CT_{Max} , CT_{Min} , desiccation rate and body water content (Table 4). CT_{Max} increased significantly with treatment temperature, but the change was less than 0.5 °C. By contrast, at 19 °C CT_{Min} declined by *c.* 3 °C relative to the other treatments. Water loss rate showed similar threshold effects, declining significantly at the highest acclimation temperature, whereas body water content showed little change with acclimation temperature (Table 4).

In the time-course experiments, changes in CT_{Min} were induced in the 16 °C and 19 °C treatments only, and the magnitude of the induced change was quite similar (Fig. 2). By contrast, the 21 °C and 24 °C treatments had no effect. In the former treatments, CT_{Min} had reached its nadir by the fifth day of treatment and typically remained at a similar level for the remainder of the acclimation period (day 9). Ten days at 24 °C were sufficient to completely reverse the effects of the acclimation treatments (i.e. at day 19 no groups differed from the 24 °C group). Thus, significant day*treatment interactions were found for CT_{Min} ($F_{15, 216} = 3.99$; $P < 0.0001$). The low treatment temperatures clearly came at some cost to normal adult development. While flies gained mass in the 21 °C and 24 °C treatments, and showed low and typically consistent among-individual variation, the flies at the lower temperatures did not show any steady progression in mass, and variation among individuals was considerable (Fig. 3). The differential effect of treatment on change in mass over the 19 days is evidenced by a significant day*treatment interaction in a two-way analysis of variance examining the effects of day and treatment on mass ($F_{15, 120} = 2.7$, $P = 0.0013$).

Colony vs. field flies

Upper and lower critical thermal limits differed among the laboratory-acclimated (colony) flies and field populations (Fig. 4a, b). However, the extent of this difference was only *c.* 1 °C for CT_{Max} , but from 4 to 10 °C for CT_{Min} . Water loss rates also varied among the field and laboratory populations, and here the maximum difference among groups (Kwale and the 29 °C treatment group) was substantial, although none of the other groups differed from each other (Fig. 4c). By contrast, dry mass varied little among the populations and groups (Fig. 4d).

Discussion

Physiological implications

The temperature and water balance traits differed substantially from each other in the extent of their variability. Of the three traits, CT_{Max} varied least among populations (0.6 °C), seasons (0.4 °C) and acclimation treatments (0.5 °C), and the full range of variation among all of these samples was only 1.1 °C (maximum value 1.02 × larger than the minimum). By contrast,

CT_{Min} was more variable, differing by 3.3 °C, 1–2 °C, and 3.3 °C, among the populations, seasons and acclimation treatments, respectively, and the full range of variation was 9.3 °C (maximum value = 3.1 × minimum). Water loss rate showed similar, substantial variation, with the overall maximum value, across laboratory and field populations, being 3.7 × the minimum. These results are consistent with work across a range of insect taxa. In insects, upper critical temperatures show much less variation than lower critical temperatures among populations and acclimation treatments,³⁸ between selection regimes,³⁹ and among species at a variety of spatial scales.⁴⁰ Substantial differences in water loss rates among populations, among seasons within a population, and between humidity acclimation treatments are also common.^{38,41, 42} However, only a few studies have examined temperature acclimation or acclimatization effects on desiccation resistance, with some finding notable effects of temperature,^{38,43} and others finding no influence.³⁰ Thus, upper thermal critical limits that are not especially variable and water loss rates and lower thermal limits that are more flexible are typical of insects in general. In consequence, their occurrence in *G. pallidipes* is not entirely unexpected.⁴⁴

A further significant finding is that the magnitude of the among-population differences was identical to or smaller than the magnitude of the acclimation effects in the laboratory-reared flies. Phenotypic plasticity might therefore account for all of the among-population variation in *G. pallidipes*. However, because of logistic constraints, each population was not examined to determine the extent of its phenotypic plasticity. Therefore, the extent to which among-population variation really is accounted for by phenotypic plasticity could not be fully determined. Nonetheless, seasonal acclimatization of CT_{Min}, CT_{Max} and water loss rates in the Lambwe and Nguruman populations encompassed 50 – 100% of the among-population variation in these traits. Together with the results from the laboratory acclimation trials, these findings provide substantive evidence for the idea that most of the variation among populations could be accounted for by phenotypic rather than genetic differences. Similar results have been obtained for two independent assessments of *Drosophila melanogaster*.^{38,45} Thus, it seems likely that much of the among-population variation in these traits examined in *G. pallidipes* is a consequence of phenotypic plasticity.

In the case of CT_{Min} these plastic changes take place rapidly. Within five days of exposure to a different temperature, flies held at the lower temperatures had substantially altered their CT_{Min}, and by nine days after their return to a common, higher temperature environment CT_{Min} had returned to its original value. Whether the rapid phenotypic change found in *G. pallidipes* is typical of CT_{Min} or similar traits in other insect species is not clear, largely because the time course of thermal acclimation (rather than developmental plasticity) has not been thoroughly investigated in the group. Nonetheless, several studies have documented similar rapid changes in physiological traits.^{46,47,48} Therefore, reasonably rapid acclimation effects are probably not unusual in insects.

Finally, whilst CT_{Max}, water loss rate and dry mass were quite similar across all of the populations and acclimation treatments, CT_{Min} differed substantially between the field-collected and laboratory-reared flies (Fig. 4). The latter had CT_{Min} that were 3–9 °C lower than those of the field-collected flies. In consequence, it seems likely that differences in the phenotypic plasticity of the upper and lower critical thermal limits extend to the ease with which these traits can evolve. Higher acclimation temperatures in laboratory flies, or lower acclimation temperatures in field flies, might have further altered CT_{Min}. However, this seems unlikely given that more extreme temperatures are debilitating to flies over the longer term.^{7,20} Therefore, the differences between the field and laboratory flies might have evolved as a consequence of conditions in the laboratory. The IAEA laboratory colony used in this study was established in 1975. Assuming linear evolutionary change, approximately 175 generations in the laboratory, and CT_{Min} values of the original Ugandan field population similar to those found at Lambwe, this would equate to approximately 1 °C change every 29 generations (i.e.

6 °C/175 generations). Whilst the reasons for laboratory evolution are not always clear, substantial changes in animals held in colonies have been documented in several studies.^{49, 50} Whatever the reason for the evolved difference, the most noteworthy feature thereof is that this evolution has been restricted to CT_{Min} and has not taken place in the other traits. In consequence, it seems likely that most of the variation in the other traits is a consequence of phenotypic plasticity, whilst the variation found in CT_{Min} is a mixture of plasticity and evolved differences between laboratory and field-collected *G. pallidipes*. In addition, it raises questions about the extent to which the laboratory flies are genetically representative of field-collected flies, at least in the case of this trait. Future work exploring the underlying genetic basis of CT_{Min} would be required to resolve this question.⁵¹

Implications for distributional modelling

Although the critical thermal maxima of adult *G. pallidipes* documented here are higher than the temperatures that induce mortality in the field,²⁰ the present findings nonetheless substantiate correlative work on the determinants of the distribution of this species, and provide several insights into the likely responses of tsetse to climate change. In the first instance, univariate and multivariate analysis of climate and productivity (normalized difference vegetation index, NDVI) variables demonstrated that temperature variables are much better predictors of *G. pallidipes* presence and absence than NDVI variables.^{5,6} These results are in keeping with population level analyses that show a strong relationship between increasing temperature and elevated mortality in *G. pallidipes* adults.²⁰ Our results suggest a causal basis for the determination of *G. pallidipes* distributions by temperature. It appears that in this species tolerance to high temperatures cannot change much either by phenotypic plasticity or by evolution. Such change in tolerance would result in a weaker relationship between temperature and population dynamics, and in doing so would reduce the strength of the relationships between temperature and the presence/absence of the fly species.⁵² The limited variation in CT_{Max} also provides a causal explanation for why, in statistical models of tsetse distribution, the mean temperature difference between areas of fly presence and absence may be less than 1 °C.¹⁹

Limited scope for change in upper critical temperatures might also explain the narrow variation in this trait among populations despite the fact that gene flow between *G. pallidipes* populations is typically low.^{25,34} From a theoretical perspective, limited gene flow between populations should promote diversification so long as gene flow is not so low that novel variants that have successfully survived similar conditions elsewhere are seldom introduced into a population.^{53,54} However, if the trait itself cannot evolve because it is globally constrained,⁵⁵ then changes in gene flow are unlikely to affect trait diversification.

From the present data it might seem difficult to discern why flies are not able to colonize lower temperature areas. However, lower temperature thresholds are likely set by development of the larvae, rather than by adult capabilities.⁵⁶ Moreover, in our acclimation experiments we showed that although flies are able to survive longer-term maintenance at low temperatures, their muscle growth and development (which accounts for the largest change in mass in adult tsetse⁵⁷) is compromised, given that mass declines substantially by days 7–9, and variance in mass between individuals was larger in the lower than in the high temperature treatments (Fig. 3). Therefore, low temperatures are likely to compromise adults too.

To date, most climate change predictions suggest that tsetse species would show a reduction in the range of the flies owing to increases in temperature.¹⁷ Our data support these findings because they show limited plasticity in upper critical temperatures. If upper thermal tolerances are inflexible, then increases in temperature are likely to lead to a reduction in range size as suggested by modelling approaches.⁷ What altered rainfall patterns are likely to mean is more difficult to discern because water loss rates in adult flies respond to altered ambient conditions,

44 water balance effects are likely to be most significant for puparia^{20,56} which were not investigated here, and fine-scale predictions of rainfall change are typically not available.¹ Therefore, it is difficult to be able to predict what the outcome of the interaction between fly desiccation resistance and changing water availability might be. Nonetheless, our results provide further evidence in support of the idea that, owing to changes in land use and climate, at least some tsetse may be brought under autonomous control in parts of their natural ranges.
58

Acknowledgements

We gratefully acknowledge Andrew Parker and the IAEA/FAO Laboratories for contributing laboratory-rearing equipment, puparia and source colony information. The fieldwork in Kenya was supported by KETRI who aided us with staff and logistics. Russ Jurenka performed the lipid extractions at ISU. In Stellenbosch, Henry Davids and Larissa Heyns provided laboratory support. Sue Jackson, Allen Gibbs, Saskia Goldberg and Brent Sinclair are thanked for comments and discussion. Two anonymous referees are thanked for their comments. This work was funded by NIH grant AI-52456 to E. S. Krafur.

References

1. Watson RT, 2002. *Climate change 2001: Synthesis report*. Cambridge: Cambridge University Press.
2. Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont LJ, Collingham YC, Erasmus BFN, de Siqueira MF, Grainger A, Hannah L, Hughes L, Huntley B, van Jaarsveld AS, Midgley GF, Miles L, Ortega-Huerta MA, Peterson AT, Phillips OL, Williams SE. Extinction risk from climate change. *Nature* 2004;427:145–148. [PubMed: 14712274]
3. Pearson RG, Dawson TP. Predicting the impacts of climate change on the distribution of species: are bioclimate envelope models useful? *Global Ecol Biogeog* 2003;12:361–371.
4. Rogers DJ, Randolph SE. Mortality rates and population density of tsetse flies correlated with satellite imagery. *Nature* 1991;351:739–741. [PubMed: 2062367]
5. Robinson T, Rogers D, Williams B. Univariate analysis of tsetse habitat in the common fly belt of Southern Africa using climate and remotely sensed vegetation data. *Med Vet Entomol* 1997;11:223–234. [PubMed: 9330253]
6. Robinson T, Rogers D, Williams B. Mapping tsetse habitat suitability in the common fly belt of Southern Africa using multivariate analysis of climate and remotely sensed vegetation data. *Med Vet Entomol* 1997;11:235–245. [PubMed: 9330254]
7. Rogers DJ. Satellites, space, time, and the African trypanosomiasis. *Adv Parasitol* 2000;47:129–171. [PubMed: 10997206]
8. Davis AJ, Jenkinson LS, Lawton JH, Shorrocks B, Wood S. Making mistakes when predicting shifts in species range in response to global warming. *Nature* 1998;391:783–786. [PubMed: 9486646]
9. Davis MB, Shaw RG. Range shifts and adaptive responses to Quaternary climate change. *Science* 2001;292:673–679. [PubMed: 11326089]
10. Helmuth B, Kingsolver JG, Carrington E. Biophysics, physiological ecology, and climate change: does mechanism matter? *Ann Rev Physiol* 2005;67:177–201. [PubMed: 15709956]
11. Rogers DJ, Randolph SE. The global spread of malaria in a future, warmer world. *Science* 2000;289:1763–1766. [PubMed: 10976072]
12. Kovats RS, Campbell-Lendrum DH, McMichael AJ, Woodward A, Cox J, St H. Early effects of climate change: do they include changes in vector-borne disease? *Phil Trans R Soc Lond B* 2001;356:1057–1068. [PubMed: 11516383]
13. Burse BV, Mellor PS, Rogers DJ, Samuel AR, Mertens PPC, Baylis M. Climate change and the recent emergence of bluetongue in Europe. *Nature Rev Microbiol* 2005;3:171–181. [PubMed: 15685226]
14. Rogers DJ, Randolph SE, Snow RW, Hay SI. Satellite imagery in the study and forecast of malaria. *Nature* 2002;415:710–715. [PubMed: 11832960]
15. Leak SGA, 1999. *Tsetse biology and ecology*. Wallingford: CABI Publishing.
16. Watson RT, Zinyowera MC, Moss RH, Dokken DJ, 1998. *The regional impacts of climate change. An assessment of vulnerability*. Cambridge: Cambridge University Press.

17. Hulme M, 1996. Climate change in Southern Africa: An exploration of some potential impacts and implications in the SADC region. Climatic Research Unit, Norwich: University of East Anglia.
18. World Health Organization, 2004. *Using climate to predict disease outbreaks*. Geneva, World Health Organization.
19. Rogers DJ, Robinson TP, 2004. Tsetse distribution. Maudlin I, Holmes PH, Miles MA, eds. *The Trypanosomiases*. Wallingford: CABI Publishing, 139–179.
20. Hargrove JW, 2004. Tsetse population dynamics. Maudlin I, Holmes PH, Miles MA, eds. *The Trypanosomiases*. Wallingford: CABI Publishing, 113–138.
21. Hargrove JW. The effect of temperature and saturation deficit on mortality in populations of male *Glossina m. morsitans* (Diptera : Glossinidae) in Zimbabwe and Tanzania. *Bull Ent Res* 2001;91:79–86.
22. Bursell E. The water balance of tsetse flies. *Trans R Ent Soc Lond* 1959;111:205–235.
23. Langley PA, Maudlin I, Leedham MP. Genetic and behavioural differences between *Glossina pallidipes* from Uganda and Zimbabwe. *Entomol Exp Appl* 1984;35:55–60.
24. Krafur ES. Tsetse fly population genetics: an indirect approach to dispersal. *Trends Parasitol* 2003;19:162–166. [PubMed: 12689645]
25. Ouma JO, Marquez JG, Krafur ES. Macrogeographic population structure of the tsetse fly, *Glossina pallidipes* (Diptera: Glossinidae). *Bull Ent Res* 2005;95:437–447.
26. Williams B, Dransfield R, Brightwell R. The control of tsetse flies in relation to fly movement and trapping efficiency. *J Appl Ecol* 1992;29:163–179.
27. Lutterschmidt WI, Hutchison VH. The critical thermal maximum: history and critique. *Canadian Journal of Zoology* 1997;75:1561–1574.
28. Klok CJ, Sinclair BJ, Chown SL. Upper thermal tolerance and oxygen limitation in terrestrial arthropods. *J Exp Biol* 2004;207:2361–2370. [PubMed: 15159440]
29. Chown SL, Nicolson SW, 2004. *Insect physiological ecology. Mechanisms and patterns*. Oxford: Oxford University Press.
30. Terblanche JS, Sinclair BJ, Klok CJ, McFarlane ML, Chown SL. The effects of acclimation on thermal tolerance, desiccation resistance and metabolic rate in *Chirodica chalcoptera* (Coleoptera: Chrysomelidae). *J Ins Physiol* 2005;51:1013–1023.
31. Hoffmann AA, Hallas R, Anderson AR, Telonis-Scott M. Evidence for a robust sex-specific trade-off between cold resistance and starvation resistance in *Drosophila melanogaster*. *J Evol Biol* 2005;18:804–810. [PubMed: 16033551]
32. Gibbs AG, Chippindale AK, Rose MR. Physiological mechanisms of evolved desiccation resistance in *Drosophila melanogaster*. *J Exp Biol* 1997;200:1821–1832. [PubMed: 9225453]
33. Brady J. Circadian ontogeny in the tsetse fly: A permanent major phase change after the first feed. *J Ins Physiol* 1988;34:743–749.
34. Krafur ES, Wohlford DL. Breeding structure of *Glossina pallidipes* populations evaluated by mitochondrial variation. *J Hered* 1999;90:635–642. [PubMed: 10589514]
35. Terblanche JS, Klok CJ, Chown SL. Metabolic rate variation in *Glossina pallidipes* (Diptera: Glossinidae): gender, ageing and repeatability. *J Ins Physiol* 2004;50:419–428.
36. Gooding RH, Feldmann U, Robinson AS, 1997. Care and maintenance of tsetse colonies. Crampton JM, Beard CB, Louis C, eds. *The molecular biology of insect disease vectors*. London: Chapman & Hall, 41–55.
37. Sombroek WG, Braun HMM, van der Pouw BJA, 1980. *Exploratory soil map and agroclimatic zone map of Kenya*. Kenya Soil Survey, Ministry of Agriculture, Nairobi: National Agricultural Laboratories.
38. Hoffmann AA, Shirriffs J, Scott M. Relative importance of plastic vs genetic factors in adaptive differentiation: geographical variation for stress resistance in *Drosophila melanogaster* from eastern Australia. *Funct Ecol* 2005;19:222–227.
39. Gilchrist GW, Huey RB, Partridge L. Thermal sensitivity of *Drosophila melanogaster*: evolutionary responses of adults and eggs to laboratory natural selection at different temperatures. *Physiol Zool* 1997;70:403–414. [PubMed: 9237300]

40. Chown SL. Physiological variation in insects: hierarchical levels and implications. *J Ins Physiol* 2001;47:649–660.
41. Addo-Bediako A, Chown SL, Gaston KJ. Revisiting water loss in insects: a large scale view. *J Ins Physiol* 2001;47:1377–1388.
42. Gibbs AG, Matzkin LM. Evolution of water balance in the genus *Drosophila*. *J Exp Biol* 2001;204:2331–2338. [PubMed: 11507115]
43. Parkash R, Tyagi P, Sharma I, Rajpurohit S, 2005. Adaptations to environmental stress in altitudinal populations of two *Drosophila* species. *Physiol Entomol* doi:10.1111/j.1365–3032.2005.00470.x.
44. Rogers DJ. A general model for tsetse populations. *Ins Sci Appl* 1990;11:331–346.
45. Ayrihac A, Debat V, Gibert P, Kister AG, Legout H, Moreteau B, Vergilino R, David JR. Cold adaptation in geographical populations of *Drosophila melanogaster*: phenotypic plasticity is more important than genetic variability. *Funct Ecol* 2004;18:700–706.
46. Zeilstra I, Fischer K. Cold tolerance in relation to developmental and adult temperature in a butterfly. *Physiol Entomol* 2005;30:92–95.
47. Fischer K, Eenhoorn E, Bot ANM, Brakefield PM, Zwaan BJ. Cooler butterflies lay larger eggs: developmental plasticity vs acclimation. *Proc R Soc Lond Biol Sci* 2003;270:2051–2056.
48. Gibert P, Huey RB, Gilchrist GW. Locomotor performance of *Drosophila melanogaster*: interactions among developmental and adult temperatures, age, and geography. *Evolution* 2001;55:205–209. [PubMed: 11263741]
49. Sgrò CM, Partridge L. Evolutionary responses of the life history of wild-caught *Drosophila melanogaster* to two standard methods of laboratory culture. *Am Nat* 2000;156:341–353.
50. Terblanche JS, Klok CJ, Marais E, Chown SL. Metabolic rate in the whip-spider, *Damon annulatipes* (Arachnida: Amblypygi). *J Ins Physiol* 2004b;50:637–645.
51. Loeschcke V, Sorensen JG, Kristensen TN. Ecologically relevant stress resistance: from microarrays and quantitative trait loci to candidate genes - A research plan and preliminary results using *Drosophila* as a model organism and climatic and genetic stress as model stresses. *J Biosci* 2004;29:503–511. [PubMed: 15625405]
52. Hoffmann AA, Blows MW. Species borders: ecological and evolutionary perspectives. *Trends Ecol Evol* 1994;9:223–227.
53. Case TJ, Taper ML. Interspecific competition, environmental gradients, gene flow, and the coevolution of species' borders. *Am Nat* 2000;155:583–605. [PubMed: 10777432]
54. Forde SE, Thompson JN, Bohannan BJM. Adaptation varies through space and time in a coevolving host-parasitoid interaction. *Nature* 2004;431:841–844. [PubMed: 15483611]
55. Hoffmann AA, Hallas RJ, Dean JA, Schiffer M. Low potential for climatic stress adaptation in a rainforest *Drosophila* species. *Science* 2003;301:100–102. [PubMed: 12843394]
56. Langley PA. Physiology of tsetse flies (*Glossina* ssp.)(Diptera: Glossinidae): a review. *Bull Ent Res* 1977;67:523–574.
57. Bursell E. Post-teneral development of the thoracic musculature in tsetse flies. *Proc R Ent Soc Lond A* 1961;36:69–74.
58. Rogers DJ, Randolph SE. A response to the aim of eradicating tsetse from Africa. *Trends Parasitol* 2002;18:534–536. [PubMed: 12482537]

Fig. 1.

Geographic variation of water loss rate in *Glossina pallidipes*. Note that desiccation rate is highest in the low altitude population from Kwale (388 m). All flies were sampled in the same season (Nov-Dec, 2003). (Narok: 1691 m; Lambwe: 1353 m; Nguruman: 670 m.)

Fig. 2.

The time-course of acclimation for CT_{Min} in *Glossina pallidipes* adults. After nine days of acclimation to 16, 19, 21 and 24 °C, all acclimation groups were returned to 24 °C for ten days recovery (tested on Day 19). Within acclimation treatment (between-day) comparisons of the means are presented in Table 5. Between acclimation treatments (within-day) comparisons are presented in Table 6. (Recorded climate chamber temperatures (mean $\pm$ S.D.): 16: 15.7 $\pm$ 0.37; 19: 18.6 $\pm$ 0.22; 21: 20.6 $\pm$ 0.50; 24: 24.4 $\pm$ 1.38 °C.) N = 10 flies for all days and all acclimation temperature groups. There was no effect of acclimation on 21 and 24 °C acclimation groups, but both 16 and 19 °C had significantly reduced CT_{Min} by Day 5 (ANOVA: 16 °C: $F_{5, 54} = 8.70$; $P < 0.0001$; 19 °C: $F_{5, 54} = 9.31$; $P < 0.0001$; 21 °C: $F_{5, 54} = 1.99$; $P = 0.095$; 24 °C: $F_{5, 54} = 1.38$; $P = 0.25$). By Day 19 all acclimation groups were similar ($F_{3, 36} = 2.64$; $P = 0.064$).

Fig. 3A-D.

Body mass (mean $\pm$ S.E.) of virgin female *Glossina pallidipes* during the course of acclimation to A) 24, B) 21, C) 19 and D) 16 °C. All flies were returned to 25 °C after day nine of acclimation. All mass measurements were obtained on non-feeding days, i.e. 24 h post-feeding. Although no differences were found in body mass upon completion of the acclimation treatments, there was a significant day * treatment interaction (see text for statistics).

Fig. 4A-D.

Traits means ($\pm$ S.E.), in field and laboratory populations of *Glossina pallidipes*, of **A**) critical thermal maxima (ANOVA, $F_{6,161} = 22.38$; $P < 0.0001$); **B**) critical thermal minima (ANOVA, $F_{6,195} = 453.20$; $P < 0.0001$); **C**) water loss rate (ANCOVA, covariate: dry body mass; $F_{6,126} = 12.378$; $P < 0.0001$; least squares adjusted means $\pm$ 95% confidence limits; **D**) body size as dry body mass from all desiccation experiment flies (ANOVA; $F_{6,239} = 8.43$; $P < 0.0001$; means $\pm$ 95% confidence limits). (The numerals in Lab19, Lab24 and Lab29 refer to the temperature of acclimation in laboratory-reared flies for periods of 9-10 days).

Table 1

Field sampling sites (coordinates in decimal degrees) and time of sampling of *Glossina pallidipes* for this study (both seasons were sampled during 2003).

Region	Coordinates (°)	Altitude (m)	Sampling season
Narok	1.12S, 35.20E	1691	After short rains
Lambwe	0.64S, 34.31E	1353	After long and short rains
Nguruman	1.85S, 26.10E	670	After long and short rains
Kwale	4.18S, 39.46E	388	After short rains

Table 2

Summary statistics for seasonal variation of physiological traits in *Glossina pallidipes* from the Lambwe and Nguruman populations. The results of either ANOVA or ANCOVA (with dry mass as covariate, indicated with superscript A) are shown.

Population	Trait	Season		F-ratio	DF	P
		Long rains	Short rains			
Lambwe	CT _{Max} (°C)	45.0 ± 0.1	44.6 ± 0.1	11.18	58	< 0.01
	CT _{Min} (°C)	12.3 ± 0.2	13.4 ± 0.3	9.83	58	< 0.01
	Water loss rate (mg H ₂ O.h ⁻¹) ^A	0.620 ± 0.027	0.536 ± 0.032	6.95	51	< 0.05
	Body (dry) mass (mg)	11.74 ± 0.42	13.73 ± 0.73	6.38	59	< 0.05
	Body lipid content (mg) ^A	2.00 ± 0.07	1.89 ± 0.08	30.03	106	< 0.0001
Nguruman	CT _{Max} (°C)	44.6 ± 0.1	45.0 ± 0.1	12.00	56	< 0.01
	CT _{Min} (°C)	11.8 ± 0.2	13.8 ± 0.2	74.79	58	< 0.0001
	Water loss rate (mg H ₂ O.h ⁻¹) ^A	0.789 ± 0.042	0.474 ± 0.033	40.15	54	< 0.0001
	Body (dry) mass (mg)	11.62 ± 0.39	10.84 ± 0.41	7.13	47	< 0.05
	Body lipid content (mg) ^A	1.53 ± 0.07	0.99 ± 0.08	3.308	98	0.072

Table 3

Geographic variation in mean population critical thermal maxima (CT_{Max}) and minima (CT_{Min}) for *Glossina pallidipes*. All flies were collected in the same season (Nov-Dec, 2003). (ANOVA: CT_{Max} : $F_{3, 114} = 14.0$; $P < 0.0001$; CT_{Min} : $F_{3, 116} = 70.35$; $P < 0.0001$.)

Trait	Mean ° C $\pm$ S.E.	Min	Max	N
CT_{Max}				
Narok	44.4 $\pm$ 0.1	43.5	45.1	30
Lambwe	44.6 $\pm$ 0.1	43.7	45.2	30
Nguruman	45.0 $\pm$ 0.1	44.3	45.7	28
Kwale	45.0 $\pm$ 0.1	44.2	45.5	30
CT_{Min}				
Narok	10.5 $\pm$ 0.2	8.9	12.4	30
Lambwe	13.4 $\pm$ 0.3	11.8	16.0	30
Nguruman	13.8 $\pm$ 0.2	11.8	15.2	30
Kwale	13.8 $\pm$ 0.2	12.6	15.9	30

Upper (CT_{Max}) and lower (CT_{Min}) critical thermal limits, water loss rate at 24 °C and initial body water content after ten days of temperature acclimation in laboratory-reared *Glossina pallidipes* adults. Recorded climate chamber temperatures presented in parentheses (mean ± S.D.) following treatment temperature. Similar superscript letters denote statistically homogeneous groups (*Post-hoc* unequal sample HSD). All analyses by means of ANOVA unless otherwise specified.

Table 4

Acclimation Temperature	CT _{Max} (°C)	CT _{Min} (°C)	Water loss rate ^I (mg H ₂ O.hr ⁻¹)	Body water content (mg H ₂ O)
19 °C (19.5 ± 0.4)	43.9 ± 0.1 ^A (20)	4.5 ± 0.1 ^A (20)	0.464 ± 0.061 ^A (8)	29.9 ± 0.7 ^A (8)
24 °C (23.4 ± 0.9)	44.4 ± 0.1 ^B (10)	7.8 ± 0.3 ^B (12)	0.393 ± 0.017 ^A (9)	25.2 ± 2.8 ^B (9)
29 °C (28.8 ± 1.7)	44.2 ± 0.1 ^{A,B} (20)	7.2 ± 0.2 ^B (20)	0.213 ± 0.047 ^B (8)	23.4 ± 1.5 ^B (8)
F	5.10	97.34	7.89	2.77
P	<0.01	<0.001	<0.003	0.085

^I ANCOVA with dry body mass as covariate