Journal of Zhejiang University SCIENCE B
 ISSN 1673-1581 (Print); ISSN 1862-1783 (Online)
 www.zju.edu.cn/jzus; www.springerlink.com
 E-mail: jzus@zju.edu.cn

Guava extract (*Psidium guajava*) alters the labelling of blood constituents with technetium-99m^{*}

ABREU P.R.C.¹, ALMEIDA M.C.¹, BERNARDO R.M.¹, BERNARDO L.C.¹,
 BRITO L.C.¹, GARCIA E.A.C.², FONSECA A.S.^{†3}, BERNARDO-FILHO M.¹

(¹Department of Biophysics and Biometry, Institute of Biology Roberto Alcântara Gomes,
 University of Estate of Rio de Janeiro, Rio de Janeiro 20551030, Brazil)

(²Department of Physiology, Biological Sciences and Health Center, Federal University of Sergipe, São Cristóvão, Sergipe 49100000, Brazil)

(³Department of Pharmacology and Psychobiology, Institute of Biology Roberto Alcântara Gomes,
 University of Estate of Rio de Janeiro, Rio de Janeiro 20551030, Brazil)

[†]E-mail: adenilso@uerj.br

Received Dec. 8, 2005; revision accepted Mar. 31, 2006

Abstract: *Psidium guajava* (guava) leaf is a phytotherapeutic used in folk medicine to treat gastrointestinal and respiratory disturbances and is used as anti-inflammatory medicine. In nuclear medicine, blood constituents (BC) are labelled with technetium-99m (^{99m}Tc) and used to image procedures. However, data have demonstrated that synthetic or natural drugs could modify the labelling of BC with ^{99m}Tc. The aim of this work was to evaluate the effects of aqueous extract of guava leaves on the labelling of BC with ^{99m}Tc. Blood samples of *Wistar* rats were incubated with different concentrations of guava extract and labelled with ^{99m}Tc after the percentage of incorporated radioactivity (%ATI) in BC was determined. The results suggest that aqueous guava extract could present antioxidant action and/or alters the membrane structures involved in ion transport into cells, thus decreasing the radiolabelling of BC with ^{99m}Tc. The data showed significant ($P<0.05$) alteration of ATI in BC from blood incubated with guava extract.

Key words: Antioxidant, Blood constituents, *Psidium guajava*, Technetium-99m

doi:10.1631/jzus.2006.B0429

Document code: A

CLC number: R282.71

INTRODUCTION

Psidium guajava (guava), belonging to the family of *Myrtaceae*, is a native of tropical America and has long been naturalized in Southeast Asia. The positive effects of guava extracts on human ailments have been described (Lozoya, 1999).

The pharmacological actions and the medicinal uses of aqueous extracts of guava leaves in folk medicine include the treatment of various types gastrointestinal disturbances such as vomiting, diarrhea, inhibition of the peristaltic reflex, gastroenteritis, spasmolytic activity, dysentery, abdominal distention, flatulence and gastric pain (Lutterodt, 1992; Aguilar

et al., 1994; Lozoya et al., 1994). These extracts have also been indicated to cause disturbances of the central nervous system: insomnia, convulsions and epilepsy (Lutterodt and Maleque, 1988; Meckes et al., 1996). Bronchitis, asthma attacks, cough, pulmonary diseases could be also treated with guava teas (Batick, 1984; Khan and Ahmad, 1985) and could also be useful as anti-inflammatory and hemostatic agent (Liu, 1988).

Moreover, aqueous extracts of guava leaves were described to be effective against a number of microbial strains: *Aeromonas hydrophila*, *Shigella* spp. and *Vibrio* spp. (Chulasiri et al., 1986), *Staphylococcus aureus* and β -*streptococcus* group A (Jaiarj et al., 1999), *Escherichia coli*, *Pseudomonas aeruginosa*, *Bacillus subtilis* (Abdelrahim et al., 2002). In addition,

^{*} Project supported by CAPES, CNPq and FAPERJ, Brazil

anti-rotavirus activity has also been reported to exist in these extracts (Gonçalves *et al.*, 2005).

Phytochemical studies have identified more than 20 compounds in guava extracts (Osman *et al.*, 1974; Begum *et al.*, 2002). The major constituents of its leaves were identified to be tannins, β -sitosterol, maslinic acid, essential oils, triterpenoids and flavonoids (Osman *et al.*, 1974; Arima and Danno, 2002; Begum *et al.*, 2002; 2004).

Technetium-99m (^{99m}Tc) has been the most used radionuclide in nuclear medicine diagnosis procedures (single photon emission computed tomography, SPECT) and in labelling of molecular and cellular structures that can be used as radiopharmaceuticals (radiobiocomplexes) (Saha, 2003).

Red blood cells labelled with ^{99m}Tc (^{99m}Tc -RBC) are radiobiocomplexes widely used in clinical nuclear medicine for several important applications (Wong *et al.*, 2004; Zaman *et al.*, 2004; Olds *et al.*, 2005; Harel *et al.*, 2005; Artiko *et al.*, 2004; Verdu *et al.*, 2005; Cortes *et al.*, 2003; Jin *et al.*, 2004; Slart *et al.*, 2004).

Sequential steps of the intracellular labelling process of blood constituents include: (1) transmembrane transport of reducing agent (Sn^{2+}) and $^{99m}\text{TcO}_4^-$ ions into the internal compartment of the RBC, (2) reduction of ^{99m}Tc ($^{99m}\text{TcO}_4^-$) by Sn^{2+} , and (3) subsequent binding of the reduced ^{99m}Tc to hemoglobin (Dewanjee *et al.*, 1982). The band-3 anion transport system and calcium channels may be involved in transport of these ions (Callahan and Rabito, 1990; Gutfilen *et al.*, 1992; Sampson, 1996).

Several authors have reported the effect of artificial and natural drug on the radiolabelling process (de Oliveira *et al.*, 2003a; 2003b; Frydman *et al.*, 2004; Moreno *et al.*, 2004; Fonseca *et al.*, 2005). The drugs could alter the labelling of blood constituents acting as antioxidant agent, modify the membrane structure or decrease the efficiency of transmembrane transport system of stannous and pertechnetate ions into cells. In this context, compounds in aqueous extracts of guava leaves have been reported to present antioxidant action and some of their pharmacological effects could be related to interaction in calcium channels (Morales *et al.*, 1994; Qian and Nihorimbere, 2004). However, to the authors' knowledge, there is no report on the possible effects of guava extracts on radiolabelling of blood constituents as RBC or

plasma proteins.

This work was aimed at evaluating the effects of aqueous extract of guava leaves on the labelling of blood constituents with technetium-99m.

MATERIALS AND METHODS

Animals

Adult male *Wistar* rats (3~4 months of age, body weight 250~350 g) were maintained in a controlled environment and allowed free access to water and food and ambient temperature was kept at $(25 \pm 2)^\circ\text{C}$. Experiments were conducted in accordance with the rules of the Institutional Committee of Animal Care.

Preparation of guava extract

The guava extracts were prepared with leaves of *Psidium guajava* L. (Conde Garcia *et al.*, 2003) obtained from the Campus of the Federal University of Sergipe (Aracaju, Brazil), and collected between June and July of 2000 (plants were free of toxic compounds). A botanic specimen was compared with the exsiccata at Federal University of Sergipe Herbarium, Brazil (code ASE 03304, collectors T.S. dos Santos and V.S. Paes, collectors No. 4122, registering book No. 1, page 138, exsiccata No. 04531).

To prepare the extracts, 20 mg of dry leaves were ground in 2 ml of 0.9% NaCl (100°C , 5 min). The crude extract was filtered, centrifuged (1500 r/min, 5 min) to obtain the final extract. The supernatant was considered as the concentration at 10 mg/ml of guava leaves.

Radiolabelling in vitro of blood constituents

Heparinized blood (500 μl), was withdrawn from the *Wistar* rats and incubated with 100 μl of guava extract at different concentrations (6.25, 12.5, 25.0, 50.0 and 100.0 mg/ml) or with a saline solution alone, as control, for 1 h (room temperature). After this the radiolabelling of blood constituents with ^{99m}Tc ($\text{Na}^{99m}\text{TcO}_4$, Institute of Energetic and Nuclear Research, National Commission of Nuclear Energy, São Paulo, Brazil) was performed as described in (Bernardo-Filho *et al.*, 1983). The samples were centrifuged (1500 r/min, 5 min) and aliquot of plasma (P) and blood cells (BC) were isolated. Another aliquots of P and BC were separated and pre-

cipitated in trichloroacetic acid (5%) and centrifuged (1500 r/min, 5 min) to isolate soluble (SF) and insoluble fractions (IF). The radioactivity in P, BC, SF-P, IF-P, SF-BC and IF-BC were determined in a well counter (Packard, model C5002, Illinois, USA) and the percentage of incorporated radioactivity (%ATI) was calculated as described in (Bernardo-Filho *et al.*, 1983; 1986).

Statistical analysis

Data obtained are reported as means±standard deviation of %ATI. To compare the treated (5 blood samples for each extract concentration) and control groups (10 blood samples), the one way analysis of variance (ANOVA) test was performed to verify possible statistical difference between the experimental groups. After this, a rigorous statistical post test (Tukey) was chosen to identify the *P* value ($P<0.05$ as lesser significant level) and compare each treated group with control. InStat GraphPad software was used to perform statistical analysis (GraphPad InStat version 3.00 for Windows 95, GraphPad Software, San Diego California, USA).

RESULTS

Fig.1a shows the %ATI in insoluble (IF-P) and soluble (SF-P) fractions isolated from plasma separated from whole blood treated with different concentrations of guava extract. Analysis of these data indicates that guava extract significantly ($P<0.05$) reduced the radioactivity uptake in IF-P at the highest concentration studied (100.0 mg/ml) but no alteration was obtained with lesser concentration used.

Fig.1b shows the %ATI in insoluble (IF-BC) and soluble (SF-BC) fractions isolated from blood cells separated from blood treated with different concentrations of guava extract. Analysis of these data indicates that incubation with guava extract at low concentrations (6.0 and 12.5 mg/ml) did not reduce the %ATI of IF-BC. However, when the highest concentrations (25, 50 and 100 mg/ml) were used significant alterations of radioactivity uptake in insoluble fractions of blood cells resulted.

Fig.1c shows the %ATI in blood cells (BC) and plasma (P) from blood treated with different concentrations of guava extract. Analysis of these data indicated that guava extract alters the distribution of

Fig.1 Effect of guava extract on (a) uptake of ^{99m}Tc by insoluble (IF-P) and soluble (SF-P) fractions of plasma (P), (b) uptake of ^{99m}Tc by insoluble (IF-BC) and soluble (SF-BC) fractions of blood cells (BC), and (c) the distribution of the ^{99m}Tc in the plasma and blood cells (BC) compartments, in the radiolabeling procedure of blood elements. Heparinized blood samples of *Wistar* rats were incubated with different concentrations of guava extract (1 h) and after with SnCl_2 (1.20 $\mu\text{g/ml}$, 1 h) and in sequence with $\text{Na}^{99m}\text{TcO}_4$ (3.7 MBq, 10 min). (a) Insoluble and soluble fractions of plasma (IF-P and SF-P) and (b) Insoluble and soluble fractions of blood cells (IF-BC and SF-BC) were obtained by precipitation with trichloroacetic acid (5%) and centrifugation (1500 r/min, 5 min). The radioactivity in these fractions was counted and the %ATI was calculated; (c) After centrifugation, plasma (P) and blood cells (BC) were isolated, the radioactivity was counted and the %ATI calculated

$P<0.05$; ## $P<0.01$; ### $P<0.001$, when compared to control group of plasma; * $P<0.05$; ** $P<0.01$; *** $P<0.001$, when compared to control group of blood cells

radioactivity in these two compartments (BC and P) at all guava extract concentrations studied. The data in this figure present an inverse dose-dependent relation with greater decreasing of cells %ATI at highest concentrations of guava extract.

DISCUSSION

Analysis of data presented in this study showed that aqueous guava extract could modify the fixation of ^{99m}Tc in plasma proteins only at high concentrations (Fig.1a). In cellular proteins, guava extract could affect the uptake of ^{99m}Tc at high but also at lesser concentrations (Fig.1b). However, the effects of this extract were more evident on the distribution of ^{99m}Tc between the blood compartments. The data analysis results suggest that aqueous extract of guava leaves used in this work could decrease the fixation of radioactivity in blood cell in a typical dose-dependent relation (Fig.1c).

Some authors have described that natural and synthetic products can alter the labelling of blood constituents with ^{99m}Tc (Frydman *et al.*, 2004; Valenca *et al.*, 2005; Fonseca *et al.*, 2005). The labelling of blood constituents could decrease due to the action of drugs in (1) binding at the same sites on the blood constituents, (2) direct inhibition (chelating action) of the stannous and pertechnetate ions, (3) direct oxidation or generation of free radicals that could oxidize the stannous ion, (4) antioxidant action impeding or decreasing the stannous ion oxidation, and (5) alteration of the plasma membrane structure or modifying the transport systems of stannous and pertechnetate ions into cells.

Several compounds (as flavonoids) present in guava extracts are transported in blood attached to plasma proteins (Podhajcer *et al.*, 1980). The fixation of ^{99m}Tc in plasma proteins occurs at different proteins sites with albumin being the principal protein involved (Early and Soddee, 1995). The binding sites in these proteins could be the same binding sites of ^{99m}Tc and thus in the presence of guava extract compounds the plasma protein labelling with ^{99m}Tc could be reduced. However, the data shown in this work suggest that at only highest concentration of aqueous guava extract these compounds (flavonoids and others phenolic compounds) could compete with

the binding of ^{99m}Tc with plasma proteins.

The antioxidant activity of some compounds could be used to prevent various chronic diseases such as heart-disease, diabetes, cancer, arterial thrombosis, cataract and may provide health-promoting effects (Kimura *et al.*, 1985; Qian and Nihorimbere, 2004). Chemical analysis of guava extracts has revealed the presence of essential oils, tannins, saponins, carotenoids, flavonoids and triterpenes (Cuellar *et al.*, 1984; Mercadante *et al.*, 1999; Arima and Danno, 2002; Begum *et al.*, 2002). The different phenolic phytochemicals in guava leaves have been described to possess antioxidant action and able to inhibit peroxidation reaction in living systems (Kimura *et al.*, 1985; Vanacker *et al.*, 1995; Qian and Nihorimbere, 2004). Thus, some compounds present in guava extracts can inhibit the oxidation of the stannous ions and to alter the labelling of plasma and cellular proteins with ^{99m}Tc as well as interfere with the distribution of this radionuclide between plasma and cellular compartments.

Moreover, data have demonstrated that aqueous and hidroalcoholic extracts of guava leaves present anti-microbial activity (Arima and Danno, 2002; Qa'dan *et al.*, 2005). Flavonoid glycosides and flavonoids as quercetin and quercetin have been isolated and present activity against *Salmonella enteritidis* and *Bacillus cereus* (Arima and Danno, 2002). This anti-microbial activity may be related to action on plasma membrane modifying its structure and decreasing or impeding the transport of stannous and pertechnetate ions into blood cells that can alter the labelling efficiency of blood constituents with ^{99m}Tc . On the other hand, the above cited compound quercetin has been described to decrease the spasmolytic activity in experimental models (Morales *et al.*, 1994). This effect was related to a possible narcotic (opioid) component in the guava leaves (Lutterodt and Maleque, 1988; Lutterodt, 1992) while other data have indicated inhibition of acetylcholine release in neuromuscular junctions by calcium-antagonistic action of quercetin decreasing the inward calcium membrane current leading to a decrease of smooth muscle contractile force (Lozoya *et al.*, 1994; Morales *et al.*, 1994; Re *et al.*, 1999) and depression of myocardial inotropism (Conde Garcia *et al.*, 2003). The radiolabelling process of red blood cells with ^{99m}Tc depends on the entry of stannous and pertechn-

netate ions into these cells through ionic channels (Callahan and Rabito, 1990; Gutfilen *et al.*, 1992). The action of quercetin and/or other compounds present in aqueous guava extract blocking and/or binding on calcium and anions (as band 3) channels could make difficult the transport of stannous ions through plasma membrane and to decrease the labelling of blood cells with ^{99m}Tc explaining in part the data obtained in this work.

In conclusion, our data suggest that aqueous extracts of guava leaves could present antioxidant action and/or effects on the membrane structures involved in ions transport altering the radiolabelling of blood constituents with ^{99m}Tc and that precaution should be taken in examinations of nuclear medicine based on this procedure in patients using guava extracts.

References

- Abdelrahim, S.I., Almadboul, A.Z., Omer, M.E.A., Elegami, A., 2002. Antimicrobial activity of *Psidium guajava* L. *Fitoterapia*, **73**(7-8):713-715. [doi:10.1016/S0367-326X(02)00243-5]
- Aguilar, A., Argueta, A., Cano, L., 1994. Flora Medicinal Indígena de México. Treinta Y Cinco Monografías del Atlas de las Plantas de la Medicina Tradicional Mexicana. INI, México, p.245 (in Spanish).
- Arima, H., Danno, G., 2002. Isolation of antimicrobial compounds from guava (*Psidium guajava* L.) and their structural elucidation. *Bioscience Biotechnology and Biochemistry*, **66**(8):1727-1730. [doi:10.1271/bbb.66.1727]
- Artiko, V., Obradovic, V., Petrovic, M., Perisic-Savic, M., Suvajdzic, N., 2004. Hemangioma of the spleen confirmed by blood pool scintigraphy. *Clinical Nuclear Medicine*, **29**:670-671. [doi:10.1097/00003072-200410000-00023]
- Batick, M.J., 1984. Ethnobotany of Palms in the Neotropics. In: Prance, G.T., Kallunki, J.A. (Eds.), *Advances in Economic Botany: Ethnobotany in the Neotropics*. New York Botanical Garden, New York, USA, p.9-23.
- Begum, S., Hassan, S.I., Siddiqui, B.S., Shaheen, F., Ghayur, M.N., Gilani, A.H., 2002. Triterpenoids from the leaves of *Psidium guajava*. *Phytochemistry*, **61**(4):399-403. [doi:10.1016/S0031-9422(02)00190-5]
- Begum, S., Hassan, S.I., Ali, S.N., Siddiqui, B.S., 2004. Chemical constituents from the leaves of *Psidium guajava*. *Natural Product Research*, **18**(2):135-140. [doi:10.1080/14786410310001608019]
- Bernardo-Filho, M., Moura, I.N.S., Boasquevisque, E.M., 1983. ^{99m}Tc -labelled red blood cells "in vitro". *Arquivos de Biological Tecnologia*, **26**:455-461.
- Bernardo-Filho, M., Caniné, M.S., Lopes, R.L.F., Boasquevisque, E.M., 1986. Effect of temperature on the "in vitro" labelling of red blood cells with technetium- ^{99m}Tc . *Arquivos de Biological Tecnologia*, **29**:407-412.
- Callahan, R.J., Rabito, C.A., 1990. Radiolabelling of erythrocytes with technetium- ^{99m}Tc : role of band-3 protein in the transport of pertechnetate across the cell membrane. *Journal of Nuclear Medicine*, **31**:2004-2010.
- Chulasiri, M., Suthienkul, O., Pavaro, C., Wongkrajang, Y., 1986. Herbal extracts for diarrheal treatment: antibacterial activity in vitro. *Journal of Public Health*, **16**:21-35.
- Conde Garcia, E.A., Nascimento, V.T., Santiago Santos, A.B., 2003. Inotropic effects of extracts of *Psidium guajava* L. (guava) leaves on the guinea pig atrium. *Brazilian Journal of Medical and Biological Research*, **36**(5):661-668. [doi:10.1590/S0100-879X2003000500014]
- Cortes, J., Alonso, J.I., Ruiz-Oliva, F., Alvarez, S., Ormijana, J.S., Caton, B., Alcorta, P., 2003. Renal cell carcinoma detected on Tc-^{99m} -labelled red blood cell imaging. *Clinical Nuclear Medicine*, **28**:920-922. [doi:10.1097/01.rlu.0000093089.53875.3e]
- Cuellar, A.C., Lara, R.A., Zayas, J.P., 1984. *Psidium guajava* L. tamizaje fotiquímico y estudio del aceite esencial. *Revista Cubana de Farmacia*, **18**:92-99 (in Spanish).
- de Oliveira, J.F., de Oliveira, M.B.N., Ávila, A.S., Braga, A.C.S., Catanho, M.T.J.A., Jales, R.L.C., Cardoso, V.N., Bernardo-Filho, M., 2003a. Assessment of the effect of *Fucus vesiculosus* extract on the labelling of blood constituents with technetium- ^{99m}Tc and the histological modifications on the shape of the red blood cells. *Food and Chemical Toxicology*, **41**(1):15-20. [doi:10.1016/S0278-6915(02)00206-5]
- de Oliveira, J.F., Santos-Filho, S.D., Catanho, M.T.J.A., Srivastava, S.C., Lima-Filho, G.L., Bernardo-Filho, M., 2003b. Effect of extract of medicinal plants on the labelling of blood elements with technetium- ^{99m}Tc and on the morphology of red blood cells (RBC): toxicological actions of roast coffee beans (*Coffea arabica*). *Indian Journal of Nuclear Medicine*, **18**:52-56.
- Dewanjee, M.K., Rao, S.A., Penniston, J.T., 1982. Mechanism of red blood cell labelling with ^{99m}Tc -pertechnetate and the role of cation pumps at RBC membrane on distribution and binding of Sn^{+2} and ^{99m}Tc with membrane proteins and hemoglobin. *Journal of Labelled Compounds and Radiopharmaceuticals*, **11**:1464-1466.
- Early, P.J., Soddee, D.B., 1995. *Principles and Practice of Nuclear Medicine*. Mosby—Year Book, Inc., Toronto, Canada, p.877.
- Fonseca, A.S., Frydman, J.N.G., Santos, R., Bernardo-Filho, M., 2005. Influence of antipyretic drugs on the labelling of blood elements with technetium- ^{99m}Tc . *Acta Biologica Hungarica*, **56**(3-4):275-282. [doi:10.1556/ABiol.56.2005.3-4.10]
- Frydman, J.N.G., Oliveira, M.B.N., Santos, A.E.O., Fonseca, A.S., Santos, R., Bernardo-Filho, M., 2004. Influence of methylxanthines on the labelling of blood elements with ^{99m}Tc . *Pakistan Journal of Biological Science*, **4**:521-524.
- Gonçalves, J.L.S., Lopes, R.C., Oliveira, D.B., Costa, S.S.,

- Miranda, M.M.F.S., Romano, M.T.V., Santos, N.S.O., Wigg, M.D., 2005. In vitro anti-rotavirus activity of some medicinal plants used in Brazil against diarrhea. *Journal of Ethnopharmacology*, **99**(3):403-407. [doi:10.1016/j.jep.2005.01.032]
- Gutflen, B., Ribeiro, B.L.A.R., Mattos, M.F., Ribeiro, C.R., Bernardo-Filho, M., 1992. Labelling of thymine with technetium-99m: suggestion of a chemical model. *Archives of Biological Technology*, **39**:69-74.
- Harel, F., Dupuis, J., Benelfassi, A., Ruel, N., Gregoire, J., 2005. Radionuclide plethysmography for non-invasive evaluation of peripheral arterial blood flow. *American Journal of Physiology, Heart and Circulation Physiology*, **289**(1):H258-H262. [doi:10.1152/ajpheart.01021.2004]
- Jaiarj, P., Khoohaswan, P., Wongkrajang, Y., Peungvicha, P., Suriyawong, P., Saraya, M.L.S., Ruangsomboon, O., 1999. Anticough and antimicrobial activities of *Psidium guajava* Linn. Leaf extract. *Journal of Ethnopharmacology*, **67**(2):203-212. [doi:10.1016/S0378-8741(99)00022-7]
- Jin, R.B., Ma, X.L., Wen, J.L., Tang, W.J., 2004. Application of radionuclide imaging to hepatic impact injury in rabbits. *Chinese Journal of Traumatology*, **7**:45-48.
- Khan, M.L.H., Ahmad, J., 1985. A pharmacognostic study of *Psidium guajava* L. *International Journal of Crude Drug Research*, **23**:95-103.
- Kimura, S., Tamaki, T., Aoki, N., 1985. Acceleration of fibrinolysis by the N-terminal peptide of alpha 2-plasmin inhibitor. *Blood*, **66**:157-160.
- Liu, H.N., 1988. Chinese Medicinal Herbs of Hong Kong, Vol. 2. Hang Chiewing As Kwang, Hong Kong, p.104-105.
- Lozoya, X., 1999. Xiuhpatli, *Herba officinalis*. SSA/UNAM, Mexico, p.127 (in Spanish).
- Lozoya, X., Meckes, M., Aboud-Zaid, M., Tortoriello, J., Nozolillo, C., Arnason, J.T., 1994. Quercetine glycosides in *Psidium guajava* L. leaves and determination of a spasmolytic principle. *Archives of Medical Research*, **25**:11-15.
- Lutterodt, G.D., 1992. Inhibition of Microlax-induced experimental diarrhea with narcotic-like extracts of *Psidium guajava* leaf in rats. *Journal of Ethnopharmacology*, **37**(2):151-157. [doi:10.1016/0378-8741(92)90073-Z]
- Lutterodt, G.D., Maleque, A., 1988. Effects on mice locomotor activity of a narcotic-like principle from *Psidium guajava* leaves. *Journal of Ethnopharmacology*, **24**(2-3):219-231. [doi:10.1016/0378-8741(88)90155-9]
- Meckes, M., Calzada, F., Tortoriello, J., González, J.L., Martinez, M., 1996. Terpenoids isolated from *Psidium guajava* with depressant activity on central nervous system. *Phytotherapy Research*, **10**(7):600-603. [doi:10.1002/(SICI)1099-1573(199611)10:7<600::AID-PTR918>3.0.CO;2-7]
- Mercadante, A.Z., Steck, A., Pfander, H., 1999. Carotenoids from guava (*Psidium guajava* L.): isolation and structure elucidation. *Journal of Agricultural and Food Chemistry*, **47**(1):145-151. [doi:10.1021/jf980405r]
- Morales, M.A., Tortoriello, J., Meckes, M., Paz, D., Lozoya, X., 1994. Calcium-antagonist effect of quercetin and its relation with the spasmolytic properties of *Psidium guajava* L. *Archives of Medical Research*, **25**:17-21.
- Moreno, S.R., Freitas, R.S., Rocha, E.K., Lima-Filho, G.L., Bernardo-Filho, M., 2004. Protection of plasmid DNA by a *Ginkgo biloba* extract from the effects of stannous chloride and the action on the labelling of blood elements with technetium-99m. *Brazilian Journal of Medical and Biological Research*, **37**(2):267-271. [doi:10.1590/S0100-879X2004000200015]
- Olds, G.D., Cooper, G.S., Chak, A., Sivak, M.V.Jr, Chitale, A.A., Wong, R.C., 2005. The yield of bleeding scans in acute lower gastrointestinal hemorrhage. *Journal of Clinical Gastroenterology*, **39**(4):273-277. [doi:10.1097/01.mcg.0000155131.04821.f3]
- Osman, A.M., Younes, M.E., Sheta, A.E., 1974. Triterpenoids of the leaves of *Psidium guajava*. *Phytochemistry*, **13**(9):2015-2016. [doi:10.1016/0031-9422(74)85152-6]
- Podhajcer, O.L., Friedlander, M., Graziani, Y., 1980. Effect of liposome-encapsulated quercetin on DNA synthesis, lactate production, and cyclic adenosine 3':5'-monophosphate level in Ehrlich ascites tumor cells. *Cancer Research*, **40**:1344-1350.
- Qa'dan, F., Thewaini, A.J., Ali, D.A., Affi, R., Elkhawad, A., Matalka, K.Z., 2005. The antimicrobial activities of *Psidium guajava* and *Juglans regia* leaf extracts to acne-developing organisms. *The American Journal of Chinese Medicine [An International Journal of Comparative Medicine East and West]*, **33**(2):197-204. [doi:10.1142/S0192415X05002783]
- Qian, H., Nihorimbere, V., 2004. Antioxidant power of phytochemicals from *Psidium guajava* leaf. *Journal of Zhejiang University Science*, **5**(6):676-683. [doi:10.1631/jzus.2004.0676]
- Re, L., Barocci, S., Capitani, C., Vivani, C., Ricci, M., Rinaldi, L., Paolucci, G., Scarpantonio, A., Leon-Fernandez, O.S., Morales, M.A., 1999. Effects of some natural extracts on the acetylcholine release at the mouse neuromuscular junction. *Pharmacological Research*, **39**(3):239-245. [doi:10.1006/phrs.1998.0433]
- Saha, G.B., 2003. Normal Biodistribution of Diagnostic Radiopharmaceuticals. In: Hladik, W.B., Saha, G.B., Study, K.T. (Eds.), *Essentials of Nuclear Medicine Science*. Williams and Wilkins, Baltimore, USA, p.3-19.
- Sampson, C.B., 1996. Complications and difficulties in radiolabelling blood cells: a review. *Nuclear Medicine Communications*, **17**:648-658.
- Slart, R.H., Phan, T.T., Talsma, M.D., Jager, P.L., 2004. Different splenic uptake of Tc-99m sulfur colloid and Tc-99m heat-denatured red blood cells in an infant with complete situs inversus. *Clinical Nuclear Medicine*, **29**:590-591. [doi:10.1097/01.rlu.0000134998.60745.1a]
- Valenca, S.S., Lima, E.A., Diré, G.F., Bernardo-Filho, M., Porto, L.C., 2005. Sodium pertechnetate ($\text{Na}^{99\text{m}}\text{TcO}_4$) biodistribution in mice exposed to cigarette smoke. *BMC Nuclear Medicine*, **5**(1):1. [doi:10.1186/1471-2385-5-1]
- Vanacker, S.A.B.E., Tromp, M.N.J.L., Haenen, G.R.M.M.,

- Vandervijgh, W.J.F., Bast, A., 1995. Flavonoids as scavengers of nitric oxide radical. *Biochemical and Biophysical Research Communications*, **214**(3):755-759. [doi:10.1006/bbrc.1995.2350]
- Verdu, J., Martinez, A., Anton, M.A., Munoz, J.M., Riera, M., Jover, R., Caballero, O., 2005. Increased thallium-201 uptake and Tc-99m red blood cell accumulation in hemangioma. *Clinical Nuclear Medicine*, **30**(1):25-26. [doi:10.1097/00003072-200501000-00007]
- Wong, K.T., Beauvais, M.M., Melchior, W.R., Snyder, S.P., 2004. Enhanced liver uptake of Tc-99m-labelled RBCs during gastrointestinal bleed scintigraphy using transfused RBCs compared with autologous RBCs. *Clinical Nuclear Medicine*, **29**:522-523. [doi:10.1097/01.rlu.0000132956.48831.14]
- Zaman, M.U., Hussain, R., Sajjad, Z., Ahmad, M.N., 2004. Localization of upper GI bleed on delayed imaging with ^{99m}Tc tagged RBC angiogram. *Journal of Pakistan Medicine Association*, **54**:490-491.

Editors-in-Chief: Pan Yun-he & Peter H. Byers
ISSN 1673-1581 (Print); ISSN 1862-1783 (Online), monthly

Journal of Zhejiang University

SCIENCE B

www.zju.edu.cn/jzus; www.springerlink.com

jzus@zju.edu.cn

JZUS-B focuses on "Biomedicine, Biochemistry & Biotechnology"

JZUS-B online in PMC: <http://www.pubmedcentral.nih.gov/tocrender.fcgi?journal=371&action=archive>

➤ Welcome your contributions to JZUS-B

Journal of Zhejiang University SCIENCE B warmly and sincerely welcome scientists all over the world to contribute Reviews, Articles and Science Letters focused on **Biomedicine, Biochemistry and Biotechnology**. Especially, Science Letters (3~4 pages) would be published as soon as about 30 days (Note: detailed research articles can still be published in the professional journals in the future after Science Letters is published by *JZUS-B*).

➤ Contribution requests

- (1) Electronic manuscript should be sent to jzus@zju.edu.cn only. If you have any questions, please feel free to visit our website (<http://www.zju.edu.cn/jzus>) and hit "For Authors".
- (2) English abstract should include Objective, Method, Result and Conclusion.
- (3) Tables and figures could be used to prove your research results.
- (4) Full text of the Science Letters should be in 3~4 pages. The length of articles and reviews is not limited.
- (5) Please visit our website (<http://www.zju.edu.cn/jzus/pformat.htm>) to see paper format.