in the airways and help patients expectorate. Despite these drugs having no effect on lung function, some studies have found that their regular use reduces the frequency of exacerbations of COPD and is associated with fewer days of ill health.

Two mucolytic agents are currently licensed for use in the United Kingdom (carbocysteine and mecysteine hydrochloride) and are generally well tolerated. They may be tried in patients with productive cough who are troubled by frequent exacerbations, although further data are required to clarify their place in COPD management. Increased oxidative stress has been implicated in the inflammatory response in COPD, and N-acetyl cysteine, another mucolytic, may confer some benefit in terms of its antioxidant properties.

Other drugs

There are no convincing data that prophylactic antibiotics are of use in stable COPD. Indeed, such treatment may only encourage the emergence of strains of bacteria resistant to conventional antibiotics. Cough is a troublesome symptom in many patients, but it may be advantageous, especially in individuals who produce copious amounts of sputum. Cough suppressants are not known to provide any benefit other than perhaps short term symptomatic control of cough, and their regular use is not indicated.

In patients with cor pulmonale there is little evidence that drugs such as angiotensin converting enzyme inhibitors, digoxin, or calcium channel blockers are of any benefit. However, if measures such as leg elevation and, where appropriate, long term oxygen therapy fail to control symptomatic peripheral oedema, low dose diuretics can be tried.

Further reading

- Rennard SI. Treatment of stable chronic obstructive pulmonary disease. *Lancet* 2004;364:791-802
- Ram FSF, Jones PW, Castro AA, de Brito Jardim JR, Atallah AN, Lacasse Y, et al. Oral theophylline for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2002;(4):CD003902
- Barnes PJ. Theophylline: new perspectives on an old drug. *Am J Respir Crit Care Med* 2003;167:813-8
- Lipworth BJ. Phosphodiesterase-4 inhibitors for asthma and chronic obstructive pulmonary disease. *Lancet* 2005;365:167-75
- Poole PJ, Black PN. Oral mucolytic drugs for exacerbations of chronic obstructive pulmonary disease: systematic review. *BMJ* 2001;322: 1271-4

Graeme P Currie is specialist registrar in the Respiratory Unit, Aberdeen Royal Infirmary, Aberdeen. Daniel K C Lee is specialist registrar in the Department of Respiratory Medicine, Papworth Hospital, Papworth Everard, Cambridge. Brian J Lipworth is professor in the Asthma and Allergy Research Group, Division of Medicine and Therapeutics, Ninewells Hospital and Medical School, Dundee.

The ABC of chronic obstructive pulmonary disease is edited by Graeme P Currie. The series will be published as a book by Blackwell Publishing in autumn 2006.

Competing interests: GPC has received funding for attending international conferences and honoraria for giving talks from pharmaceutical companies GlaxoSmithKline, Pfizer, and AstraZeneca. BJL has received funding for attending conferences, payment for speaking and consulting activity, and research funding from pharmaceutical companies AstraZeneca, Atlanta, Aerocrine, Sepracor, MSD, Neolab, Cipla, Innovata, UCB-Celltech, and Schering-Plough.

The figure of the additive effects of theophylline and salmeterol was reproduced with permission from Zu Wallack et al. *Chest* 2001;119: 1661-70.

BMJ 2006;332:1497-9

bmjlearning.com

Class war

"We can formulate the genetic law of cultural development in the following way: any function in the child's cultural development appears on stage twice, on two planes. First it appears on the social plane, then on the psychological, first among people as an interpsychical category and then within the child as an intrapsychical category." So claimed Lev Vygotsky, a Soviet constructivist psychologist in the first half of the last century.¹ Quite what he meant I couldn't say, but I do know that Vygotsky's name has become fashionable among educators, and not least medical educators. Many of his ideas have been enthusiastically applied to medical educational theory.

Educational theory is no doubt a good thing, but can you get too much of it? Are we in danger of missing the wood for the trees? Many traditionalists would argue that we are. We may be educating doctors in a sound format, but are they learning important content? A worrying study by Griffiths and Roberts showed that 45% of junior doctors don't know how to put in a chest drain in a patient with a pneumothorax and, if asked to do one, would insert the drain outside the "triangle of safety" recommended by the British Thoracic Society.² The triangle of safety is bordered by the anterior border of the latissimus dorsi, the lateral border of the pectoralis major muscle, a line superior to the horizontal level of the nipple, and an apex below the axilla.

This is an important piece of information—inserting a drain outside this triangle could do some serious damage.

In his book *Class War*, the former chief inspector of schools Chris Woodhead says that we have got lost in a maze of pedagogy, and, as a result, our children are failing to learn important pieces of information. He berates academics who claim that being able "to critically analyse work" is more important than learning content (and also bemoans their inability to spot a split infinitive). Could medical education become lost in a similar sea of pedagogy?

At BMJ Learning, we concentrate on content rather than theory—one of our most recent modules is on chest x ray interpretation. It enables you to train and test your skills in diagnosing pleural effusions and pneumothoraces. You might also learn the right place to put in a chest drain, rather than a Bermuda triangle of your own choosing.

Kieran Walsh *clinical editor, BMJ Learning*
(bmjlearning@bmjgroup.com)

1 The Vygotsky project. <http://webpages.charter.net/schmolze1/vygotsky/vygotsky.html>.

2 Griffiths JR, Roberts N. Do junior doctors know where to insert chest drains safely? *Postgrad Med J* 2005;81:456-8.