paigns were more likely to support voter registration in the clinic (18 of 23 vs 4 of 12, $P = .02$). PCPs who primarily served low-income patients were less likely to regard discussions about politically oriented health care issues as appropriate (0 of 5 vs 13 of 22, $P = .04$). In conclusion, the majority of our study PCPs discussed voting or health care politics with their patients. To our knowledge, this is the first study to demonstrate that the personal political activities of physicians are associated with their willingness to address political issues with clinic patients. Moreover, we found that gender was associated with political activity inside and outside the clinic, a finding consistent with a study of English PCPs.⁵ Our study was limited by the small sample size, the single health care system source of our cohort, unadjusted analyses, and self-report. Future studies should continue to explore the political attitudes and activities of physicians in different populations and settings, in addition to addressing the influence of such activities on the patients they serve. Moreover, we believe that a debate on the ethical issues surrounding political discourse in the practice setting is warranted.—**Craig A. Umscheid, MD, Bruce Y. Lee, MD, MBA, Division of General Internal Medicine; (DGIM) and Center for Clinical Epidemiology and Biostatistics (CCEB); Robert Gross, MD, MSCE, Division of Infectious Disease and CCEB; and Barbara Turner, MD, MEd, DGIM, University of Pennsylvania School of Medicine, Philadelphia, Pa, USA.**

REFERENCES

1. **Jameson MG.** Physicians and American political leadership. *JAMA*. 1983;249:929–30.
2. **Frank E.** Political self-characterization of US women physicians. *Soc Sci Med*. 1999;48:1475–81.
3. **Myllykangas M, Ryyanen OP, Kinnunen J, Takala J.** Comparison of doctors', nurses', politicians' and public attitudes to health care priorities. *J Health Serv Res Policy*. 1996;1:212–6.
4. **Wakeford RE, Allery L.** Doctors' attitudes, medical philosophy, and political views. *Br Med J (Clin Res Ed)*. 1986;292:1025–7.
5. **Cooke M, Ronalds C.** Women doctors in urban general practice: the doctors. *Br Med J (Clin Res Ed)*. 1985;290:755–8.

Mortality Associated with Hormone Replacement Therapy in Younger and Older Women

In Reply:—I appreciate the opportunity to respond to Dr. Grant's letter concerning our meta-analysis on mortality associated with hormone replacement. It is true that no single trial of younger women has shown a significant reduction in mortality with hormone use. In fact, no trial of hormone replacement has shown a decrease or increase in mortality. Death is a very rare event, especially in younger postmenopausal women, and it would require a very large trial to provide the statistical power necessary to produce significant results. A more precise estimate can be made by pooling the results of many smaller trials in the form of a meta-analysis.

In the analysis of younger women, approximately one half of the deaths were from one trial with ovarian cancer survivors that provided 40% of the weight. However, as described in the article, when this trial was excluded from the analysis there still was a significant reduction in mortality for the younger age

group, with an odds ratio of 0.56 (confidence interval [CI], 0.31 to 0.99). Since our analysis was published, the Women's Health Initiative estrogen-only trial has shown a hazard ratio for total mortality of 0.73 (CI, 0.47 to 1.13) in those 50 to 59 years old.¹ If these data were added to our meta-analysis, it would provide 50% of the weight, with a revised odds ratio for mortality in the younger age group of 0.67 (CI, 0.49 to 0.92).

It is not possible to completely control for previous exposure to hormone replacement in clinical trials, but randomized trials help to reduce baseline differences between the treatment and control groups. In this meta-analysis, the women in the younger age group had a mean age of 53.7 years and had no recent exposure to hormone replacement, which indicates that their previous exposure to hormone replacement was probably minimal. Longer trials of hormone replacement will be needed in the future to assess whether this mortality benefit seen in younger women persists over the years.—**Shelley Salpeter, MD, FACP, Santa Clara Valley Medical Center, San Jose, CA.**

REFERENCE

1. **Anderson GL, Limacher M, Assaf AR, et al.** Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: the Women's Health Initiative randomized controlled trial. *JAMA*. 2004;291:1701–12.

To the Editor:—We commend Dr. Shanafelt et al.¹ on their recent study noting an association between higher mental well-being and residents' capacity for empathy, comprising the "cognitive" capacity for insight into others' experiences and the "emotive" capacity to respond to these experiences. Although the use of the SF-8 to measure well-being only presents a limited assessment of aspects of physical and mental health, their findings support the need for further empiric work about the nature and promotion of resident well-being. In this study sample, female residents had higher emotive and cognitive empathy scores compared with men, but a lower proportion of female residents possessed "high mental well-being." This intriguing finding raises further questions about the interactions between empathy and well-being and the implications for future research.

A recent review described clinical empathy toward patients as emotional labor.² Physicians may use 2 different modes of acting to express empathy: (1) "surface acting"—displaying behaviors unmatched by underlying emotional content and (2) "deep acting"—modifying their underlying emotions to create congruency with these behaviors. Larson and Yao argue that recurrent surface acting by physicians may actually lead to cynicism and burnout, although there is a lack of empiric studies to support this belief. If this conjecture is true, residents who possess a high capacity for empathy may have decreased mental well-being over time because of this persistent dissonance between their underlying emotional state and their performance as empathic physicians.

Another concern is that too much empathy for patients could lead physicians to sacrifice aspects of their own well-being. Some physicians with high empathy may be unable to set the necessary boundaries to protect their own well-being. Huggard³ describes the phenomenon of "compassion fatigue," whereby physicians who engage empathically with their patients develop burnout because of secondary traumatic stress.

Perhaps in Dr. Shanafelt's study, the wellness promotion practices of residents with high mental well-being enable them to demonstrate empathic concern while still maintaining their own well-being and averting burnout.

Striking this balance between empathy and self-care may be a critical skill for residents to master and may help to ensure well-being throughout their careers. Further explorations of gender differences in clinical empathy may help elucidate how the risk of burnout or compassion fatigue may differ between female and male physicians. We appreciated the contribution of Dr. Shanafelt and his colleagues to our understanding of these important connections.—**Neda Ratanawongsa, MD, Scott M. Wright, MD, Rachel B. Levine, MD, MPH, Johns Hopkins School of Medicine, Baltimore, MD.**

REFERENCES

1. **Shanafelt TD, West C, Zhao X, et al.** Relationship between increased personal well-being and enhanced empathy among internal medicine residents. *J Gen Intern Med.* 2005;20:559–64.
2. **Larson EB, Yao X.** Clinical empathy as emotional labor in the patient-physician relationship. *JAMA.* 2005;293:1100–6.
3. **Huggard P.** Compassion fatigue: how much can I give? *Med Educ.* 2003;37:163–4.

Authors' Response

In Reply:—We thank Drs. Ratanawongsa and colleagues for their insightful comments. Larson and Yao's¹ concept of "surface acting" and "deep acting" is a useful method to conceptualize the ways physicians express empathy. In addition to potential effects on the patient's experience, the way physicians respond to patients can positively (enhanced meaning/personal fulfillment) or negatively (burnout, compassion fatigue) impact physician outcomes.

Although we believe surface acting is appropriate in many situations, the motivation behind use of this technique is important. We agree with Larson and Yao's¹ assertion that surface acting motivated by a "value guided commitment to care" can contribute to both patient and physician satisfaction. In contrast, repetitive use of surface acting simply to avoid dealing with patients' emotional concerns is likely to contribute to cynicism.

Deep acting has the potential to create an even greater connection between physicians and patients as well as enhance physicians' sense of meaning in their work. This approach, however, involves physicians altering their own emotional state to identify with the patient's experience and has the potential to create reactive emotions (anger, distress, concern, and grief) for the physician. Physicians who are able to recognize and process such personal reactions are better able to express empathy without experiencing "compassion fatigue." We believe training medical students and residents to calibrate their use of surface and deep acting and identify personal reactions in each patient encounter is an important step in developing physicians who are both empathetic and resilient.^{2–4}

The energy physicians can devote to emotional labor is ultimately a limited resource. We hypothesize that attention to

personal well-being through mindful practice and nurturing of personal interests and relationships allows physicians to renew the energy they have to devote to this work.⁵ This concept holds that empathy and well-being have reciprocal feedback loops. The complexity of these interactions highlights the need for additional studies on the relationship between well-being and empathy and research that provides insight into how physician well-being can be promoted. We hope that support for such studies will grow in the coming years.—**Tait Shanafelt, MD, Colin West, MD, PhD, Joseph Kolars, MD, and Thomas Habermann, MD, Mayo Clinic College of Medicine, Rochester, MN.**

REFERENCES

1. **Larson E, Yao X.** Clinical empathy as emotional labor in the patient-physician relationship. *JAMA.* 2005;293:1100–6.
2. **Epstein R.** Mindful practice. *JAMA.* 1999;282:833–9.
3. **Novack DH, Suchman AL, Clark W, Epstein RM, Najberg E, Kaplan C.** Calibrating the physician. Personal awareness and effective patient care. Working group on promoting physician personal awareness, American academy of physician and patient. *JAMA.* 1997;278:502–9.
4. **Novack D, Epstein R, Paulsen R.** Toward creating physician-healers: fostering medical students' self-awareness, personal growth, and well-being. *Acad Med.* 1999;74:516–20.
5. **Shanafelt T, Sloan J, Habermann T.** The well-being of physicians. *Am J Med.* 2003;114:513–7.

To the Editor:—Franks et al.'s¹ article entitled "Racial disparities in the content of primary care office visits" observed that African-American (AA) patients are less likely to receive preventive care during office visits to primary care. The authors present an interesting study and do a good job enumerating and addressing the limitations of their dataset and analyses. Still, we feel 2 points merit further discussion:

1. That AA receive more of some preventive services and less of others does not make intuitive sense. If ethnicity were a direct determinant of preventive services, then the relationship should be independent of the type of service in question. In absence of a consistent relationship between ethnicity and the type of services offered, we must consider the existence of an unadjusted confounder. Attempts to explain this phenomenon should consider not only provider characteristics, but also those of the patient and site of care as well.
2. That the disparities observed for one time period do not persist into the following one suggests that, regardless of the mechanism for the disparities, they are waning over time. Rather than focus on the causal mechanism for disparities, we should celebrate that the only remaining significant disparity by 1995 to 2001 was in rectal screening (a procedure not recommended by the U.S. Preventive Services Task Force <http://www.ahcpr.gov/clinic/uspstf/uspstfprca.htm>).

Future research should try to understand what might account for this improvement so as to perpetuate its benefits.—**Maria Silveira, MD, MA, MPH, and Eve Kerr, MD, MPH, Ann Arbor VA Health Services Research and Development Center of Excellence, Ann Arbor, MI.**

REFERENCE

1. **Franks P, Fiscella K, Meldrum S.** Racial disparities in the content of primary care office visits. *J Gen Intern Med.* 2005;20:599–603.