Obesity is less common in richer countries

Roger Dobson *Abergavenny*

Almost one in five people in Hungary are obese, three times the proportion in Norway, Italy, and France, says a new report.

The study, which was based on data from 24 countries in Europe and published online before print publication on 14 June in the *European Journal of Public Health* (www.oxfordjournals.org, doi: 10.1093/eurpub/ckl073), shows that Central and Eastern European nations have some of the highest rates, although the prevalence of obesity among men is higher in the United Kingdom than in Russia, where women are nearly twice as likely to be obese as men.

The results show that the prevalence of obesity varies with economic and other indicators. There was less obesity in countries with a higher gross domestic

product and in nations with higher percentages of city dwellers.

The authors used national surveys to determine the sex specific prevalence of obesity—the percentage of people with a body mass index ≥30.

The results show a wide range of prevalence across countries and between the sexes. The prevalence among men ranged from 7.1% in Italy to 18.4% in

Hungary and among women from 5.8% in Norway to 20.4% in Hungary. The overall prevalence ranged from 6.2% in Norway to 19.5% in Hungary.

The highest prevalence for both sexes was reported from Central and Eastern European countries, ranging from 9% to 18% (median 15%) in men and from 12% to 20% (17%) among women. □

NEJM corrects its study on rofecoxib

Susan Mayor *London*

An increase in cardiovascular events associated with the cyclooxygenase-2 inhibitor rofecoxib (Vioxx) occurred much earlier in the course of treatment than the original study reporting the problem had indicated, says a correction published online in the *New England Journal of Medicine* on 26 June (<http://content.nejm.org>, doi: 10.1056/NEJMc0606).

The correction to the study of cardiovascular events associated with rofecoxib in the colorectal adenoma chemoprevention trial (*New England Journal of Medicine* 2005;352:1092-102) explains that the test for proportionality of hazards used linear time rather than the logarithm of time as had been specified in the methods section.

Analysis using the logarithm of time means that the Kaplan-Meier curves separated 18 months after randomisation. However, the results of an overall test of the proportional hazards assumption for the entire 36 month observation period were not significant ($P=0.07$).

In a letter also published online in the journal on 26 June (doi: 10.1056/NEJMc066260) Steven Nissen, of the Cleveland Clinic Foundation, Cleveland, Ohio, notes: "The original article included a post hoc hypothesis that curves for confirmed thrombotic events would not begin to diverge until after 18 months of exposure to rofecoxib. However, all intention-to-treat analyses in the newly released report show that the event curves begin to diverge much earlier, generally within four to six months.

"The most useful Kaplan-Meier curves, involving intention-to-treat analysis of the APTC [Antiplatelet Trialists' Collaboration] end point, show divergence after only three months of exposure to rofecoxib," the letter says.

Merck voluntarily withdrew rofecoxib from the market in 2004 because of questions raised by data from the study and because other treatment choices for patients were available. □

Direct to consumer medical tests are offered in United States

Janice Hopkins Tanne *New York*

"Internet companies and some hospitals in the United States are offering medical tests directly to consumers without the need for a doctor's recommendation, reports the *Wall Street Journal* (20 Jun, p D4).

Although welcomed by consumers' groups, the move has sparked concerns among doctors about patients' ability to interpret results and about legal implications.

Gene Herbek, head of the pathology centre at Methodist Hospital in Omaha, Nebraska, told the *BMJ* that his hospital would begin offering tests within a few weeks at two walk-in clinics. Its tests include those for blood glucose concentrations, thyroid disease, blood lipid concentrations, hepatitis B, Epstein-Barr virus, pregnancy, and prostate specific antigen, as well as screening for drug misuse. Patients will also be able to order a kit for faecal occult blood testing.

Dr Herbek said, "It's a nation-

al trend—a service patients want.

"The idea is to encourage patients to know their family history and to take charge of their health," he said. Pharmacies already sell over the counter tests for blood glucose concentrations, pregnancy, and melanin concentrations (which are commonly believed to be related to sleep problems). Many also offer free blood pressure testing by machine.

Patients will be able to ask for the tests they want, pay up front, and receive the results by post. Abnormal results will prompt a phone call advising the patient to see a doctor. Patients will be given an explanation of the test and a number to call if they have questions. The hospital can refer them to one of its staff doctors if patients don't have a doctor.

Internet sites such as MedLabUSA.com, MyMedLab.com, and DirectLabs.com offer a nationwide service by acting as online brokers, allowing users to

enter their postcode and get the addresses of several nearby laboratories. Results from tests these laboratories offer are sent directly to the consumer. Patients are contacted if serious abnormalities are picked up.

Privacy is a key factor in the growth of direct to consumer tests, said Dr Herbek. "Patients want to be the first to know," he said. Abnormal tests can affect people's ability to get life or health insurance. Convenience is another reason. Tests are cheaper, because patients pay up front and administrative costs are less.

But insurance companies will not reimburse patients for the cost of tests unless they have been ordered by a doctor. The College of American Pathologists says that pathology clinics offering these tests should consider whether a patient can interpret the results without the help of a doctor.

Arthur Levin, director of the Center for Medical Consumers in New York, said that the companies were selling "worry to the worried well" and that money was being wasted. Tests might have false positive results, leading to further inappropriate testing, he added. □