In brief

European alcohol consumption is double the global average: In 2002 drinking in Europe was responsible for 6.1% of deaths, 12.3% of all years of life lost, and 10.7% of all disability adjusted life years, a report in *Addiction* says (doi: 10.1111/j.1360-0443.2006.01491.x). Each European drank on average 12.1 l of pure alcohol, more than double the average global consumption of 5.8 l.

Dutch council calls for trials of chlamydia screening: The Dutch Health Council is recommending randomised trials of screening of 15 to 30 year olds in the country's largest cities for chlamydia. It says that this is justified by the 3.2% prevalence of the infection among people in cities.

Germany revises law on tobacco advertising: Germany will introduce its own bill to outlaw the advertising of tobacco products after pressure from the European Court of Justice. Germany had challenged a European Union law banning tobacco advertising in print, on the radio, and online.

Mental ill health among UK employees triples in five years: UK occupational health doctors saw three times as many people with stress and mental illness in 2001 as they did in 1996. The proportion of all visits involving mental ill health more than tripled, rising to 36.7% from 11.4% (*Occupational Medicine*, doi: 10.1093/occmed/kql041).

Red Cross accepts Israel's first aid society as member: The International Committee of the Red Cross and Red Crescent Movement has accepted Israel's national first aid and ambulance organisation, Magen David Adom, as a full member 57 years after it first applied. (*BMJ* 2005;331:654).

NICE issues guidance on atrial fibrillation: The UK's National Institute for Health and Clinical Excellence and the Royal College of Physicians have jointly produced new guidance on atrial fibrillation to increase its detection and treatment (www.nice.org.uk/CG036).

NICE removes β blockers as first line treatment for hypertension

Susan Mayor *London*

β blockers are no longer recommended for the first line or second line treatment of hypertension, says an updated guideline for England and Wales.

The guideline recommends that for hypertensive patients aged 55 or older or for black patients of any age the first choice for initial treatment should be either a calcium channel blocker or a thiazide-type diuretic. Black patients are considered to be those of African or Caribbean descent, not people of Asian descent or mixed race people. For patients under the age of 55 an angiotensin converting enzyme (ACE) inhibitor (or an angiotensin receptor blocker if an ACE inhibitor is not tolerated) is recommended as first line treatment.

The updated guideline was developed by the National Institute for Health and Clinical Excellence (NICE)—the body that advises on use of treatments by the NHS in England and Wales—the National Collaborat-

ing Centre for Chronic Conditions, and the British Hypertension Society after an expert review of data from new studies and data on adverse events and a detailed health economic analysis that compared the cost effectiveness of the main classes of antihypertensives.

Previous guidelines included β blockers as an option for first line treatment, but this was changed after the guideline development group reviewed evidence from clinical trials indicating that these drugs reduced blood pressure less effectively than other classes of antihypertensives, particularly in elderly people. Head to head trials showed that β blockers were usually less effective than the other drugs in reducing major cardiovascular events, particularly stroke. β blockers were also less effective than an ACE inhibitor or a calcium channel blocker in reducing the risk of type 2 diabetes; patients taking a β blocker and a thiazide-type

diuretic were at particular risk.

Bryan Williams, professor of medicine at University Hospitals of Leicester NHS Trust and a member of the guideline group, said: "This new guidance has benefited from a rigorous evaluation of the evidence and has resulted in a simple, pragmatic, and practical approach to treating blood pressure, one of the most important causes of premature death in the UK. The decision to recommend that β blockers should no longer be used as a routine initial treatment for high blood pressure is a bold—but correct—decision."

The guideline also advised on combination treatments. For patients whose blood pressure is not adequately controlled by monotherapy with a calcium channel blocker or a thiazide-type diuretic, it recommends adding an ACE inhibitor (or an angiotensin receptor blocker if an ACE inhibitor is not tolerated). If initial treatment was with an ACE inhibitor, a calcium channel blocker or a thiazide-type diuretic should be added. □

Hypertension: Management of Hypertension in Adults in Primary Care—Partial Update is available at www.nice.org.uk.

Publicity will highlight problem of obstetric fistula

Andrew Cole *London*

Hundreds of thousands of women in developing countries suffer the devastating injury during child bearing of obstetric fistula. But the agency behind a global campaign to eradicate the condition says it is simple to prevent and easy to treat.

The global campaign, led by the international development agency the United Nations Population Fund, has launched a month long advertising drive in the United Kingdom to raise awareness of a condition that is believed to affect between 50 000 and 100 000 women each year.

Obstetric fistula is usually caused by several days of obstructed labour without prompt medical intervention and leaves the woman with ago-

A woman attends the Addis Ababa fistula hospital, Ethiopia

nising, long term pain, chronic incontinence, and—in most cases—a stillborn baby.

Reconstructive surgery for the condition has a 90% success rate and costs, together with rehabilitation, around £170 (£250; \$310). But most women in developing countries with the condition are not treated at all.

"Obstetric fistula was elimi-

nated here in the UK more than 100 years ago," said the singer and model Natalie Imbruglia, a spokeswoman for the campaign. "It's unacceptable that women and girls in developing countries are still suffering from this entirely preventable and treatable condition."

The campaign to end fistula, launched in 2003, aims to help women with fistula in more than 35 countries of Africa and Asia. It focuses on three key areas: preventing fistula, treating affected women, and supporting women after surgery.

"Women with obstetric fistula are dead women—physically and psychologically," said Gloria Esegbona, a British doctor of Nigerian descent who has treated women with the condition in Africa. "Their injuries render them little more than cripples and scar their souls."

Ninety six per cent of these women lose their babies, she said.

For more on the campaign see www.endfistula.org.