interrogation of prisoners.

At a workshop in Washington, DC, to plan the alliance's approach, Steven Xenakis, a retired brigadier general and currently director of child and adolescent psychiatry at the Psychiatric Institute of Washington in Washington, DC, said that incidents such as the abuses of prisoners in Baghdad's Abu Ghraib prison "became possible because our men and women in uniform were given ambiguous instructions, which in some cases authorised treatment that went beyond what was allowed in the *Army Field Manual* and because the chain of command chose to look the other way."

The lobbyist Chris Anders said that most Congressmen and women feel that they have addressed the issue of torture, after having voted last year on the McCain amendment barring cruel, inhuman, or degrading treatment or punishment. They are therefore reluctant to take on the associated issue of extraordinary rendition, he said.

Bob Roehr *Washington, DC*

(See BMJ 2006;333:1462, 2 Jun.)

Incomes of US doctors are falling, survey finds

The average income of doctors in the United States fell by 7% from 1995 to 2003, says a new report from the Center for Studying Health System Change, a non-profit research group in Washington, DC.

Worst affected were primary care doctors, whose income—after deduction of expenses such as malpractice insurance but before taxes—fell by 10% over the period, while surgeons' income fell by 8%. Medical specialists' income remained relatively unchanged.

The incomes of lawyers and other professionals rose by 7% during the same period.

Static or declining fees from both public and private sources seem to be a major factor in doctors' lower incomes, says the report.

Yet the volume of services provided by doctors increased substantially, largely because of

the growth in the number of tests and procedures, says the report, which partly explains why medical specialists have seen their income falling at a slower pace than that of primary care doctors.

Cecil Wilson, chairman of the board of the American Medical Association, said that the survey "confirms what they [doctors] already know from their own practices: payments are not keeping up with inflation."

Fred Charatan *Florida*

For more information see www.hschange.org.

Drug company chief urges faster introduction of new vaccines in poor nations

The president of GlaxoSmith-Kline Biologicals, Jean Stéphenne, has urged the global health community to speed up the introduction of new life saving vaccines in poor countries and avoid the customary lengthy delays.

With the exception of the rotavirus vaccine, which was introduced in Latin America and Europe at the same time, so far vaccines had been introduced in developing countries some 10 to 20 years after developed countries had received them, he said.

Mr Stéphenne said that the rotavirus vaccine had proved that it was perfectly possible to hasten the process. "The infrastructure exists. If there is the willingness to introduce it it can be done within one or two years."

International health charities and funding initiatives could make a real difference, he said, but much depended on WHO and the Global Alliance for Vaccines and Immunization, he suggested.

Thomas Cherian, acting joint coordinator of the WHO's expanded programme on immunisation, said that the agency saw itself as more as a facilitator, helping countries make their own decisions. And decisions needed to be based on hard evidence, he said.

John Zarocostas *Geneva*

Faster delivery of drugs to heart attack patients cuts mortality

Adrian O'Dowd *London*

The NHS in England and Wales is the best healthcare system in the world for people who have had heart attacks, states the fifth annual report of the myocardial infarction national audit project.

The level of care for such patients is unequalled, says the report. It gathered data from all hospitals and ambulance services in England and Wales that cared for patients with suspected heart attack in 2005-6.

More lives are being saved now, and there has been a decrease in the number of deaths from heart attack, says the report, because of the improving performance of NHS organisations.

The project's work was carried out primarily by the clinical effectiveness and evaluation unit of the Royal College of Physicians on behalf of a multiprofessional steering group. The project is funded by the health standards watchdog body the Healthcare Commission in collaboration with the Central Cardiac Audit Database.

The new data show that the time taken to give patients thrombolytic treatment has become shorter recently and that the number of patients getting this treatment from ambulance staff before they reach hospital has risen dramatically.

Among the findings from 2005-6 are that: 83% of eligible patients in England received thrombolytic treatment within 30 minutes of arrival at hospital, compared with 44% in early 2001 (74% in Wales, compared with 65% in 2003-4, when Welsh hospitals were first included in

the report); 58% of patients received thrombolytic treatment within 60 minutes of calling for professional help in England, compared with 22% in early 2001 (30% in Wales, compared with 22% in 2003-4); and 2231 patients received pre-hospital thrombolytic treatment, compared with 1374 patients in 2004-05.

The decrease in the number of deaths is due, the report claims, to two developments. Firstly, more patients than ever are being treated very early by paramedics before they reach hospital—28 of the 31 ambulance services in England and Wales can now do this. Secondly, more patients are being treated by immediate angioplasty: 1647 patients in 2005-6 and 1087 in 2004-5.

John Birkhead, the project's clinical director, said: "This large audit confirms the provision of a high standard of care to people suffering heart attack. It allows those hospitals performing less well to put in place policies to improve performance and ambulance services to work with hospitals to deliver even more rapid treatments."

Jonathan Boyce, head of audit at the Healthcare Commission, said: "The fact that levels of pre-hospital thrombolysis have steadily increased is good news. This reassures patients who have had a heart attack that they will be treated in a timely way that limits the damage to the heart." □

How the NHS Manages Heart Attacks: Fifth Public Report 2006 can be seen at www.rcplondon.ac.uk/pubs.