LETTERS TO THE EDITOR

To the Editor:—I applaud Straus et al.¹ for evaluating a combination of clinical findings—history of chronic obstructive pulmonary disease (COPD), wheezes, and forced expiratory time (FET)—in evaluating test measures for COPD and for assessing time intervals for performing bedside maneuvers. However, I take issue with several points in the article. First, the authors downplay the significance of a forced expiratory time >9 seconds because the post-test probability increases only from 10% to 40%. A positive FET does have merit, insofar as spirometric testing, the gold standard, will be ordered. Second, the authors infer that the absence of the combination excludes COPD. A pretest probability of 40% decreases only to 17% with a likelihood ratio (LR) of 0.3. I doubt any clinician would dismiss COPD with a post-test probability of 17%. Third, the authors arrived at the LR +59 for the combination by multiplying LRs for each. They have included “crude” LRs without corresponding sensitivities and specificities for history of COPD or FET. The sensitivities, specificities, and likelihood ratios could be calculated from a 2 × 2 table using spirometry positive (disease) and spirometry negative (no disease) with index test positive (combination of findings) and index test negative (2/3 positive, 1/3 positive, or 0/3 positive) categories. Finally, I find it odd that the sensitivity for wheezing is 35% and the “estimated” sensitivity for the combination is 70% based on the LR −0.30. In order to obtain an LR +59, the false-positive rate must be slightly higher than 1%. As a result, LR− is represented by 30%/99% or 0.30. Intuitively, combinations of findings are much more likely to have lower sensitivities than single findings, whereas the specificity would be expected to be higher with combinations than with one finding.²—**DAVID A. NARDONE, MD**, Hillsboro, Ore, VHA Medical Center, Professor Emeritus, Oregon Health Sciences University.

REFERENCES

1. Straus SE, McAlister FA, Sackett DL, Deeks JJ. Accuracy of history, wheezing, and forced expiratory time in the diagnosis of chronic obstructive pulmonary disease. *J Gen Intern Med*. 2002;17:684–88.
2. Nardone DA, Roth KM, Mazur DJ, McAfee JM. Usefulness of physical examination in detecting the presence or absence of anemia. *Arch Intern Med*. 1990;150:201–4.

Reply:—We appreciate Dr. Nardone’s comments and his interest in our paper. We concur with Dr. Nardone that spirometry must be done in people with prolonged forced expiratory time.

As we stated in our article, forced expiratory time is not sufficient on its own to diagnose chronic obstructive pulmonary disease.¹ Likelihood ratios greater than 10 and less than 0.10 are very useful in either ruling in or ruling out a target disorder.² The pretest probability of disease becomes particularly important when likelihood ratios are between 0.10 and 10. In our previous paper, we stated that if the patient’s pretest probability is 10%,³ and applying a likelihood ratio of 0.3, the posttest

probability is sufficiently low that we would feel comfortable in ruling out the disease. However, if the pretest probability for a particular patient were higher, the posttest probability would be higher, such that we might not be comfortable ruling out the disease. Finally, the combined likelihood ratio of 59 was obtained from multiplying the adjusted likelihood ratios (adjusted for their nonindependence), not from the crude likelihood ratios. Therefore, they cannot be used directly to calculate sensitivity and specificity.—**SHARON E. STRAUS, MD, FINLAY A. MCALISTER, ON BEHALF OF THE CARE-COAD INVESTIGATORS**, Toronto General Hospital, Toronto, Ontario, Canada.

REFERENCES

1. Straus SE, McAlister FA, Sackett DL, Deeks JJ, On Behalf of the CARE-COAD2 Investigators. Accuracy of history, wheezing, and forced expiratory time in the diagnosis of chronic obstructive pulmonary disease. *J Gen Intern Med*. 2002;17:684–8.
2. Badgett RG, Tanaka DJ, Hunt DK, et al. Can moderate chronic obstructive pulmonary disease be diagnosed by historical and physical findings alone? *Am J Med*. 1993;94:188–96.
3. Sackett DL, Straus SE, Richardson WS, Rosenberg WM, Haynes RB. *Evidence-based Medicine: How to Practise and Teach It*. London: Harcourt Brace; 2000.

To the Editor:—McGee¹ does a service for us by focusing on the likelihood ratio as one of—if not the most important—operating characteristic of a diagnostic test. He suggests that its lack of use is due to the burdensome calculations necessary for conversion between odds and probabilities. Although this may be true, we offer an alternative explanation.

Probabilities differ from likelihoods in important ways.² With probabilities, the *data are random and hypotheses are fixed*, allowing one to develop conditional probabilities. Conversely, with likelihoods, the *data are fixed and the hypotheses are random*, as seen in a differential diagnosis. Likelihoods do not follow the rules of probability theory. Odds have made intuitive sense to the man on the street who bets on categorical events like winning a horserace. Odds also apply to such things as whether a solitary pulmonary nodule is malignant or benign or if a woman is pregnant or not. That we in medicine have chosen to remain with probabilities and dismiss odds shows that we have yet to grasp this subtle but important distinction.

With respect to an interpretation of likelihood ratios such as that given in McGee’s Table 2, Jaeschke et al.³ from the Evidence-Based Medicine Working Group note the following:

- “Likelihood ratios greater than 10 or less than 0.1 generate large and often conclusive changes from pretest to post-test probability.
- Likelihood ratios of 5 to 10 and 0.1 to 0.2 generate moderate shifts in pretest to posttest probability.
- Likelihood ratios of 2 to 5 and 0.2 to 0.5 generate small (but sometimes important) changes in probability.

- Likelihood ratios of 1 to 2 and 0.5 to 1 alter probability to a small (and rarely important) degree."

The divergent relationship between odds and probabilities is important in decision-making involving probabilities less than 10% and greater than 90%. If a man has a solitary pulmonary nodule with a 50% probability (odds of 1) of malignancy, would his doctor and he want a hypothetical test that had a likelihood ratio of 13 or one with 101 to rule it in? If the first test was positive, the probability would be 93%, compared with a probability of 99% with a positive second test—a mere difference of 6%. Using odds, after a positive first test, he would have 13 chances of having the malignancy to 1 chance of its being benign, whereas with the second test positive, his odds increase to 101 chances of having a cancer to 1 not. This example involves probability ranges beyond the formula proposed by McGee.

Dreyfus and Dreyfus⁴ have shown that the novice makes decisions in a protracted and laborious manner, using context-free rules, with no base of experience, having no responsibility for the problem, and using conscious analysis. Conversely, the expert gathers information fluently and expeditiously by focusing on the situation at hand, using his or her base of experience, having a strong sense of responsibility that drives an intuitive decision. The range suggested by McGee limits us to probabilities between 10% (odds 1 to 9) and 90% (odds 9 to 1), in which an intuitive sense is meager. While this is sufficient for decisions regarding appendicitis or streptococcal pharyngitis, it is inadequate in dealing with problems such as malignancy. —**JOHN C. PEIRCE, MD, MA, MS, RICHARD D. GERKIN, MD, MS**, Good Samaritan Regional Medical Center, Phoenix, Ariz.

REFERENCES

1. McGee S. Simplifying likelihood ratios. *J Gen Intern Med*. 2002;17:646–9.
 2. Goodman SN, Royall R. Evidence and scientific research. *Am J Public Health*. 1988;78:1568–74.
 3. Jaeschke R, Guyatt G, Sackett DL for the Evidence-Based Medicine Working Group. User's guide to the medical literature: III. How to use an article about a diagnostic test, B. What are the results and will they help me in caring for my patients? *JAMA*. 1994;271:703–7.
 4. Dreyfus HL, Dreyfus SE. Five steps from novice to expert. In: *Mind Over Machine: The Power of Human Intuition and Expertise in the Era of the Computer*. New York: The Free Press; 1986.
- Reply:*—I appreciate Peirce and Gerkin's comments and welcome the opportunity to respond.
- Peirce and Gerkin's distinction between probability and likelihood is unnecessary when applied to diagnostic testing (the subject of my paper), because unlike measures of therapeutic effect, diagnostic testing admits only 2 possible hypotheses: the proposed diagnosis of interest is either present or it is absent. These hypotheses are not random, and therefore the likelihood ratio (LR) reduces to only 2 possible probabilities: the probability of a finding if the disease is present divided by that of the same finding if the disease is absent.
- "Odds" have always appealed to epidemiologists and statisticians because of their attractive mathematical properties, but whether clinicians use "probability" or "odds" to describe the likelihood of a particular hypothesis is only a matter of personal taste. Most of us are more comfortable using "probability." Similarly, LRs are nothing more than diagnostic weights, describing how much a particular finding argues for or against a particular hypothesis (and thus changes "probability" or "odds"). The beauty of LRs is that they place multiple findings (or tests) on the same scale, thus allowing clinicians to quickly compare them and identify the most discriminatory tests for (or against) a particular diagnosis and discover the clinical setting in which a finding is most discriminatory.
- The only limit mentioned in my paper is that pretest probability must be between 10% and 90% for my method to work. Therefore, my method should not be used to combine large LRs (e.g., LR = 10) with pretest probabilities less than 10% (e.g., screening tests) nor should it be used to combine tiny LRs (e.g., LR = 0.1) with pretest probabilities near 100% (a clinically unlikely scenario). For virtually all clinical problems of symptomatic patients, however, pretest probabilities do range between 10% and 90%, and if clinicians unfamiliar with odds desire to gauge how much the value of a particular LR changes this pretest probability, the method described in my paper accomplishes this simply and accurately. —**STEVE MCGEE, MD**, University of Washington, Seattle, Wash.
-
- JGIM welcomes your letters and comments. Letters may contain brief commentaries on articles published in the Journal, illustrative case reports, general suggestions for improving the Journal, or other information of interest to readers. Letters to the Editor should not exceed 450 words in length and should be sent to the Editor via e-mail at jgim@jhmi.edu.*