LETTERS TO THE EDITOR

To the Editor:—We agree with the recent article in the Journal by Sippel et al.,¹ which concludes that routine spirometry does not motivate smoking cessation. These results agree with three prior randomized controlled trials.² This is a very important question as the National Heart, Lung, and Blood Institute (NHLBI) has recently considered whether to recommend that our nation screen smokers for obstructive airways disease (OAD) and has also initiated the National Lung Health Education Program.³

Sippel's study advances the case against screening for OAD. In addition, Sippel found a trend that subjects with abnormal spirometry were less likely to quit smoking over the ensuing year. Although long-term trials suggest that abnormal spirometry predicts smoking cessation, prior trials of a year or less duration consistently find smokers with abnormal spirometry are less likely to quit.²

In conclusion, smokers who receive spirometry are not more likely to quit smoking, even if their results are abnormal. As smoking cessation is the only efficacious treatment for asymptomatic OAD, this study adds to evidence against screening. The public health burden of smoking and OAD is large. Our nation should put a tremendous effort into methods to reduce smoking, but not by burdening physicians with the additional responsibility of unjustified screening for asymptomatic OAD.—**BOB BADGETT, MD**, University of Texas Health Science Center at San Antonio, San Antonio, Tex, and **DAVID TANAKA, MD**, University of Colorado Health Sciences Center, Denver, Colo.

REFERENCES

1. Sippel JM, Osborne ML, Bjornson W, Goldberg B, Buist AS. Smoking cessation in primary care clinics. *J Gen Intern Med* 1999;14:670–6.
2. Badgett RG, Tanaka DJ. Is screening for chronic obstructive pulmonary disease justified? *Prev Med* 1997;26:466–72.
3. Petty TL, Weinmann GG. Building a national strategy for the prevention and management of and research in chronic obstructive pulmonary disease. National Heart, Lung, and Blood Institute Workshop Summary. Bethesda, Maryland, August 29–31, 1995 [see comments]. *JAMA* 1997;277:246–53.

In reply:—As the reader correctly concludes, routine use of spirometry was not beneficial in this study, which comprised highly nicotine-addicted and lower socioeconomic status smokers.

JGIM welcomes your letters and comments. Letters may contain brief commentaries on articles published in the Journal, illustrative case reports, general suggestions for improving the Journal, or other information of interest to readers. Letters to the Editor should not exceed 450 words in length and should be sent in triplicate (typed, double-spaced). We cannot guarantee publication and we may edit to improve readability.

You may send your letters to us by mail or fax: JGIM, JHU Division of General Internal Medicine, 1830 East Monument Street, Room 8068, Baltimore, MD 21205; fax (410) 955-0825.

ers. However, extrapolating these conclusions to demographically different populations, which have not been well-studied, may not be appropriate. The role of spirometry as part of a smoking cessation program, combined with nicotine replacement therapy or bupropion, has also not been investigated. Additional clinical trials need to be undertaken to fully elucidate what populations, if any, would benefit from screening spirometry. Lastly, the results of this study do not negate the important role lung function studies have in the diagnosis of lung disease or their ability to provide assistance with therapy, pre-operative risk assessment, and prognosis.¹—**JEFFREY SIPPEL, MD, MPH**, and **MOLLY OSBORNE, MD, PhD**, Oregon Health Sciences University, Portland, Ore.

REFERENCES

1. Crapo RO. Pulmonary-function testing. *N Engl J Med* 1994;331:25–30.

To the Editor:—Doctors Riley and Smith in their recent article¹ recommend a combination of alpha-fetoprotein (AFP) and ultrasound in screening for hepatocellular carcinoma (HCC). After reviewing the literature, in particular the articles to which they make reference, we find no support for the use of AFP as a screening tool for HCC.

The first article referenced, by Oka et al.,² screened 269 patients for HCC with both AFP and liver ultrasounds for 38 months. The sensitivity of an AFP greater than 20 but less than 100 was 39%. The second article, by Zoli et al.,³ monitored 164 consecutive patients with liver cirrhosis for clinical, biochemical, and radiographic evidence of HCC. In only 2.4% of patients did the AFP exceed 100 at time of HCC diagnosis. In a study by Cottone et al.,⁴ of 147 patients with Child's A cirrhosis, 30 (20%) developed HCC. None of those patients had an elevated AFP at time of diagnosis. Of the 30 (20%) diagnosed, only 10 (33%) were eligible for treatment. Of the original 30 diagnoses, there was an overall survival rate of 46%.

Although the evidence does not support the use of AFP as a screening tool, AFP does provide prognostic information. In Oka's study,² 26% of patients with AFP less than 20 developed HCC, while 46% of patients with AFP greater than 20 developed HCC. Other studies supporting AFP as a prognostic tool include the aforementioned article by Cottone.⁴

Thus, we feel that the sensitivity of AFP does not support its use as a screening tool. Additionally, many patients at risk for HCC are so ill that the development of HCC does not markedly affect their overall prognosis. We would recommend obtaining an initial AFP only in those patients, at risk for HCC, who are relatively healthy, and only for prognostic purposes. We would not recommend further AFPs to screen for HCC.—**JANE E. O'RORKE, MD**, and **MARY O'KEEFE, MD**, University of Texas Health Science Center, San Antonio, Tex.

REFERENCES

1. Riley TR, Smith JP. Preventative care in chronic liver diseases. *J Gen Intern Med*. 1999;14:699–704.
2. Oka H, Tamori A, Kuroki T, Kobayashi K, Yamamoto S. Prospective

study of alphafetoprotein in cirrhotic patients monitored for development of hepatocellular carcinoma. *Hepat.* 1994;19:61–6.

3. Zoli M, Magallotti D, Bianchi G, Gueli C, Marchesini G, Pisi E. Efficacy of a surveillance program for early detection of hepatocellular carcinoma. *Cancer* 1996;78(5):977–85.
4. Cottone M, Turri M, Caltagirone M, et al. Screening for hepatocellular carcinoma in patients with Child's A cirrhosis: an 8-year prospective study by ultrasound and alphafetoprotein. *J. Hepat.* 1994;21:1029–34.

In reply:—Dr. O'Rourke and Dr. O'Keefe correctly point out that alpha-fetoprotein (AFP) has a limited sensitivity for detection of hepatocellular carcinoma. The references they refer to give the lower range of sensitivity. The sensitivity of AFP from the literature is more typically 65 to 75%.¹ Given this relatively low sensitivity, it is important to apply this test to a high-risk population. The population that appears appropriate for this type of AFP screening would include patients with cirrhosis. The highest-risk groups among these are the hepatitis C and B patients. In a recent article by Chalasani, however, the other forms of cirrhosis accounted for 20% of the cancers and therefore they made the argument they should not be excluded from screening.² The highest yield of AFP screen comes with the initial value; for example, in Chalasani's study, 22 of 27 tumors detected were detected on the initial screen.

The reason for follow-up studies with AFP is to look for a subsequent rise from baseline. Many patients with hepatitis C and cirrhosis have elevated AFP at baseline, with no detectable lesions on triphasic CT scan. The current thought process is that liver regeneration is occurring in this group, which leads to increased AFP. Follow-up AFP can benefit in deciding which cases need further evaluation by more sensitive tests (such as triphasic CT scan), based on subsequent rises in levels. Because of a relative lack of data in the US population, it is impossible to make firm recommendations for the most cost-effective strategy. For these reasons the individual care provider has to choose a strategy he or she feels is justified by the current available information. In my opinion, a one-time AFP measurement ignores your patient's much higher risk for developing hepatocellular carcinoma and will miss opportunities to detect hepatocellular carcinoma development at early stages.—**THOMAS R. RILEY, III, MD,** *Milton S. Hershey Medical Center, Hershey, Penn.*

REFERENCES

1. Riley TR, Smith JP. Preventative care in chronic liver diseases. *J Gen Intern Med.* 1999;14:699–704.
2. Chalasani N, Horlander JC, Said A, et al. Screening for hepatocellular carcinoma in patient with advanced cirrhosis. *Am J Gastroenterol.* 1999;94:2988–93.