reports four deaths in 30 years attributed to milk-related infection as against a yearly toll of 150 000 deaths from coronary disease in England and Wales. The evidence which began to accumulate in the last 15 or so years about the possible connection between coronary disease and milk consumption comes from several independent studies: epidemiological, immunological, experimental, and clinical.

Virtually every plant we eat—cereals, potatoes, carrots, onions, pulses, tomatoes—contains substances toxic to the human consumer—lectins, saponins, cyanogens, glucosinolates, lathrogens, gossypol, oestrogen mimics, antivitamin, enzyme inhibitors¹—most of which are probably their defensive weapons against their innumerable enemies—viruses, fungi, phytophagous insects, and large herbivores. The fact that we can eat them with impunity is largely because we cook them. Cows' milk is the back door through which many of these substances—originating in grasses, clover, lucerne, soya beans, cottonseed meal, etc.—can reach us in an uncooked form. As there is no blood-milk barrier, many of these substances are likely to pass unchanged into milk and the pasteurisation of the milk is not nearly sufficient to destroy them.

STEPHEN SEELY

University of Manchester Medical School,
Manchester M13 9PT

¹ Liener IE, ed. *Toxic constituents of plant foodstuffs*. New York: Academic Press, 1980.

Laser treatment of port wine stains

SIR,—Although I share the views expressed by Mr J A S Carruth (17 April, p 1195) and in Dr J A Cotterill's leading article (13 March, p 766) about detailed reporting of the beam dosage, the mature port wine stain often varies in colour intensity and skin thickness and may accordingly require quite a varied dosage regimen. My early experience with the Coherent 901 argon laser photocoagulator would suggest that some areas have to be treated repeatedly to achieve a result acceptable to the patient. Incidentally, the use of a local anaesthetic would reduce the energy absorption of the dermal vessels and at times can be withheld.¹ The technique has attracted commercial attention and where the inappropriate CO₂ laser has been used for treating port wine stains the resultant atrophy or scarring could be disastrous. The method has been barely established in the UK and the safety aspects only recently emphasised.² Cannot the BMA urgently induce the DHSS to introduce suitable restrictive legislation?

V K DAVE

Skin Hospital,
Manchester M3 3HL

¹ Arndt K. *Arch Dermatol* 1982;118:220-4.

² Carruth JAS, McKenzie AL. *Clin Exp Dermatol* 1982;7:247-54.

Hormone replacement treatment and benign intracranial hypertension

SIR,—In his report of benign intracranial hypertension occurring in a woman on hormone replacement therapy (5 June, p 1675) Dr John P Sheehan repeats the suggestion that oral contraceptives can cause this condition. The available evidence indicates

that oral contraceptives do not cause benign intracranial hypertension and that any observed association occurs by chance. Of the 30 reported cases in the world literature of benign intracranial hypertension in oral contraceptive users, only 16 meet accepted diagnostic criteria (as set out, for example, by Johnston and Paterson¹) and two of these discontinued the drug one and three months before the onset of their illness. Since there are an estimated 80 million women on oral contraceptives,² and since benign intracranial hypertension predominantly affects women of child-bearing age,¹ a chance association is likely.

If oral contraceptives do have a causative role the rapid increase in use of this type of drug over the past 20 years might be expected to increase the incidence of benign intracranial hypertension. In this region at least there does not appear to have been any such trend.³ Finally, a large cohort study of oral contraceptive users has not revealed any evidence of susceptibility to this condition.⁴

Hormone replacement therapy seems even less likely to cause the condition since the hormone dose is lower, as Dr Sheehan points out. It is probable that this too represents a chance association.

K L WOODS

University Department of Therapeutics and Clinical
Pharmacology,
Queen Elizabeth Hospital,
Birmingham B15 2TH

¹ Johnston I, Paterson A. *Brain* 1974;97:301-12,313-6.

² World Health Organization. *Technical Report Series* 1978;619.

³ Bickerstaff EM. *Neurological complications of oral contraceptives*. Oxford: Oxford University Press, 1975:78.

⁴ Royal College of General Practitioners. *Oral contraceptives and health*. Tunbridge Wells: Pitman Medical, 1974.

Anxiety caused by a short-life hypnotic

SIR,—The reply by Mr K Morgan and Professor I Oswald (12 June, p 1785) to our letter on their report (27 March, p 942) that anxiety was induced by a short-life hypnotic did not answer most of the questions we raised, and indeed raises further difficulties with their study.

There were, of course, two groups of subjects—those who ingested triazolam first and those who ingested loperazolam first—and we would point out that neither the nine out of 12 subjects nor the seven out of nine who reported increased anxiety supports the rejection of a null hypothesis. Statistically such ratios do not reach an acceptable level of significance and so do not indicate increased anxiety in the two groups separately even by the third week of ingestion.

However, our main criticisms are concerned with design and graphical presentation. There was no equivalent period (three weeks) of placebo ingestion. There could have been an expectation of an improvement in anxiety with the ingestion of a tablet, and if this was not fulfilled the subjects could have reported increased anxiety even though the medication was devoid of such activity. Furthermore, the absence of an equal period of placebo ingestion works against double-blindness. The same period (three weeks) at the same time in each of the five weeks of each study (four in all) was used for drug ingestion only, and so even single-blindness may not have been preserved later on.

In their graphic presentation the authors

implied an increase in anxiety with triazolam by comparison with a drug (loperazolam) which had clear-cut anxiolytic activity. We would emphasise, once more, that an inactive compound could easily and equally have been said to have induced anxiety if it had been compared in this way with an anxiolytic drug.

Our criticisms are essentially those of Iversen and Mackay¹ concerning a previous study by Bržinová and Oswald.² We can do no better than to repeat that the validity of the conclusions which one may draw from any clinical trial is directly related to the amount of care taken to exclude the influence of placebo reactions and the preconception of response on the part of either patient or physician. There are serious reservations concerning this study by Morgan and Oswald and about the misleading way in which the data have been presented.

A N NICHOLSON

A J BELYAVIN

B M STONE

M B SPENCER

Royal Air Force Institute of Aviation Medicine,
Farnborough, Hampshire

¹ Iversen LL, Mackay AVP. *Br Med J* 1972;ii:766.

² Bržinová V, Oswald I. *Br Med J* 1972;ii:431.

Laboratory accuracy

SIR,—Most clinical chemists are obsessional in their approach to the quality control of their assays and will resent the suggestion by Dr John Crammer (19 June, p 1852) that clinicians, faced with the problem of unexpected variation in the concentrations of lithium in a patient's serum, should check the laboratory's competence by means of hidden duplicate samples. When a clinician does suspect that a laboratory is producing erroneous results the appropriate first action should be to discuss the problem with the head of the department concerned.

A R W FORREST

Royal Hallamshire Hospital,
Sheffield S10 2JF

Families in high-rise flats

SIR,—Your leading article "Families in high-rise flats" (20 March, p 846) may have overstated the case for flat dwelling causing mental illness. Comparing families in flats with those in houses is not comparing like with like. Mothers in houses are older, have older children, and are better off financially. Any increase in the prevalence of mental illness of flat dwellers is just as likely to be due to differences in maturity and social class as to the accommodation itself. In my study,¹ after matching for age, social class, and other variables I found that the difference in the incidence of mental illness disappeared. Two other studies,^{2,3} referred to in your editorial, support my findings. Fanning,⁴ whose work is often quoted to support the opposite point of view, did not match these variables. It is also worth noting that the accommodation he studied was all low rise—of four storeys or lower. The problems in high-rise flats may be quite different.

Nevertheless, there is a small but important subgroup of flat dwellers which has a slightly increased risk of mental illness.⁵ Those of neurotic personality are more likely to develop

symptoms when living in flats than in houses. Flats lack privacy and are noisy, and their occupants feel isolated and lonely and have a poor indoor social life.⁶ It is not surprising that some of those of neurotic personality cope badly with this way of life and use it as a convenient explanation for their unhappiness. Although flat dwelling for the majority does not cause mental illness, there is no dispute that flats are socially inferior and that the quality of life for most families would be improved if they could be avoided altogether.

N C MOORE

RAF Hospital Wegberg,
BFO 40

- ¹ Moore NC. *Br J Psychiat* 1974;125:500-7.
² Richman N. *Dev Med Child Neurol* 1974;16:53-8.
³ Ineichen B, Hooper D. *Social Science and Medicine* 1974;8:369-74.
⁴ Fanning DM. *Br Med J* 1967;iv:382-6.
⁵ Moore NC. *Br J Psychiat* 1976;128:259-61.
⁶ Moore NC. *Publ Hlth Lond* 1975;89:109-15.

Quality of diet and blood folate concentrations

SIR,—We have previously reported the red cell folate concentrations in women at increased risk for neural tube defect during prospectively studied pregnancies.¹ We wish to emphasise the close relationship between folate concentrations in the non-pregnant state and quality of diet for all the women included in that study. Women resident in Glamorgan and Gwent who had a pregnancy complicated by open neural tube defects (anencephaly, encephalocele, and spina bifida)

general pattern for meals and a check list designed to show any deficiency of foods rich in folic acid consumed during the average week and listing protein foods, dairy products, vegetables, fruits and salads, brown and white bread, confectionery, and soft drinks. The diets were classified into three broad groups. Good diets were those with a good intake of all essential nutrients, including protein, vitamins, and folic acid, calcium, and iron, and no excessive amounts of sweets, chocolates, or soft drinks. Poor diets were those with a deficient intake of protein, vitamin and folic acid foods, calcium, and iron, and generally an imbalance with excessive amounts of refined carbohydrates.² Fair diets were intermediate and were those containing less than the optimum quantities for non-pregnant women of protein and foods containing vitamins and folic acid, often only small quantities of fresh fruits and vegetables, with or without an excess of refined carbohydrates.

Blood samples for folate estimation were taken, but the conditions under which they were taken (time of month and day and relationship to meals) could not be standardised. Serum and red cell folic acid concentrations were measured by a modified lactobacillus method.^{3,4}

Of the 905 women visited, 443 were not pregnant at the time but were planning to have further pregnancies and were willing to co-operate. Of these, 11 had to be excluded because the patient could not be positively identified from the labelled blood specimens, and blood samples were not obtained, did not arrive at the laboratory, or arrived haemolysed in a further 64.

The accompanying tables show a highly significant association between the dietary rating and red cell folate and serum folate, but the latter was less marked. The red cell folate values, generally regarded as a good indicator of folate concentrations in the body, are of interest because they suggest not only that those women on the poorer diets have lower folate concentrations but that the

alone in the prevention of neural tube defects due to be launched this autumn.

K M LAURENCE

Department of Child Health

NANSI JAMES
H CAMPBELLDepartment of Medical Statistics,
Welsh National School of Medicine,
Cardiff CF4 4XN

- ¹ Laurence KM, James N, Miller M, Tennant GP, Campbell H. *Br Med J* 1981;282:1509-11.
² Laurence KM, James N, Miller M, Campbell H. *Br Med J* 1980;281:1542-4.
³ Hoffbrand AV, Newcombe BFA, Mollin DL. *J Clin Pathol* 1966;19:171-81.
⁴ Tennant GB, Withey JL. *Medical Laboratory Technology* 1972;29:171-81.
⁵ Smithells RW, Sheppard S, Schorah CJ, et al. *Arch Dis Child* 1981;56:911-8.
⁶ Smithells RW, Sheppard S, Schorah CJ, Seller MJ, Nevin NC, Harris R, Read AP, Fielding DW, Walker S. *Lancet* 1981;ii:1424-5.

Blood pressures that fall on rechecking

SIR,—Dr S Mann and others (15 May, p 1477) state that "many studies have failed to demonstrate a positive relation between variability and mean pressure." They therefore imply that the apparently highly significant correlation between mean blood pressure and its standard deviation shown by the Framingham study¹ is incorrect and that these other studies, in contrast, all show a highly significant negative correlation between mean blood pressure and its coefficient of variation. The authors should be well aware that when one replies to a letter or an article to point to the existence of evidence which conflicts with that presented originally, it is customary to cite one's sources. If there is definite evidence that the coefficient of variation is significantly negatively correlated with mean blood pressure they should produce it. If their statement was based on investigations which were unable to show any significant correlation between blood pressure and its standard deviation or its coefficient of variation, these results do not conflict with the evidence cited in the Framingham study and should not have been mentioned.

P M GAYLARDE

Department of Dermatology,
Royal Free Hospital,
London NW3 2QG

- ¹ Kannel WB, Sorlie P, Gordon T. *Circulation* 1980; 61:1183-7.

TABLE I—Quality of diet and concentrations of folate in blood of 368 women with history of pregnancy with a fetal neural tube defect

Quality of diet	No of subjects	Mean serum folate (±SEM) (in µg/l)	Mean RBC folate (±SEM) (in µg/l of red cells)
Good	62*	8.87 ± 1.03	295 ± 16.6
Fair	181	5.39 ± 0.16	237 ± 17.6
Fair but disordered	46	5.66 ± 0.42	232 ± 15.5
Poor	79	4.88 ± 0.28	197 ± 7.8
All women	368	5.90 ± 0.23	238 ± 5.4

*This includes one woman on a good diet with an anomalously high serum folate level of 51.4 µg/l whose red blood cell folate was 461 µg/l of red cells. The significance of the differences remain unchanged if this case is excluded.

TABLE II—Significance of differences between dietary grades

Comparison	Serum folate			RBC folate		
	Mean difference	t	p	Mean difference	t	p
Good v fair	3.48	3.06	<0.01	58	9.92	<0.0001
Good v poor	3.99	3.45	<0.001	98	19.83	<0.0001
Fair v poor	0.51	1.59	>0.1	40	7.92	<0.0001

between 1954 and 1969 were traced through malformation registers, maternal and paediatric records, local authority records, and other sources. Those under 35 at the time of the study were visited in their homes by medically qualified field workers as part of a prospective study of diet² and possible folic acid depletion in early pregnancy. During the home visits a questionnaire was completed giving details of the woman's diet when she was not pregnant, and blood was taken for folate estimation only from those women planning further pregnancies and willing to co-operate in the prospective study.

A simple diet sheet was used that provided a

dietary ratings used in the South Wales studies of prevention of recurrence of neural tube defects after preconceptional dietary counselling and also after supplementation with folic acid are valid. The association between the quality of the diet and the serum folate concentration is slightly less marked, as might be expected, for this is a measure only of folate in transit and would be subject to variation due to the content and time of the last meal, etc.

These associations are of relevance to the ongoing trials^{5,6} of periconceptional supplementation with Pregnavite Forte F and to the large multicentre randomised controlled MRC trial of Pregnavite Forte F, a multivite preparation without folic acid, and of folic acid

A happier old age in Denmark?

SIR,—Let us give thanks for the timely and farsighted synthesis from Dr Colin Godber (12 June, p 1729). Let us hope and pray that those in position to take action will understand his quiet sensibility and accept his guidance to take an initiative which has been delayed too long.

Our clinical practice, tempered by the results of research work carried out in part III homes of our district,¹ confirms the view that the best way forward from our present gross inadequacy is by developing existing part III homes as comprehensive centres of care for the elderly of the immediate locality. Beds, day care, and co-ordination of services in the community could be provided from the single "nucleus."² This is very similar to the Danish model in its shape, and we would suggest a capacity "twice [that] of our part III and long-stay