Conference Report

Alcohol update

RICHARD SMITH

Anatomists, the cynics at medical school used to say, compensated for never discovering anything new by repeatedly changing their nomenclature. Changing names and definitions, reassessing priorities, assimilating mistakes, and reanalysing old data may, however, be important and may be one way for knowledge to advance. In such ways rather than in the publishing of important new data, judging from last week's Federation of Alcoholic Rehabilitation Establishments symposium in York, has knowledge on alcohol problems advanced in the last year since I did the research for a series of articles on alcohol problems.¹

Indeed, the one crucially important paper published in the last year is one that seems to have undermined completely the germinally important work of Sobell and Sobell, the doyens of the controlled-drinking school. They published results in the early 1970s²⁻³ showing that using behaviour psychotherapy and aiming for controlled drinking rather than abstinence they could achieve better results than people using other techniques. A paper published in *Science*⁴ this year has, however, looked again at the subjects from the Sobells's study and showed that far from being controlled drinkers most were dead, drinking heavily, or abstinent. The Sobells have denied emphatically the imputation that their results were fixed, but although independent inquiries have been established and nothing is yet proved the feeling at the FARE symposium was that their work would probably be discredited. Would this be a fatal blow to the controlled-drinking movement? Although the Sobells's work is the main star in the controlled-drinking firmament, people at the conference felt that there was too much other work showing that controlled drinking was a viable alternative to abstinence for the whole technique to be discredited.

Other treatment controversies

A more British controversy that was considered at the conference is a good example of the problem of nomenclature—is there such a thing as the alcohol dependency syndrome? Bill Saunders, director of the Alcohol Studies Centre in Paisley, brought up this question in his review of current treatment issues. Professor Griffith Edwards, the most influential figure in British alcoholology (yes, there is such a word, and it's here to stay), has argued strongly in his recent book⁵ that there is such a syndrome and that it is important for those concerned with treating alcohol problems to recognise it.

The term has been around since 1976⁶ but seems to go against the fashion, which is for the abandoning of the concept that there is any discrete group of people (alcoholics or whatever) who are especially vulnerable to alcohol. The fashionable idea is more that given enough to drink we will all run into serious alcohol problems and that if we drink less many of these problems will

disappear. Griffith Edwards is arguing, however, that after drinking so much some people move into a phase (the alcohol dependency syndrome) where they move rapidly towards disaster and find it very much more difficult to abandon alcohol; one important part of the concept is that for such people controlled drinking may not be a treatment option.

Some of those at the symposium felt that the pushing of this idea was a rearguard action by doctors to keep alcohol problems in their patch, while others, including some doctors, thought that even if the syndrome did exist it didn't really change much as only a minority of patients would fit the syndrome anyway. The controversy is another one that is unresolved but that will not go away.

Bill Saunders in his excellent paper identified other important treatment controversies. One important one that has dogged alcohol treatment since it ever began is the conflict between the therapists who are "craftsmen" and those that are "scientists." The craftsmen are those who learn their craft at their master's knee after a long apprenticeship, and their main characteristic is that they do not criticise the master or their craft. They know they are doing good and they have no need to prove it. The "scientists," however, are characterised by their critical approach to their activities. Bill Saunders's sympathies lay with the scientists, and he thought that the time had come for the craftsmen to prove themselves. He quoted from our recent leading article by Dr Jonathan Chick (3 July, p 3): "Treatment may actually make some alcoholics worse by sheltering them from the consequences of their drinking or fostering passivity. The gauntlet has been thrown down: the onus of proof is now on therapists who try to provide more than just succour to show that they are necessary agents in the recovery of certain of their patients and to show which these are."

A closely related issue is the relative importance of the characteristics of the patient, the therapist, and their relationship. "Scientists" are increasingly convinced that what matters most are the characteristics of the patient: if he is married, employed, and "socially stable" a man is likely to recover from alcohol problems, whereas if he is not he is likely not to. The "craftsmen" resent this assertion and say that the therapist's characteristics are vitally important (some said that his motivation is more important than the patient's), but again evidence is lacking. Many of the psychologists at the conference took a different line and said that empathy between the therapist and the patient, which they say can be reliably and reproducibly measured, was the most important predictor of outcome. Bill Saunders concluded this part of his talk by saying that Anthony Clare's much quoted statement of 1977 still stood. "It remains to be established," he said, "that minimal forms of treatment are any more effective in modifying alcoholic patterns of drinking than no treatment at all."

That many people who drink heavily and experience problems can spontaneously reduce their drinking to unarmful levels is, however, well established, and some have suggested that it is important to study how these people stop drinking. Various papers, including Vaillant's long-term study of Boston teenagers⁷

(which I⁸ and Dr Chick referred to), have looked at this issue, and an important one from Tuchfield was published last year.⁹ People who spontaneously have overcome severe alcohol problems were found by media advertising, and the reasons for their recovery were analysed: 17 had reduced their drinking after an illness or accident; 15 because of extraordinary events such as humiliation or attempted suicide; 13 because of religious conversion or experience; and 11 because of financial problems. These factors are similar to those identified by Vaillant.

Compulsory treatment of alcoholics

An old idea for responding to alcohol problems—compulsory treatment—is resurfacing around the world,¹⁰ including Britain,¹¹ and was debated at the conference. In about 10 cities in Britain young drunken offenders are being put on probation orders and as part of the order are required to attend courses on alcohol education. Ian Robertson, a senior clinical psychologist from Edinburgh, described the course he had run in Dundee, and Malcolm Smith, a probation officer, described his course in Corby. One part of the debate was just how compulsory these courses were: all of the “students” had committed an offence and could if they wanted opt for traditional sentencing rather than the course. So in one sense they did have a choice, but others felt that when the alternative was prison or some other sentence there was no real choice. Once they have opted to attend the course the “students” have to attend or they are returned to the court.

The course in Dundee consists of six one-and-a-half hour sessions; and about eight offenders under 25 who have alcohol problems but not severe ones and who have no other psychiatric problems attend. In the first session rapport is established. Thereafter, topics covered include: basic information about alcohol and the body; high- and low-risk ways of drinking; rules for safer drinking; alternative ways of using leisure time; ways of coping with a craving for drink; and the problems that may arise with severe alcohol dependency. Malcolm Smith's scheme was similar but lasted longer.

None of the therapists who had undertaken to run such schemes had done so without much soul searching and discussion, argued Robertson and Smith. They thought that it was much better to try in some way to help these young men rather than do nothing because of “high falutin” philosophical and ethical worries: “A libertarian conscience is a luxury on a Dundee council estate,” said Ian Robertson. Doug Cameron, a psychiatrist from Leicester, thought differently, however, and was convinced that compulsory treatment was an ominous development. Members of the caring profession by running such courses were, he argued, becoming “agents of social control” rather than therapists. People had the right to damage themselves and refuse treatment, and, anyway, trying to treat somebody with alcohol problems against his will was an impossible task.

People in the audience were divided on the issue. Some took the pragmatic line that these young offenders were going to be punished anyway and better to attend a course that might help rather than go to prison, which certainly would not help them. Others found the ethical implications too horrible and were against the whole idea. Many delegates agreed, however, that if such courses were to continue they should be subjected to controlled trials and hard evidence should be produced that they worked. Ian Robertson also made what I thought was a very important point when he said the Government should not be allowed to make such programmes part of its policy and then claim that it was doing something positive about alcohol problems.

Prevention in the North-east

In a much less sinister session Anthony Thornley, a psychiatrist from Newcastle, looked back at the lessons to be learnt

from the Health Education Council's campaign on alcohol in the North-east. The campaign had been in three stages—in 1974, between 1977 and 1979, and in 1980. The first two stages included many mistakes, but Anthony Thorley thinks that the present campaign should achieve much more. The first two parts of the campaign made a huge impact in the North-east but left people more confused than knowledgeable, and the 1974 campaign led to hundreds of people coming forward for help with their alcohol problems when none was available.

Many of the failures, Anthony Thorley thought, were due not just to the newness of the art of health education but also to the fact that a London-based organisation and a London-based advertising agency were designing a campaign for an area about which they knew next to nothing. Their clever but obscure messages were lost on the people of the North-east. The 1980 campaign used a local advertising agency, a local man (Peter Bellamy) whose accent was acceptable to people all over the North-east (for Newcastle and Gateshead are divided by more than a river, and Sunderland, as Anthony Thorley kept reminding his audience, is part of Denmark as far as many North-easterners are concerned), and a clear message—“Stick to your two or three pints two or three times a week—don't spoil a good thing.” The early analyses of this campaign, which are to be published very soon, suggest that the message has reached many North-easterners and many have remembered the message.

So in a year thinking on neither treatment nor prevention has changed much: nobody is certain that treatment is very effective, and preventive campaigns are achieving only the most modest results for large expenditure of time and money. Politically, things have not progressed very much either: the Government's response to the suggested reorganisation of the antialcohol groups¹² is still anxiously awaited; the alcohol equivalent of Action on Smoking and Health has not been launched; and although the Government “think tank's” report on alcohol policy has been published in Sweden¹³ the Government continues to ignore its recommendations.

References

- Lock S, ed. *Alcohol problems*. London: BMJ, 1982.
- Sobell MB, Sobell LC. Alcoholics treated by individualised behaviour therapy: one-year treatment outcome. *Behav Res Ther* 1972;11:599-618.
- Sobell MB, Sobell LC. Second year treatment outcome of alcoholics treated by individualised behaviour therapy: results. *Behav Res Ther* 1976;14:195-215.
- Pendery ML, Maltzman IM, Jolyon West L. Controlled drinking by alcoholics? New findings and a reevaluation of a major affirmative study. *Science* 1982;217:169-75.
- Edwards G. *The treatment of drinking problems*. London: Grant McIntyre, 1982.
- Edwards G, Gross MM. Alcohol dependence; provisional description of a clinical syndrome. *Br Med J* 1976;i:1058-61.
- Vaillant GE, Milofsky ES. Natural history of male alcoholism. IV. Paths to recovery. *Arch Gen Psychiatry* 1982;39:127-33.
- Smith R. Treating alcohol problems: making ends meet. *Br Med J* 1981;283:1043-5.
- Tuchfield BS. Spontaneous remission in alcoholics. Empirical observations and theoretical implications. *J Stud Alcohol* 1981;42:626-41.
- Anonymous. Is forced treatment the future? *Legal Issues* 1981;November: 1.
- Hamson C, Knapman E, Mallett J, Smith M. *Corby alcohol therapy group 1979-81. Report*. Northampton: Northamptonshire Probation and Aftercare Service, 1982.
- Department of Health and Social Security. *National voluntary organisations and alcohol misuse*. London: DHSS, 1982.
- Brunn K, ed. *Alcohol policies in the United Kingdom*. Stockholm Sociologiska institutionen, 1982.

Can indoor flash photography of young babies damage their retinas?

There is no known risk provided flashes are not applied rapidly (high-speed) in succession.—R A WEALE, professor of visual science, London.