Commentary: Patient's view

Kenneth Neville with the support of Mark Fox

Oesophageal
Laboratory,
St Thomas'
Hospital, London
SE1 7EH
Mark Fox
specialist registrar

Correspondence to:
M Fox
markfox@
doctors.org.uk

As a child I remember eating well and enjoying my food, although I did get heartburn now and again. That all changed in 2001 when I started to bring up what I ate. At first this was occasional, but then my stomach settled into a new routine. On eating even small amounts I felt unpleasantly full, then I had to belch and the food came back to my mouth. I usually spat this out, but sometimes I swallowed it again. If I tried to eat larger amounts my stomach hurt and I had to vomit everything up.

In spring 2004 I was sicking up more and more, but my stomach just felt fuller and fuller. I was admitted to hospital. At first everybody was sympathetic. However, after a while I could sense that the doctors were becoming frustrated at not reaching a diagnosis and then I felt that they began to get frustrated with me as a person. In the end I was sent home without an explanation for my problems. I got the impression that the doctors thought that I was wasting their time.

At home I forced myself to eat, but I was keeping down only a third of my food. At Christmas I had to keep leaving the dining room so I could be sick without upsetting my mother. I put off going to hospital until the New Year because I knew nothing would happen until after the holidays. Back on the ward I had a lot of

investigations. Every time the person doing the test would come to me and say, "Everything's normal"; they always wondered why I lost my smile. I was begging for something to come back positive so I could get on with my life. The doctors didn't seem to be able to agree on how to deal with me. I asked for a second opinion.

At St Thomas' I kept on losing weight and became so weak that I could not get out of bed. Finally, I had an investigation that measured what happened in my stomach after I ate. I always wondered why nobody had done that before. I had my doubts when I was told that the problem was under my control. With the doctors' help I became aware of the process. I began to feel the tension building up in my stomach after eating and tried to keep the food down. At first it was hard to distinguish wind and food coming back up, but suddenly I got the hang of it. Once I was in control it encouraged me to continue. I still remember the excitement when I moved from "underweight" to "low normal" on the weight chart. At home I am getting stronger and have only allowed food to come back up a couple of times when I ate too much. I'm really grateful and glad because I can now get my life back.

Commentary: Wide angle thinking helps patients with distressing illness

Ed Peile

Institute of Clinical
Education, Warwick
Medical School,
University of
Warwick, Coventry
CV4 7AL
Ed Peile
*professor of medical
education*
ed.peile@
warwick.ac.uk

Kenneth Neville presents interesting problems for specialists and generalists alike.¹ At the time of the case presentation, we were told about a young man with distressing symptoms of postprandial vomiting leading to weight loss and weakness. The local hospital had diagnosed gastro-oesophageal reflux disease, and a battery of other routine investigations had given negative results. We were presented with several possibly relevant

psychological factors, and it was mentioned that his asthma had been difficult to control. Readers were invited to consider the differential diagnosis and to suggest the most helpful course of management for the family doctor to take when Kenneth re-presented shortly before Christmas.

How often have I heard the expression, "This patient has been fully investigated at..."? The truth is that no patient is fully investigated. Investigation, like diagnosis, is a means to an end, and the end is that a patient is able to live as comfortably as possible with disease burden, and hopefully that the burden will be relieved altogether. On bmj.com family doctors rose to the challenge and showed their ability to think widely across the range of possibilities in the differential diagnosis.² The clinical reasoning processes were both inclusive (what do we need to consider that has not been thought about so far?) and exclusive (what can we safely rule out on the basis of the evidence presented?). Most doctors resisted the temptation to think in terms of either psyche or soma, instead considering emotional and psychological factors alongside physical illnesses.

Referral decisions

I think this is the stuff of general practice, working with the patient, starting from the point that distress persists, and showing a willingness to use professional

DAVID PARKERS/SL

Doctors need to look at all angles