alcoholics with cirrhosis histologically unrelated to their alcoholism show a very high percentage (9/15, 60%), significantly higher than that of patients with cirrhosis and alcoholic histological findings (12/45, 27%) ($p=0.05$). This last percentage is higher than that found among alcoholics with no liver disease (4/40, 10%) but the difference is not statistically significant. Among the healthy population 5% are positives.

These results show a high prevalence of hepatitis B virus active infection in alcoholics with cirrhosis without histological signs typical of alcohol-induced liver disease. Although the percentage is also high in the subgroup of alcoholic cirrhosis, it is not significantly different from that in alcoholics with normal livers but the difference from the percentage in healthy controls does reach significance ($p=0.001$). Two conclusions can be drawn: (1) In some alcoholics liver disease may be due to hepatitis B virus. (2) In alcohol-induced liver disease it is possible that hepatitis B virus may play a part.

VICTOR VARGAS
JOSÉ D PEDREIRA
RAFAEL ESTEBAN
JOSÉ M HERNÁNDEZ
JAIME GUARDIA
RICARDO BACARDÍ

Departments of Internal Medicine
and Haematology,
Ciudad Sanitaria de la Seguridad Social,
Universidad Autónoma,
Barcelona-35, Spain

Brucellosis

SIR,—I was interested in the comments on chronic brucellosis in your leading article (11 April, p 1180). Working in an area where brucellosis has been a common scourge for many years it has been my experience that chronic brucellosis occurs frequently.

In 1976, when the disease was rampant here and this practice produced more than 10% of the total disease for England and Wales (14 cases out of 129), we decided to survey the local farming population. In this survey we found that 94 farmers out of 355 (27%) had an antihuman globulin level of 1/80 or above.

Eradication in animals is taking place here now and we plan to repeat our survey in 1987 to see what effect eradication has had on our farming population. Meanwhile we continue to see patients with recurring symptoms of chronic brucellosis. These include depression, arthritis, pancreatitis, and myalgia; and I have two patients whose depressions respond very well to treatment with antibiotics. I would consider brucellosis to be a greater source of chronic ill health in a rural community than has hitherto been expected.

I frequently find it difficult to treat this chronic condition. Those who are originally treated with streptomycin in combination with tetracycline appear to have fewer relapses, but occasionally antibiotics are required for very long periods to relieve symptoms.

I hope that eradication of the disease in animals will rid us of this problem in time, but I expect that the chronic and subacute conditions will continue to be a source of diagnostic difficulty for many years to come.

ANGUS ARMSTRONG

Morpeth,
Northumberland NE65 7TE

Insulinomas

SIR,—Your leading article on insulinomas (21 March, p 927) mentions only two methods for investigating cases refractory to diagnosis by the measurement of glucose and insulin during a fast. There is striking omission of reference to the paediatric experience in this regard.

The diagnosis of insulinomas and other causes of hyperinsulinism in childhood can be assisted by measuring, in addition to glucose and insulin, the plasma β -hydroxybutyric acid and free fatty acids during the fasting study.¹ Further confirmatory evidence is obtained by administering glucagon parenterally at the end of the fast, and noting the rise in blood sugar over the next half hour.²

It is believed that these methods, so helpful in the diagnosis of childhood insulinomas, are applicable to adult cases.

KENNETH R LYEN

Division of Endocrinology and Diabetes,
Children's Hospital of Philadelphia,
Philadelphia, Pennsylvania 19104,
USA

¹ Stanley CA, Baker L. *Pediatrics* 1976;57:702-11.

² Finegold DN, Stanley CA, Baker L. *J Pediatr* 1980; 96:257-9.

Assessing elderly incontinent patients

SIR,—We read with interest the article by Dr C M Castleden and colleagues (4 April, p 1103) and the correspondence (25 April, p 1396) relating to this and to our own paper on a similar subject (21 March, p 940). We feel, however, that several further points should be made.

We would firstly refute entirely the suggestion made by Dr Castleden and his colleagues that we have applied double standards of care. Our standard for all patients is to achieve the most accurate possible diagnosis with the minimum of investigation. While we certainly accept that urodynamic investigation is necessary for an accurate diagnosis in young patients, we showed in our paper that in the elderly 95% accuracy in determining therapy can be obtained with our algorithmic process. The fact that urodynamic investigation allows 100% accuracy of diagnosis is not an argument for the wholesale investigation of elderly incontinent patients, since the available treatments are less than 100% effective. Dr Castleden and others are indeed fortunate if their local general practitioners and consultant colleagues do carry out some initial assessment of incontinence prior to referral; this has certainly not been our experience, as we showed in our article that 25% of our patients had either infected urine or faecal impaction at the time of referral. Indeed, we would see the main place of our algorithmic approach as being in the initial assessment, at a primary care level, and not necessarily in specialist urodynamic practice. We take Dr Eastwood's point that a small number of patients with acontractile bladder may present neither with retention nor with stress incontinence, and that with our algorithmic method would be treated by antispasmodic drugs. Although it is our practice to warn such patients of the possibility of voiding difficulties, it has been our experience (as stated in our article) that such patients do not go into retention but rather respond well—subjectively at least—to such therapy.

Concerning the paper by Dr Castleden and others, we would firstly question much of their terminology. Their criteria for normality are undefined; and since two of their "normal" patients had bladder capacities of less than 100 ml, and three or more had pressure rises exceeding 20 cm H₂O on filling, we are dubious about the validity of this group. "Atonic" and "irritable" as specified in their paper seem to be largely clinical diagnoses with no clearly defined urodynamic criteria. The

failure to recognise urethral sphincter incompetence (genuine stress incontinence¹) as a cause of incontinence in the elderly we consider a major omission; in 40% of our female patients this was a significant contributory factor.

Since 58% of Dr Castleden's patients required some aid to mobility, we would question his conclusion that incontinence in the elderly is not associated with immobility and cerebral disease. We feel quite strongly that even relatively minor limitations in mobility in the elderly are quite sufficient to produce incontinence in a patient who might otherwise suffer only frequency or urgency. We were surprised to find that the clinical details presented are limited to neurological and musculoskeletal aspects. Although the presence of faecal impaction was noted, no information is given as to other pelvic findings. The demonstration of palpable bladder enlargement and the physical sign of stress incontinence have been shown to be of considerable importance in the assessment of incontinence, and we feel that they should be within the scope of any competent district nurse.

Concerning their urodynamic data, in their own words "no bladder capacity or pressure appeared characteristic of any condition, including normality." There would seem only two possible conclusions from this statement: either the conditions referred to are insufficiently well defined to be of relevance to the assessment of incontinence or patients could be equally well allocated to these categories without the use of cystometry. Dr Castleden and his colleagues do not specify what treatments they have employed in their study, or show how their urodynamic data have altered management.

We feel therefore that their conclusion that urodynamic investigation is necessary in elderly incontinent patients before treatment is unproved, a sentiment apparently echoed by Dr Eastwood in his letter. The final statement of their paper could perhaps be modified: "Treatment of urinary incontinence in the elderly without prior cystometry may be likened to treating cardiac arrhythmias without electrocardiography, though much useful information may be obtained from prior palpation of the pulse."

PAUL HILTON
STUART STANTON

Urodynamic Unit,
St George's Hospital Medical School,
London SW17 0RE

¹ Bates P, Bradley WE, Glen E, et al. *J Urol* 1979; 121:551-4.

SIR,—The clinical and urodynamic evaluation of incontinence in elderly patients is more difficult and time consuming than in younger people. Dr C M Castleden and colleagues (4 April, p 1103) have performed a praiseworthy study of this group of patients, who often suffer much from their disability and whose treatment provides a significant challenge. We would like to comment on certain aspects of his communication.

It is unfortunate that the symptom findings in this study were not presented. A prediction of urodynamic findings can often be made by careful symptom analysis.¹ Although communication with elderly patients may prove time consuming, eliciting the symptom complex of frequency, nocturia, and urgency of micturition will frequently correlate well with the urodynamic findings of detrusor instability and perhaps avoid the necessity for urodynamic evaluation.

It is important to distinguish between male and female patients as the aetiology of their incontinence may differ. In the men, although incontinence of urine in this study was shown to be related to instability in 22 of 25 patients, no mention of outflow obstruction was made; but it is likely that a number would have evidence of outflow obstruction, which if treated might well render them stable. Urethral pressure profilometry