NICE issues new guidance to improve the treatment of bipolar disorder

Adrian O'Dowd *London*

The comedian Stephen Fry, who has bipolar disorder, interviews others with the condition in a BBC2 documentary, *The Secret Life of the Manic Depressive*, to be screened in the autumn

New national guidance for England and Wales encourages doctors to spot and treat bipolar disorder sooner and more accurately.

Bipolar disorder, formerly known as manic depression, affects around 1% of the population and takes an average of eight years to be correctly diagnosed. It is associated with the

highest suicide rate of any mental illness—one in seven people who are untreated commit suicide—but it is treatable.

The National Institute for Health and Clinical Excellence launched its clinical guideline on the condition last week, together with the National Collaborating Centre for Mental Health.

The guideline sets out the criteria for referring patients to specialist psychiatric assessment and treatment. It also details the options for drug treatment for people with the disorder and recommends annual physical health checks for such patients as well as monitoring their drug use.

Failure to diagnose bipolar disorder is the result of a lack of knowledge about the condition, said Richard Morris, professor of psychiatry at the University of Nottingham.

Speaking at the launch of the guideline this week, Professor Morris said, "The failure is not with doctors but comes from a lack of guidance on this area within the NHS itself. There should be more of a focus on this area during doctors' training stage.

"We want to sharpen the practice of people and encourage doctors to think [that] if one drug does not seem to be effective for a patient then there are others and they should not give up."

Professor Morris said that someone with bipolar disorder diagnosed at age 30 who lives to 70 could experience 20 years of ill health and 14 years unable to work and expect a life span shortened by nine years. People with bipolar disorder tend to have heart disease problems, he said, due to the stress on the body during mania.

Stephen Pilling, a consultant clinical psychologist and joint director of the National Collaborating Centre for Mental Health, part of the Royal College of Psychiatrists, said, "More needs to be done to improve awareness, identification, and recognition of this problem so that appropriate treatments are prescribed and symptoms can be better controlled."

At the guideline's launch Robert Westhead, who has the condition, spoke of his experience. He said it took him 14 years to find the right combination of drugs to work for him and that he attempted suicide a few years ago.

"I hope this guidance will shorten the torturous journey for others that I have had to make to get the right treatment, and I hope it will save lives," he said. □

The Management of Bipolar Disorder in Adults, Children and Adolescents in Primary and Secondary Care can be seen at www.nice.org.uk.

Senior doctors criticise government report on allergy treatment

Paul Dinsdale *London*

Doctors' leaders have criticised a report on the treatment of allergies by the Department of Health in England for not proposing strong enough action to improve services.

The report places the onus on family doctors, who, it says, need to be better trained in the detection and treatment of allergies, and it proposes that they work with the medical royal colleges to this end.

The Department of Health's report follows a year long review, which was prompted by calls from the parliamentary health select committee in 2004 for better services for allergy.

Around a third of the population have some form of allergy

at some point in their lives, and every year about three million people in England present to primary care with conditions that may be allergic in origin.

The incidence of allergies has been increasing over the past 30 years. In 2004 about 70 000 people were admitted to hospital for asthma, says the report, although many cases were not allergic in origin, and 2400 people, mostly children, were admitted for eczema. There were also more than 3000 admissions for acute allergic conditions (anaphylaxis), and the number of these seems to be on the rise.

Although it is difficult to accurately assess the number of deaths that are directly attribut-

able to allergy, anaphylaxis kills between 10 and 20 people every year, although many cases are not recorded as such on the death certificate.

The minister for care services, Ivan Lewis, said: "We need to ensure that GPs and others in primary care have clinical knowledge and support systems in order to spot allergy in the early stages, so that an effective management plan can be offered at the start."

But Pamela Ewan, secretary to a working party of the Royal College of Physicians that produced a report on allergy in 2003, said: "The review has tried to address the issues, but their recommendations will hardly scratch the surface of the problem. This needs central direction, and funding cannot be left to devolved local systems, as the [health department] proposes. That has been the situation for the last decade, and patient care has not improved."

Graham Archard, vice chairman of the Royal College of General Practitioners, said that much better educational packages for GPs needed to be developed and that mentoring by allergy specialists needed to be in place wherever possible.

He said, "There are only seven specialist allergy centres in the country, and the problem is getting enough training for GPs by allergy specialists. I know the government does not want to hear about us spending more money, but I'm not sure how else you can possibly do this."

"We need to improve allergy services available in secondary care, and in the interim we've got to train doctors in primary care up to a particular level, and then we can move forward as a team," he added. □

The Review of Services for Allergy is available at www.dh.gov.uk/ publications.