Whooping cough is quite common and can be diagnosed clinically

EDITOR—Harnden et al, with the help of a recently available blood test, have gone some way to confirm that whooping cough is still about.¹ Since 1977, I have, with the help of my practice colleagues, been recording every clinically diagnosable case of whooping cough (based on a minimum of three weeks of paroxysmal coughing). This year so far we have seen six cases in a practice of 11 000. In 2002 we recorded 44 cases. Ten had blood specimens tested; nine were positive and the 10th was lost. Twenty three had pernasal swabs taken; 14 were positive. This small practice was responsible for 5% of all whooping cough notifications in England and Wales in 2002. This surely cannot be because there is more whooping cough where I work.

I have had four papers published in the *BMJ* as a result of this study.²⁻⁵ I have decided to publish the incidence data on the web. Some years ago I set up a website to help patients diagnose their own whooping cough, and I can confirm from the feedback that doctors the world over seem to be equally poor at diagnosing it. I have published the Keyworth figures (www.whoopingcough.net/keyworth). Basic data on all cases are available, as well as year-on-year figures compared with national notifications (figure).

Whooping cough is a distressing illness, especially when it is undiagnosed. I am hopeful that the paper by Harnden et al will wake us up about this disease, which, if my

data are correct, is just as common as it was 25 years ago after the vaccine scare settled.

Doug Jenkinson *general practitioner principal*
Keyworth Health Centre, Keyworth, Nottingham
NG12 5JU
dougjenkinson@btinternet.com

Competing interests: DJ has a website on whooping cough that charges a fee for personal advice. The fees have so far never covered the expenses for the site.

- 1 Harnden A, Grant C, Harrison T, Perera R, Brueggemann AB, Mayon-White R, et al. Whooping cough in school age children with persistent cough: prospective cohort study in primary care. *BMJ* 2006;333:174-7. (22 July.)
- 2 Jenkinson D. Outbreak of whooping cough in general practice. *BMJ* 1978;277:896.
- 3 Jenkinson D. Whooping cough: what proportion of cases is notified in an epidemic? *BMJ* 1983;287:185-6.
- 4 Jenkinson D. Duration of effectiveness of pertussis vaccine: evidence from a ten year community study. *BMJ* 1988;296:612-4.
- 5 Jenkinson D. Natural course of 500 consecutive cases of whooping cough: a general practice population study. *BMJ* 1995;310:299-302.

Author's reply to review of his book on autism

EDITOR—The misrepresentations and omissions in Fitzgerald's review of my book demand correction.¹ The book overviews evidence for (and against) the apparent rise in autism and concludes that the increase is probably real—pointing to an environmental risk factor so far unknown.

A focus of the book is on physiological dysregulation that accompanies and exacerbates autism. It reviews evidence that limbic damage can both cause and reflect physiological problems, with environmental toxins including heavy metals having a causal role, although concluding that the rise in autism cannot be uniquely attributed to such exposure (but it remains a strong suspect). Just one short chapter refers to possible strategies for treatment and prevention.

Fitzgerald focuses his critical attentions only on this chapter: most of the book is not addressed. He asserts that there is no coherent scientific rationale for the treatments and little evidence of their efficacy. He omits to note the considered position taken by the book, that the field is fraught with uncertainty because few logical therapeutic approaches have been evaluated systematically. Reviewer and author are therefore substantially in agreement.

Fitzgerald then draws attention to an autistic boy who died while receiving chelation treatment. This is also misleading: use of the wrong drug was responsible for the death.²

He then says that guidelines from the American Academy of Neurology explicitly reject a series of biomedical tests.³ The recommendations speak not of rejection but of inadequate supporting evidence, a different matter. This is an area of debate.

Regarding biomedical intervention, affected children should be spared unproved treatments until efficacy has been shown in controlled trials. Trials (examples include haloperidol, risperidone, methylphenidate hydrochloride (Ritalin)) have been conducted through mainstream medicine and not by (in the words of Fitzgerald) quacks and charlatans. Valproate continues to be prescribed for epilepsy associated with autism, even though valproic acid is a known cause of autism.⁴

Fitzgerald's exclusive focus on therapeutic options is unfortunate, for this is a minor aspect. The central thesis offered by the book is of a rational and plausible sequence of events that causes as well as exacerbates autism—addressed by Fitzgerald as "speculative."

Rushing into biomedical remediation without fullest consideration of justification and efficacy is a mistake, but it is an equally grave error to dismiss without objective consideration of all the evidence (as laid out in the book), and new research under way,⁵ an environmental and physiological contribution to autism.

Richard Lathe *neuroscience consultant*
Pieta Research, PO Box 27069, Edinburgh
EH10 5YW
rlathe@pieta-research.org

Competing interests: None declared.

Incidence of whooping cough in Keyworth practice and in England and Wales both equated to 100 000 population

- 1 Fitzpatrick M. Autism, brain and environment. *BMJ* 2006;333:205. (29 July.)
- 2 Centers for Disease Control and Prevention. Deaths associated with hypocalcemia from chelation therapy—Texas, Pennsylvania, and Oregon, 2003-2005. *Morb Mortal Wkly Rep MMWR* 2006;55:204-7.
- 3 Filipek PA, Accardo PJ, Ashwal S, Baranek GT, Cook EH Jr, Dawson G, et al. Practice parameter: screening and diagnosis of autism: report of the Quality Standards Subcommittee of the American Academy of Neurology and the Child Neurology Society. *Neurology* 2000;55:468-79.
- 4 Moore SJ, Turnpenny P, Quinn A, Glover S, Lloyd DJ, Montgomery T, et al. A clinical study of 57 children with fetal anticonvulsant syndromes. *J Med Genet* 2000;37:489-97.
- 5 Hertz-Picciotto I, Croen LA, Hansen R, Jones CR, Van de WJ, Pessah IN. The CHARGE study: an epidemiologic investigation of genetic and environmental factors contributing to autism. *Environ Health Perspect* 2006;114:1119-25.

What works in schizophrenia

BSIP, PHO/SPL

Depot preparations may improve outcomes

EDITOR—The study by Tiihonen et al indicates that antipsychotics are not equally effective in treating schizophrenia.¹ After a first admission with schizophrenia, patients treated with clozapine, olanzapine, or perphenazine depot had substantially lower risks of readmission, and discontinuation of treatment for any reason, than patients treated with oral haloperidol.

The increased effectiveness of clozapine is understandable in terms of its superior efficacy in treatment resistant schizophrenia.² The increased effectiveness of perphenazine depot presumably derives from its ability to reduce non-compliance, a common problem in schizophrenia that can be overt or covert.³ In the study reported by Tiihonen et al, the less favourable outcome for oral perphenazine, in contrast to the depot preparation, supports this hypothesis.

The effectiveness of clozapine and perphenazine depot may also partly reflect the regular clinical contact and supervision that are integral to both treatments; with a depot this relates to its administration and with clozapine to regular haematological monitoring. The results are unlikely to reflect a prescribing bias as depots are largely used in patients who comply poorly with oral drugs and clozapine is restricted to use in treatment resistant schizophrenia.

Despite its effectiveness, perphenazine depot was not commonly used. This may reflect negative perceptions about depots.⁴ In a Cochrane review a greater proportion of patients treated with depot than with oral drugs showed global improvement, perhaps reflecting depots reducing partial compliance.⁵ The work of Tiihonen et al adds to the view that depot antipsychotic drugs are effective for some patients. Like all antipsychotic drugs, they should be prescribed on an individual basis after full discussion with the patient and considering a range of factors.

Pharmacologically, clinical outcomes in schizophrenia can be improved by using drugs that are either more efficacious or improve compliance, and one way to do this is to use a depot preparation.

Peter M Haddad *consultant psychiatrist*
Bolton, Salford and Trafford Mental Health NHS Trust, Salford M30 0GT
Peter.Haddad@bstmt.nhs.uk

Omar Niaz *specialist registrar in psychiatry*
Sheffield Care Trust, Fulwood House, Sheffield S10 3TG

Competing interests: PMH has received honorariums for lecturing or attending advisory boards from Eli Lilly, Janssen-Cilag, and Novartis.

- 1 Tiihonen J, Wahlbeck K, Lonnqvist J, Klaukka T, Ioannidis JP, Volavka J, et al. Effectiveness of antipsychotic treatments in a nationwide cohort of patients in community care after first hospitalisation due to schizophrenia and schizoaffective disorder: observational follow-up study. *BMJ* 2006;333:224-7. (29 July.)
- 2 Chakos M, Lieberman J, Hoffman E, Bradford D, Sheitman B. Effectiveness of second-generation antipsychotics in patients with treatment-resistant schizophrenia: a review and meta-analysis of randomized trials. *Am J Psychiatry* 2001;158:518-26.
- 3 Lindstrom E, Borgefors K. Patient compliance with drug therapy in schizophrenia. Economic and clinical issues. *Pharmacoeconomics* 2000;18:106-24.
- 4 Patel MX, Nikolaou V, David AS. Psychiatrists' attitudes to maintenance medication for patients with schizophrenia. *Psychol Med* 2003;33:83-9.
- 5 Adams CE, Fenton MK, Quraishi S, David AS. Systematic meta-review of depot antipsychotic drugs for people with schizophrenia. *Br J Psychiatry* 2001;179:290-9.

Cognitive behaviour therapy is not effective

EDITOR—Kingdon's statement that more than 20 randomised controlled trials and five meta-analyses have shown cognitive behaviour therapy to be beneficial in schizophrenia gives an oversimplified picture of both the randomised controlled trials and the meta-analyses.¹ Reviewing the randomised controlled trials, Tarrier and Wykes, two supporters of cognitive behaviour therapy in schizophrenia, noted that five included groups who received befriending, supportive counselling, or problem solving to control for the non-specific effects of intervention, in other words as a psychological placebo.² They said that not one study has shown clear and significant overall differences between cognitive behaviour therapy and the non-specific control groups.³

The conclusion of the Cochrane Collaboration's meta-analysis of cognitive behaviour therapy for schizophrenia was currently that trial based data supporting the wide use of such treatment for people with schizophrenia or other psychotic illnesses are far from conclusive.³ Compared with standard care, cognitive behaviour therapy was found not to reduce relapse and readmission; it helped mental state over the medium term but after one year the difference had gone, and it did not show a consistent effect on continuous measures of mental state. Compared with supportive psychotherapy, cognitive behaviour therapy had no effect on relapse or on the outcome "no clinically meaningful improvements in mental state" over the same time periods.

Behind all the recent publicity surrounding cognitive behaviour therapy for

schizophrenia, the truth is that it works only in poorly controlled trials and not in well controlled ones.

Peter J McKenna *professor of psychiatry*
University of Glasgow, Glasgow G12 0XH
peter.mckenna@virgin.net

Competing interests: None declared.

- 1 Kingdon C. Psychological and social interventions for schizophrenia. *BMJ* 2006;333:212-3. (29 July.)
- 2 Tarrier N, Wykes T. Is there evidence that cognitive behaviour therapy is an effective treatment for schizophrenia? A cautious or cautionary tale? *Behav Res Ther* 2004;42:1377-401.
- 3 Jones C, Cormac I, Silveira da Mota Neto JI, Campbell C. Cognitive behaviour therapy for schizophrenia. *Cochrane Database of Syst Rev* 2004, Issue 4.

Fluoroquinolone resistance in *Salmonella* Typhi

EDITOR—As Parry et al point out,^{1,2} appropriate laboratory methods are crucial in detecting clinically important quinolone resistance.

We highlight the emergence of strains of *Salmonella enterica* serovar Typhi (*S* Typhi) that show reduced susceptibility to the fluoroquinolones but are susceptible to nalidixic acid (minimum inhibitory concentration <16 mg/l). In a review of all 692 isolates of *S* Typhi sent to the Laboratory for Enteric Pathogens at the Health Protection Agency in London between 2000 and 2003 we detected 49 isolates that were susceptible to nalidixic acid but had reduced susceptibility to fluoroquinolone (minimum inhibitory concentration 0.125-1.0 mg/l; table).³ When the country of acquisition was known, 18 of these isolates were from patients who had visited India; eight, Pakistan; four, Bangladesh; and one, Kenya.

Overall, of 271 isolates with reduced susceptibility to fluoroquinolone, 18% were susceptible to nalidixic acid and therefore would not have been detected by routine screening with a nalidixic acid disc. No clinical outcome data are available and so the clinical relevance of these strains is uncertain, but an enhanced surveillance study is being undertaken by the Health Protection Agency.⁴

Although resistance to nalidixic acid (minimum inhibitory concentration >256 mg/l) remains an important marker for failure of fluoroquinolone treatment in typhoid fever, several isolates show reduced susceptibility to fluoroquinolone while remaining susceptible to nalidixic acid. This is particularly true for isolates from the Indian subcontinent. Furthermore, in six of the 49 such isolates (table), none of the mutations

S Typhi susceptible to nalidixic acid but with reduced fluoroquinolone susceptibility imported into England, Scotland, and Wales, 2000-3

Year	Susceptible to nalidixic acid and reduced susceptibility to fluoroquinolones	Resistant to nalidixic acid and reduced susceptibility to fluoroquinolones	Total with reduced susceptibility to fluoroquinolone (% susceptible to nalidixic acid)
2000	10	47	57 (18)
2001	8	55	63 (13)
2002	10	46	56 (18)
2003	21	74	95 (22)
Total	49	222	271 (18)

in *gyrA* and *parC* commonly associated with reduced resistance to fluoroquinolone were present using the GAMA method.⁵ Therefore a new mechanism of fluoroquinolone resistance, which might not confer nalidixic acid resistance, may be emerging in *S Typhi*. Our findings underscore the importance of estimating the minimum inhibitory concentration of the agent selected for treatment.

Fiona J Cooke MRC clinical research training fellow
fiona@sanger.ac.uk

John Wain member of faculty
Wellcome Trust Sanger Institute, Wellcome Trust
Genome Campus, Cambridge CB10 1SA

E John Threlfall professor
Health Protection Agency, Centre for Infections,
London NW9 5HT

Competing interests: None declared.

- 1 Parry CM, Karunanayake L, Coulter JB, Beeching NJ. Test for quinolone resistance in typhoid fever. *BMJ* 2006; 333:260-1. (29 July.)
- 2 Bhutta ZA. Current concepts in the diagnosis and treatment of typhoid fever. *BMJ* 2006;333:78-82. (8 July.)
- 3 Cooke FJ, Day M, Wain J, Ward LR, Threlfall EJ. Cases of typhoid fever imported into England, Scotland and Wales (2000-2003). *Trans R Soc Trop Med Hyg* (in press).
- 4 Anonymous. A pilot of enhanced surveillance of enteric fever in England, Wales and Northern Ireland. *Commun Dis Rep CDR Wkly* 2006;16(18):3-4.
- 5 Walker RA, Skinner JA, Ward LR, Threlfall EJ. LightCycler *gyrA* mutation assay (GAMA) identifies heterogeneity in *GyrA* in *Salmonella enterica* serotypes Typhi and Paratyphi A with decreased susceptibility to ciprofloxacin. *Int J Antimicrob Agents* 2003;22(6):622-5.

An independent NHS?

Why it really is time to separate from direct government involvement

EDITOR—I received 21 items of correspondence about my personal view on separating the NHS from direct government involvement, mostly supporting the idea and many urging me to take it further.¹ I have worked in the NHS for 28 years and my conclusions are based on first hand experience of the waste associated with massive bureaucracy to support edicts from the top, short term solutions to satisfy the ruling party's election prospect, and, most damaging of all, the three yearly cycle of NHS reforms.

There is lack of confidence in any health minister, their handful of civil servants, and their chosen "advisers" in ever producing policies that are foremost for the benefit of the country without worrying about their position and their party's election prospect.

The model I am suggesting would remove this monolithic self interest. The

model is apolitical, has input from all parties, and is not an unaccountable quango or technocracy of professionals. It is new, unique, democratic, and accountable to parliament with the necessary regulations to ensure its effectiveness. Such reform would be the least costly in the history of the NHS. The only change would be the people who sit around the decision making table and their advisers. All it needs is men and women with courage and conviction to put their country before their party.

If such a body came to the conclusion that we should overtly ration health care, the country would listen and better trust its motives. If it concluded that some input from the private sector was inevitable, the country would listen and might reluctantly accept but trust the source of that decision.

To Gordon Brown, I say, one more bold decision, and the country will remember who saved the NHS. To David Cameron, if Brown misses this opportunity then it's your chance to double your prospect of winning. To my colleagues, please speak up as advocates for your patients and your communities.

Layla Jader consultant in public health medicine
National Public Health Service for Wales, Cardiff
CF10 3NW
Layla.Jader@nphs.wales.nhs.uk

Competing interests: None declared.

- 1 Jader L. It is time to separate the NHS from direct government involvement. *BMJ* 2006;332:1518. (24 June.)

There is no "UK health service"

EDITOR—Berwick and Leatherman express their steadfast support for the NHS eloquently.¹ We forgive them for their error and that of the other authors in this series of articles in writing about the NHS as a UK model when commenting only on the NHS in England as directed from Westminster. The NHS in Scotland, in particular, is different in many respects.

A need exists to survey and audit the differences with the other home nations, of course, and I communicated that suggestion to the authors of a paper last year about the NHS in England.² Feachem et al too seemed unaware from a distance that England is not the UK.

In Scotland, I was glad that two reforms ago when we got local healthcare cooperatives, they had no commissioning responsibility. There was enough to do without that. Just when these had matured and might have taken on that challenge, reform has produced community health partnerships, which in my area have yet to have a whimper of a profile some 18 months after their birth.

Philip A Gaskell general practitioner
Allan Park Medical Practice, Stirling FK8 2QD
philip.gaskell@nhs.net

Competing interests: None declared.

- 1 Berwick DM, Leatherman S. An independent NHS? Steadying the NHS. *BMJ* 2006;333:254-5. (29 July.)
- 2 Feachem RGA, Sekhri NK. Moving towards true integration. *BMJ* 2005;330:787-8.

Towards evidence based changes in NHS in all UK countries

EDITOR—As a Scottish doctor I can tell Godlee that the assumption that "NHS" means "NHS in England" is not only the *BMJ* but in government statements, the BBC, and the national British newspapers eventually produces a weary acceptance.¹ Frankly, like the *BMJ*, I have no idea what's going on in the NHS in Scotland because it is so unreported.

The degree of divergence in NHS systems in the four countries based, as the BMA briefing paper Godlee refers to points out, on profoundly different philosophies and policy groups, provides us with a unique opportunity to learn something. What are the effects of the different NHS reforms on the experience of patients? And what are the effects of the different reforms on the experience of those who are delivering health care—the NHS employees in the four different countries? If we could at least read some comparative journalism, not to mention proper research which considered these questions, then we would be able to take the first steps in all four countries to create NHS changes which were evidence based.

Robert W Leckridge associate specialist
Glasgow Homeopathic Hospital, Glasgow G12 0XQ
bobleckridge@gmail.com

Competing interests: None declared.

- 1 Godlee F. Editor's choice. B is for British. *BMJ* 2006;333:0. (13 July.)

Suffering from files

EDITOR—Thornton says that we shouldn't remove the NHS from politics—he says that the NHS wouldn't exist today without politicians like Nye Bevan.¹ He thinks that more democratic commissioning—perhaps even from local government—could increase NHS accountability and at the same time give more freedom to clinicians. Can we really square the circle in this way?

Or would this idea become just another initiative leading nowhere but more bureaucracy and an even higher paper mountain? What would Bevan say to the range of initiatives that the NHS has gone through in the past 10 years? Here's what he said about a contemporary administrator: "Poor fellow, he suffers from files."²

Sarah J Farrow specialist registrar in rheumatology and medicine
University Hospital Lewisham, London SE13 6LH
sfarrow@doctors.org.uk

Competing interests: None declared.

- 1 Thornton S. Democratic control is essential. *BMJ* 2006;333:251-2. (29 July.)
- 2 Bevan A. Bureaucracy. www.quotationsbook.com/quotes/5028/view (accessed 3 Aug 2006).

bmj.com

We select the letters for these pages from the rapid responses posted on bmj.com favouring those received within five days of publication of the article to which they refer.

Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.