and is marginally more potent.¹⁶ Concern has long been expressed that when given in modest excess aspirin may prove toxic and even fatal in children.^{17 18} This concern has increased recently because of evidence that aspirin may be implicated in the pathogenesis of Reye's syndrome.¹⁹⁻²¹ The latter is uncommon but may occur in small epidemics and has a high mortality. In 1977 and 1978 there were 655 cases reported in the United States with a mortality of 32%.²⁰ The clinical features of the disease are vomiting and rapidly progressive encephalopathy after a viral like illness, and the characteristic pathological findings are fatty infiltration of the liver and cerebral oedema.

Various viruses have been implicated, especially influenza and chickenpox, but many of the features of the disease are similar to those of salicylate poisoning,¹⁹ and case-control studies show a significant relation between administration of aspirin during the prodromal illness and the development of Reye's syndrome. In three studies a total of 141 patients with Reye's syndrome were compared with 247 controls matched for age and the nature of the prodromal illness.²⁰⁻²² Some 97% of the patients and 67% of the controls had been given salicylate, whereas 15% of patients and 37% of controls had been given paracetamol. The doses of salicylate used were not excessive but the patients tended to have had more than the controls. Despite the rarity of Reye's syndrome it seems sensible to conclude that aspirin should be avoided in children with fever, especially during epidemics of influenza or chickenpox.²³ The danger of paracetamol in overdose is well known but there is no evidence that it is harmful in therapeutic doses (admittedly, the same was held to be true of aspirin for many years). A recent survey in the United States showed that aspirin and paracetamol had been given to equal numbers of children before admission to hospital, but the doctors in two teaching hospitals prescribed paracetamol 10 times more often than aspirin.²⁵ Concern has been expressed that increased use of paracetamol might lead to an increase in accidental poisoning in children, but the drug is readily available and is commonly used by adults. Any increase in poisoning would seem more likely to be due to the general household use of the drug than to its prescription, or use by parents, for children.

At present, therefore, when antipyretic drug treatment is considered necessary for a child paracetamol must be regarded as the drug to be preferred. Single dose studies of paracetamol have shown that the antipyretic effect starts between one and two hours after the dose, the maximum effect (reduction of fever by about 70%) occurring at between one and a half and three and a half hours.^{16 26} Recommendations for dose vary from 10 to 20 mg/kg body weight and for the interval between doses from four to eight hours.^{26 27} Paracetamol elixir contains 120 mg in 5 ml. Between 1 and 5 years of age, therefore, an average appropriate dose would vary from one to two and a half standard 5 ml spoonfuls every four to six hours. Paracetamol suppositories are effective²⁷ but not readily available; they are rarely necessary.

D P ADDY

Consultant Paediatrician,
Dudley Road Hospital,
Birmingham B18 7QH

¹ Roberts KB, Borzy MS. Fever in the first eight weeks of life. *Johns Hopkins Med J* 1977;**141**:9-13.

² McGowan JE Jr, Bratton L, Klein JO, Finland M. Bacteremia in febrile children seen in a "walk-in" pediatric clinic. *N Engl J Med* 1973;**288**:1309-12.

³ McCarthy P, Dolan T. Comments. In: Oski FA, Stockman JA, eds. *Year-book of pediatrics 1979*. Chicago: Yearbook Medical Publishers, 1979:77-8.

- ⁴ Bratton L, Teele DW, Klein JO. Outcome of unsuspected pneumococemia in children not initially admitted to hospital. *J Pediatr* 1977;**90**:703-6.
- ⁵ Kluger MJ. Fever. *Pediatrics* 1980;**66**:720-4.
- ⁶ Heim T. Fever. In: Davis JA, Dobbing J, eds. *Scientific foundations of paediatrics*. 2nd ed. London: William Heinemann, 1981:118-9.
- ⁷ Camfield PR, Camfield CS, Shapiro SH, Cummings C. The first febrile seizure—antipyretic instruction plus either phenobarbital or placebo to prevent recurrence. *J Pediatr* 1980;**97**:16-21.
- ⁸ Steele RW, Tanaka PT, Lara RP, Bass JW. Evaluation of sponging and of oral antipyretic therapy to reduce fever. *J Pediatr* 1970;**77**:824-9.
- ⁹ Hunter J. Study of antipyretic therapy in current use. *Arch Dis Child* 1973;**48**:313-5.
- ¹⁰ Colgan MT, Mintz AA. The comparative antipyretic effect of N-acetyl-p-aminophenol and acetylsalicylic acid. *J Pediatr* 1957;**50**:552-5.
- ¹¹ Eden AN, Kaufman A. Clinical comparison of three antipyretic agents. *Am J Dis Child* 1967;**114**:284-7.
- ¹² Steele RW, Young FSH, Bass JW, Shirkey HC. Oral antipyretic therapy. Evaluation of aspirin-acetaminophen combination. *Am J Dis Child* 1972;**123**:204-6.
- ¹³ Tarlin L, Landrigan P. A comparison of the antipyretic effect of acetaminophen and aspirin. Another approach to poison prevention. *Am J Dis Child* 1972;**124**:880-2.
- ¹⁴ Lovejoy FH Jr. Aspirin and acetaminophen: a comparative view of their antipyretic and analgesic activity. *Pediatrics* 1978;**62**, suppl:904-9.
- ¹⁵ Yaffe SJ. Comparative efficacy of aspirin and acetaminophen in the reduction of fever in children. *Arch Intern Med* 1981;**141**:286-92.
- ¹⁶ Wilson JT. Antipyretic management of febrile seizures. In: Nelson KB, Ellenberg JH, eds. *Febrile seizures*. New York: Raven Press, 1981:231-9.
- ¹⁷ Dodd K, Minot AS, Arena JM. Salicylate poisoning: an explanation of the more serious manifestations. *Am J Dis Child* 1937;**53**:1435-46.
- ¹⁸ Craig JO, Ferguson IC, Syme J. Infants, toddlers, and aspirin. *Br Med J* 1966;**ii**:757-61.
- ¹⁹ Giles H McC. Encephalopathy and fatty degeneration of the viscera. *Lancet* 1965;**ii**:1075.
- ²⁰ Starko KM, Ray CG, Dominguez LB, Stromberg WL, Woodall DF. Reye's syndrome and salicylate use. *Pediatrics* 1980;**66**:859-64.
- ²¹ Halpin TJ, Holtzhauser FJ, Campbell RJ, et al. Reye's syndrome and medication use. *JAMA* 1982;**248**:687-91.
- ²² Waldman RJ, Hall WN, McGee H, Amburg GV. Aspirin as a risk factor in Reye's syndrome. *JAMA* 1982;**247**:3089-94.
- ²³ American Academy of Pediatrics Committee on Infectious Diseases. Aspirin and Reye syndrome. *Pediatrics* 1982;**69**:810-2.
- ²⁴ Glezen WP. Aspirin and Reye's syndrome. *Am J Dis Child* 1982;**136**:971-2.
- ²⁵ Mitchell AA, Lovejoy FH Jr, Slone D, Shapiro S. Acetaminophen and aspirin. Prescription, use, and accidental ingestion among children. *Am J Dis Child* 1982;**136**:976-9.
- ²⁶ Wilson JT, Brown DR, Bocchini JA Jr, Kearns GL. Efficacy, disposition and pharmacodynamics of aspirin, acetaminophen and choline salicylate in young febrile children. *Ther Drug Monit* 1982;**4**:147-80.
- ²⁷ Vernon S, Bacon C, Weightman D. Rectal paracetamol in small children with fever. *Arch Dis Child* 1979;**54**:469-70.

Food allergy

Early this year Fisons organised a workshop on food allergy in Harrogate, under the chairmanship of Professor R R A Coombs—a follow up to an earlier workshop on the same theme in 1980. The meeting produced paradoxical results: the clinical evidence for food intolerance was strengthened, while doubts were thrown on the diagnostic value of specific tests for that intolerance.

Clinically, the most important presentations concerned the contribution of food intolerance to gastrointestinal disease. Patients with Crohn's disease are known to respond to an elemental diet but little is known about the manner in which diet influences their risk of relapse once the elemental diet is stopped. Dr J O Hunter (Addenbrooke's Hospital, Cambridge) reported that dietary exclusion produced long term remission in 20 out of 28 patients with histologically proved Crohn's disease, in many of whom the disease had remained active despite treatment with steroids or surgery. In addition 141 out of 205 patients with the irritable bowel syndrome also responded to dietary exclusion. The foods most commonly responsible were wheat, dairy products, corn, citrus fruit, tea, and coffee. These clinical observations had been supported by appropriately controlled challenge tests.

Milk has become an accepted cause of food intolerance. A study by Dr P D Manuel (All Saints Hospital, Chatham) of acute gastroenteritis in Indonesian infants showed that at least one fifth of those affected developed enteropathy attributable to sensitivity to cows' milk, a conclusion based on clinical and histological findings reinforced with milk challenges. The importance of intolerance to both human and cows' milk as a cause of chronic gastrointestinal symptoms in infants was emphasised by many observers.

Other associations between gastrointestinal disease and food intolerance were also advanced, though in contexts which enjoined a more cautious evaluation. To purists gluten sensitivity remains synonymous with jejunal villous atrophy (coeliac disease). Dr P Asquith (East Birmingham Hospital), however, claimed that seven out of 12 patients with coeliac disease that had resisted rigorous gluten exclusion responded to exclusion of soybean proteins. A link between chronic gastritis and food intolerance was claimed by Dr E P Jewell and his colleagues (Radcliffe Infirmary, Oxford) in studies of 44 patients with immunopathological evidence of type I immune responses in gastric biopsy specimens and good clinical and histological responses to sodium cromoglycate.

Outside the gastrointestinal tract, atopic eczema has received particular attention in terms of possible food intolerance. Dr D J Atherton (Hospital for Sick Children, Great Ormond Street, London) had carried out several studies of dietary exclusion in this disorder. He emphasised that dietary measures needed to be considered in only one child in four with this disorder; the remainder could be treated with orthodox measures. Of those children needing dietary control most responded to exclusion of milk, eggs, chicken, colouring matters, and preservatives. The few remaining needed a much stricter diet with sequential reintroduction of foods to identify offending items, tartrazine being the ingredient most commonly implicated.

Enthusiasts for invoking food intolerance as an important cause of migraine will be encouraged by the observations reported by Dr J Egger (Hospital for Sick Children, Great Ormond Street, London). Not only did most of 75 children with severe migraine respond to an exclusion diet but double blind controlled challenges specifically implicated milk, tea, rye, and artificial colourings as provoking items. Data on bronchial asthma were more fragmentary, with a dearth of properly controlled observations.

The mechanisms that provoke food intolerance were subjected to timely rescrutiny. The first workshop had taken the view that the excessive absorption of food allergens was the critical step in initiating food allergic disorders, ascribing the resulting reactions variously to classic cell mediated and humoral immune responses. While IgE has received the most attention in terms of humoral responses, the possible importance of IgG₄ was emphasised by Dr R St C Barnetson (Royal Infirmary, Edinburgh). Many observers doubt the validity of the techniques on which claims of increased intestinal permeability have been based, but this important issue received little attention at the meeting. It was left to Dr J Walker-Smith (Queen Elizabeth Hospital for Children, London) to present the only relevant data—ultrastructural studies showing increased uptake of antigens by damaged enterocytes in jejunal biopsy specimens from children with enteropathies.

The focus of attention had, indeed, moved from conventional assays of cell mediated and humoral immunity to pharmacological mediators of inflammation. This possibility, first advocated by Professor Lessof and his colleagues (Guy's Hospital, London), has been followed up by other workers.

Professor A W Cuthbert (Department of Pharmacology, Cambridge) described experiments in which cows' milk protein induced inappropriate chloride secretion when applied in vitro to the colonic epithelium of guinea pigs sensitised to milk proteins. This secretion could be blocked by indomethacin, suggesting that prostaglandins may be responsible for mediating the secretory diarrhoea induced by food intolerance. Indeed, only the observation that serum from sensitised animals confers this abnormal pattern of electrolyte secretion on normal gut epithelium obtained from normal animals provided any link with classical type I immunity. Dr P McLaughlan and her colleagues (Addenbrooke's Hospital, Cambridge) described experiments in which jejunal biopsy specimens cultured from patients with gluten hypersensitivity released histamine after challenge with gliadin or anti-IgE. They also provided hints that basophils isolated from the blood of patients with suspected food intolerance may release histamine after exposure in vitro to food allergens such as cows' milk, egg, soya, and gliadin, as well as anti-IgE. Neutrophil chemotactic factors may also be important in the pathogenesis of bronchial asthma (Professor A B Kay, Cardiothoracic Unit, London). Experiments of this kind do not yet provide definitive information about the mechanisms which cause food intolerance, but they do emphasise the wisdom of using the term "food intolerance" to describe reactions in circumstances in which the part played by classical immune reactions is uncertain.

Given this reappraisal, it is not surprising that traditional means of diagnosing disorders due to food allergy were widely challenged. Both clinical studies (Dr M Kemeny, Guy's Hospital, London) and epidemiological studies (Dr M L Burr, MRC Epidemiology Unit, Cardiff) showed that serum IgE specific for food allergens can be detected by techniques such as the radioallergosorbent assay in patients with disorders such as atopic eczema and bronchial asthma. Unfortunately, the diagnostic limitations of these findings are now apparent. Firstly, both IgE and IgG antibodies reactive with food allergens have been found in a high proportion of blood donors without any obvious clinical consequences (Dr R M R Barnes, Royal Liverpool Hospital, and Dr Anne Ferguson, University of Edinburgh). Secondly, several participants reported a poor correlation between positive immunological tests for food allergens and the clinical response to withdrawing allergens. Thirdly, conventional immunological tests did not help to predict which foods would provoke a relapse of Crohn's disease (Dr J Hunter, Addenbrooke's Hospital, Cambridge). Finally, infants who have shown undoubted cows' milk sensitivity may eventually become clinically tolerant to milk despite maintaining serum IgE antibodies specifically reactive with milk proteins. Plainly, claims by commercial laboratories that food allergic disorders can be diagnosed on the basis of serum tests are unwarranted and specific "cytotoxic" tests for diagnosing food allergy by exposing white cells to specific foods should be regarded with proper suspicion.

What conclusions may be drawn from these proceedings? The diagnosis of food intolerance still depends primarily on clinical observations and the effective use of properly controlled double blind challenges. Even this time honoured method has its drawbacks, however; dietary manipulation in itself has an important placebo effect, and the reaction to food challenge is not immutable but susceptible to changing conditions such as intercurrent infections. The place of properly controlled clinical observations has not been undermined by conventional laboratory tests, which are, at best, an adjunct to clinical evaluation. Our increasing awareness of the part inflam-

matory mediators such as prostaglandins may play in food intolerance underscores the folly of attributing all food intolerance to classical immunological mechanisms. Treatment by exclusion diets is a demanding occupation for skilled dietitians, and many inconsistencies in published reports may well be attributable to the varying zeal with which exclusion diets are pursued. Dr P Hall-Smith (Brighton Health Authority) made the constructive suggestion that proper instruction should be available for dietitians wishing to undertake this task. As yet there is little evidence that sodium cromoglycate has reduced the importance of dietary measures, and only one study by Professor Louisa Businco (University of Rome) provided any controlled data in a study of atopic eczema that this drug is helpful in common disorders allegedly attributable to food intolerance.

A M DENMAN

Consultant Physician,
Connective Tissue Disease Research Group,
Clinical Research Centre,
Harrow HA1 3UJ

Clinical biochemistry nearer the patient

Despite its expansion in the past two decades clinical biochemistry has undergone little conceptual change since it first became an indispensable part of hospital practice some 60 years ago. Now, however, it is on the brink¹⁻⁵ of a revolution far more reaching in effect than the purely quantitative one arising from the introduction of automation in the early 1960s. The new factor—comparable in importance to the introduction of the stethoscope by Laennec in the nineteenth century—is our ability to perform tests much nearer to the patient than was hitherto possible. Analytical procedures previously able to be performed only in custom built laboratories staffed by highly trained technicians and scientists can now be carried out by relatively unskilled staff in the clinic or in the ward without loss of precision or accuracy.⁶

The technical background to this change is a series of advances in the construction of ion specific and enzyme electrodes, in solid phase chemistry, electronics, non-invasive "chemical" probes, microprocessors, and, above all, in the miniaturisation of apparatus. This change from the old to the new way of doing clinical chemistry will not be achieved painlessly or without a struggle. Laboratory workers will see themselves threatened by an apparent down grading of their hard learnt skills, and clinicians will need to rethink the way they request pathological tests and indeed their whole clinical behaviour.

Once insulin became available in 1922 patients suffering from diabetes mellitus could no longer reasonably be treated without ready access to blood glucose measurements—not primarily for diagnostic purposes but as a guide to dosage and monitoring treatment. This early but key monitoring function of analytical chemistry as a guide to the benefit, or otherwise, of treatment accounts for at least one third of all laboratory requests in clinical chemistry⁷ and is growing in importance with the trend towards tailoring drug treatment to the individual.⁸ It has, however, been largely overshadowed by the more dramatic use of clinical chemistry as an aid to diagnosis and its largely abortive application to screening for

early or unsuspected diseases. Indeed, as much as anything, belief in screening tests was responsible for the introduction into laboratories of expensive multichannel instruments capable of producing large numbers of mainly irrelevant, often uninterpretable, and occasionally misleading results⁹⁻¹²—though their saving grace was a commendably high standard of analytical precision and accuracy.

Since the numbers of tests performed, their precision, and their accuracy are all easy to quantify, these aspects of laboratory performance (rather than the clinical value and relevance of the service provided) have tended to become the yardstick against which quality and cost effectiveness have been judged. Similarly the status of laboratory heads and technicians has grown in direct proportion to the size and complexity of their instruments, which could reasonably be housed only in large central laboratories often remote from the patients they served. This, in turn, has demanded the introduction of elaborate systems for the collection, processing, documentation, and dissemination of information—which for many tests cost more than the actual analysis¹³ and, despite computerisation, still depend heavily on human factors, not the least of which is the sense of participation. Indeed, human factors account for most of the 1-2% of all results dispatched from laboratories that are hopelessly wrong, and since these are mostly due to errors in clerking, labelling, or transcription outside the analytical area they may often be completely beyond the control of the laboratory director.

Over the years disillusionment has grown with the cost effectiveness and clinical usefulness of many clinical chemical procedures as presently performed.¹⁴⁻¹⁷ The only remedy seemed to be better education.¹⁸⁻²¹ All this will be changed, however, with the gradual introduction of technically advanced but cheap and portable analytical systems suitable for small laboratories, hospital outpatient clinics, intensive treatment and special care units, ward side rooms, general practitioners' surgeries, and even patients themselves. These new systems are within the technical competence of virtually any literate and sufficiently well motivated person,^{5 22 23} and their advent demands an urgent reappraisal of the part played by clinical laboratories in modern medical practice.

Apart from purely technical considerations, such as the specificity, accuracy, and reliability of the apparatus to be used, several other questions need to be answered before clinical chemistry becomes established nearer the patient: these include the question of who should initiate, carry out, interpret, and assume legal responsibility for tests performed outside the laboratory. Special problems are posed by screening tests in those rare cases where they can be scientifically and ethically justified²⁴ and by some essential procedures which are required only rarely and for which reliable "black box" technology is not and is unlikely to become available. Possibly as common tests become devolved to clinics and wards certain specialist procedures will become concentrated in fewer institutions—though not necessarily all in the same place. This should ensure the maintenance of high analytical and interpretative skills required to make them clinically useful.

Some of the benefits that may be expected from decentralisation are already apparent. Self calibrating blood gas analysers capable of being operated by comparatively unskilled medical staff have improved the quality of clinical care provided by intensive care units; the effects of blood glucose analysers on the management of diabetes and of non-invasive percutaneous reflectance bilirubinometers on neonatology represent just two tips of the iceberg. The day cannot be far