United Kingdom is 30 cases per 100 000 women. Unless one is advising routine diagnostic curettages in all 50-60 year old women, the intention to start oestrogen treatment does not create such a necessity. An argument can be made for a curettage every three to five years after oestrogen treatment, but an annual endometrial biopsy as recommended by Dr Craig does not even appeal to the private practitioner in me. The epidemic of endometrial carcinoma in the late 1970s occurred only in the USA and was clearly the result of inappropriate treatment without progestogen or proper supervision (of 2500 reported cases only seven had had "progesterone" rarely).² Now that the protective effect of oestrogen-progestogen preparations on the endometrium has been established,^{3,4} I doubt whether an annual curettage is justified even on a research level as such an investigation of 2000 patients over eight years in my research clinic has yet to diagnose a single case of endometrial carcinoma in patients whose treatment we had commenced and supervised.

No doubt the Family Planning Association will successfully spread its skills past reproductive life, but I would urge them to resist the temptation to create an expensive complexity out of a simple cost effective treatment.

JOHN STUDD

King's College Hospital,
London SE5

¹ Chakravati SK, Collins WP, Thom MH, Studd JWW. Relationship between plasma hormone profiles, symptomatology and response to oestrogen in women approaching the menopause. *Br Med J* 1979;i:983-5.

² Studd JWW, Thom MH. Oestrogens and endometrial cancer. In: Studd JWW, ed. *Progress in obstetrics and gynaecology*, vol 1. London: Churchill Livingstone, 1981:182-98.

³ Sturdee D, Wade-Evans T, Paterson MEL, Thom MH, Studd JWW. Relations between bleeding pattern, endometrial histology, and oestrogen treatment in menopausal women. *Br Med J* 1978; i:1569-648.

⁴ Paterson MEL, Wade-Evans T, Sturdee D, Thom MH, Studd JWW. Endometrial disease after treatment with oestrogens and progestogens in the climacteric. *Br Med J* 1980;280:822-4.

SIR,—Dr Gillian Craig's suggestions (25 June, p 2033) for setting up menopausal clinics in district hospitals and community health clinics are long overdue. Achieving her aims may be difficult but might entail shifting the emphasis of the conventional family planning clinic away from the provision of free, safe sex for teenagers towards that of providing mental, physical, and metabolic support for the menopausal woman. Society owes it to our middle aged women many of whom have suffered privations during the long years spent producing and nurturing their families only to become afflicted by unpleasant vasomotor symptoms, feelings of worthlessness, and, in old age, a knowledge of a fourfold greater chance of enduring mechanical failure of their skeletons than men of similar age.

Dr Craig's contention that the long term benefit of a menopausal clinic would be the prevention of fractures related to osteoporosis is an exciting but unproved possibility. To confirm and expand on this aspect of menopause clinics she would have to make some provision for the regular assessment of the skeletal state of those women attending her clinic. Failure to do this would leave her open to the same justified criticism levelled against gynaecologists who prescribe prophylactic iron and folic acid to pregnant women without making a single haemoglobin estimation.

Professor Cooke's suggestions on treating the menopause (25 June, p 2001) add authority to Dr Craig's proposals, but his alarmist claims about the prevalence of appreciable spinal osteoporosis might easily stampede the impulsive into unnecessary action, whereas his pessimism about the long term benefit of oestrogen treatment might well persuade the timid to shy away from treating

"oestrogen deficiency" osteoporosis, otherwise known as postmenopausal osteoporosis, with agents other than oestrogens.

Professor Cooke's premise that 25% of women over the age of 60 have spinal compression depends on what is meant by "spinal compression." Nordin¹ has stated that no more than 8% of 80 year old women have an appreciable degree of vertebral compression, and by that most of us not privy to Nordin's data assume the presence of either multiple vertebral crush fractures or at least one vertebral crush fracture associated with vertebral wedging and biconcavity elsewhere in the spine. Nordin readily admits that 50% of 75 year old women have some degree of vertebral wedging and biconcavity, but no more than 7% have vertebral compression.

Professor Cooke's other statement that it would be worth setting up a prospective controlled trial of calcium and vitamin D (1 α -hydroxyvitamin D₃) versus sequential oestrogen-progestogen treatment, overlooks previous reports^{2,3} which have shown that vitamin D treatment, whether taken alone or with calcium supplements, was either useless or positively harmful in terms of preventing loss of bone mineral from the appendicular skeleton. One report² suggested that patients taking either calcium supplements alone or calcium and vitamin D had a slower rate of progression of their spinal disease than those taking other forms of treatment, but in fact none of these results were significant except in so far as the spines of those patients taking vitamin D alone were significantly worse. Although this study used doses of vitamin D which were 20 to 100 times greater than the usual daily requirement, it would be important to establish that "sensible" doses of vitamin D had a beneficial effect in patients with spinal osteoporosis before subjecting postmenopausal women with normal skeletons to a potentially lethal bone cracker.

The funding of menopause clinics was an issue to which neither Dr Craig nor Professor Cooke addressed themselves. In the present climate of financial constraint funding would have to come from savings in other areas, and the running expenses would have to be met by cutting back on non-essentials.

In these respects the basic funding of a menopause clinic might well be achieved by changing the terms of reference of the present family planning clinic, so that most of its existing clientele could be transferred back to the general practitioner's own family planning service, reserving its own considerable skills for women with unusual contraceptive problems. The provision of extra items of service, such as suction curettage and measurement of bone mass, might be financed by economical prescribing in this area, where the most expensive proprietary preparations may cost as much as £1 a week, while simpler preparations actually cost less than £4 a year.⁴ Prudent prescribing could therefore reduce the potential cost of providing treatment alone to 1000 women from approximately £50 000 a year to £4000 a year.

I hope that district management teams will be encouraged to look critically at this area of medical practice and make some realistic provisions for dealing with this not inconsiderable problem, which until now has been swept under the bureaucratic carpet.

J M AITKEN

Essex County Hospital,
Colchester,
Essex CO3 3NB

¹ Nordin BEC, Horsman A, Marshall DH, Hanes F, Jakeman W. The treatment of post-menopausal osteoporosis. In: Barzel U, ed. *Osteoporosis II*. New York: Grune and Stratton, 1979:183-203.

² Nordin BEC, Horsman A, Crilly RG, Marshall DH, Simpson M. Treatment of spinal osteoporosis in postmenopausal women. *Br Med J* 1980;280:451-4.

³ Finn Jensen G, Christiansen C, Transbol I. Treatment of post-menopausal osteoporosis. A controlled therapeutic trial comparing oestrogen/gestogen, 1,25-dihydroxy-vitamin D₃ and calcium. *Clin Endocrinol* 1982;16:515-24.

⁴ Aitken JM. Postmenopausal osteoporosis. *Br Med J* 1982;285:1577.

SIR,—I would question the use of the Blatt menopausal index in the assessment of menopausal patients. Only two of the symptoms in Dr Gillian Craig's list (25 June, p 2033) have been verified as hormone dependent: flushes and sweats; and atrophic vaginitis, which might be a cause of dyspareunia.^{1,2} The use of this check list of "menopausal" symptoms may easily encourage the indiscriminate prescription of hormone treatment for patients who would not benefit from it.

JEAN COOPE

Bollington,
Macclesfield SK10 5JL

¹ Utian WH. The true clinical features of postmenopause and oophorectomy, and their response to oestrogen therapy. *S Afr Med J* 1972;46:732-7.

² Coope J, Thomson JM, Poller L. Effects of "natural oestrogen" replacement therapy on menopausal symptoms and blood clotting. *Br Med J* 1975;iv:139-43.

Whooping cough and pertussis vaccine

SIR,—The comments made by Miller and Ross^{1,2} and others^{3,4} in your columns about the whooping cough and pertussis vaccine controversy in general and myself in particular can be answered only by examining verifiable epidemiological data and reliable clinical reports. Either without the other is liable to be misleading.

The relevant period is from 1970 to 1982. Before 1970 it was agreed, even in your columns,^{5,6} that pertussis vaccine was "not very effective," in the words of the summary of a comprehensive national survey. After that, some people claimed⁷ that the vaccines were more effective because antigens 1-3 were included, but this claim, unconfirmed epidemiologically and refuted microbiologically,⁸ was not supported by the international expert panel of the World Health Organisation⁹ nor by one of the main manufacturers,¹⁰ who pointed out that these antigens had been included in their vaccine since 1964. Anyway, there was an extensive outbreak in 1970-1, with 51 deaths. Nothing was published between 1970 and 1974 to show that the vaccine was more effective, but there was growing concern about its safety to such an extent that by 1975 acceptance had dropped to less than 50% of eligible infants in most parts of Britain. There has been little change since then. About half of the children born in Britain since 1974 have not received pertussis vaccine, so that the child population is naturally divided into two halves in which the incidence and risks of whooping cough and the incidence of adverse reactions to pertussis vaccine and its efficacy can be compared.

Between 1969 and 1978 available data consisted mainly of notifications,¹¹⁻¹⁴ but since then there have been several prospective surveys in Britain¹⁵⁻¹⁹ and two retrospective ones in the United States^{20,21} of whooping cough and its complications. Unfortunately, the same cannot be said of adverse effects of the vaccine, but it is possible, from data in the National Childhood Encephalopathy Study²² and various other sources,²³⁻³¹ to estimate the incidence of severe toxic effects and hence to compare benefit with risk.

Table I summarises the data from these surveys. The four British surveys show broadly similar rates of vaccine acceptance, of secondary attack in household contacts of known cases, and of vaccine protection calculated from this. Most cases were in children older than 1 year. In the group aged 1-5 years the efficacy of pertussis vaccine was only 51-59%. Admissions to hospital were infrequent (0.8-8% of cases), as were deaths and encephalopathy.

The American surveys show, in contrast, a seemingly high attack rate (62-76%) in infants and much higher rates of death, encephalopathy (0.4-1.3%), and admission to hospital (57-60%). The general attack rate appears to be much lower, but this is because the American surveys were

restricted to babies and young children admitted to hospital.³² Vaccine protection is claimed to be about 82% (63-94%), which equates to the rate claimed in Britain if reliance is placed on notifications.¹¹⁻¹⁴ In other studies of household contacts in the United States^{21,33} the protection rate is much lower, as in Britain, although vaccine acceptance is about 95%. It is possible²¹ that the booster injections of vaccine given in the United States afford clinical protection to older children, but the high incidence of infections in infants throws doubt on the notion that herd immunity is thereby obtained since infants seldom contract infection except from older children, who are missing in both US surveys.

These data, all of which are verifiable, provide a basis for estimating risk-benefit by converting rates to prior probabilities and comparing the risks of disease with those of pertussis vaccine, as calculated from various sources,²³⁻³¹ including the National Childhood Encephalopathy Study.²² It is important to realise, as Miller and Ross pointed out,² that the estimate of attributable risk provided by the national childhood encephalopathy study is epidemiologically satisfying because controls are available for the cases notified by hospitals. But, equally, by no means all children reacting to vaccines are referred to hospital,³⁰ and in detecting rare events such as encephalopathy epidemiology is an inadequate instrument unless attention is paid also to relative incidences of relevant occurrences. There is no doubt that, in this respect, pertussis vaccine has a consistent record in the published work and in unpublished reports since 1933 of neurotoxic and other severe sequelae unmatched by any other vaccine, long before there was any adverse publicity about it in the media.

Since about 1950 the risk of death or of severe disease from whooping cough has been confined essentially to infants. Comparison of risks (table II) shows that if an infant contracts whooping cough (1:14-1:100) its risk of severe disease as indicated by admission to hospital (about 1:300-1:3000) is not very different from its risk (1:750-1:3000) of a severe reaction after pertussis vaccine. The average risk of encephalopathy with permanent brain damage after whooping cough (1:38 000) is within the range (1:25 000-1:110 000) of that reported after diphtheria, pertussis, and tetanus vaccination. The risk of death (1:130 000) after whooping cough (mainly unconfirmed) is not too different from that reported (1:100 000), although also unconfirmed, after pertussis vaccine in 1979-80.

On a population basis, the risk of any child contracting infection (1:14-1:100) varies considerably from time to time between persons and between and within places. About 25-35% of those who develop whooping cough have already received three injections of diphtheria, pertussis, and tetanus vaccine and, in older children, the proportion is higher.^{15,18} Encephalopathy with permanent brain damage after diphtheria, pertussis, and tetanus vaccine is variously estimated at 1:17 000-1:178 000,^{22,23,27,29,30,31,34} which is higher than that detected after whooping cough (1:300 000) in the National Childhood Encephalopathy Study.²¹

TABLE II—Estimates of risks of whooping cough and pertussis vaccine based on incidences of reported occurrences (numerator = number of occurrences reduced to unity; denominator = corresponding number of children at risk by age groups)

Incidence	Age (yrs)	High risk areas	Low risk areas	Averages*
Whooping cough	0-15	1:32	1:100	1:60 (1:100)
	1-4	1:16	1:50	1:27 (1:80)
	0-1	1:14	1:100	1:40 (1:117)
Admission to hospital	1-1	1:336	1:3060	1:547
Permanent brain damage	0-1	1:2000	0	1:38 000
				(NCES ²² 1:300 000)
Death	0-1	1:1300	0	1:12 600
	0-15	1:34 000	0	1:130 000
England and Wales	0-15			(1:560 000)
Scotland	0-15			(1:260 000)
All causes	0-1			(1:36)
After pertussis vaccine	0-2	CSM (1979-80)		(1:100 000)
After all other vaccines	0-2	CSM (1979-80)		(1:450 000)
	0-2	Britain ^{27,29,31}		1:17 000-1:54 000
	0-2	DHSS tribunals		(1:25 000)
Permanent brain damage after pertussis vaccine	0-2			(NCES ²² 1:110 000)
	0-2	West Germany ²³		1:60 000
	0-2	Prensky ³¹		1:178 000
Neurotoxic reactions	0-2	Britain, USA, Holland ^{24,25}		Range 1:750-1:3000
Systemic reactions		Britain, USA, Holland ^{24,25}		Range 1:2 -1:10

*From surveillance of 1978-9 outbreak in Britain/Glamorgan¹⁶ is taken as a high risk area, Guildford¹⁷ as a low risk area.

Estimates in brackets are calculated from notifications to OPCS Registrars General, or DHSS (Tribunals, NCES). OPCS = Office of Population Censuses and Surveys. CSM = Committee on Safety of Medicines. DHSS = Department of Health and Social Security. NCES = National Childhood Encephalopathy Study.

It is also higher than the risk of permanent lung damage reported in other surveys.^{35,36}

Clinical reports from various sources show that in the outbreaks of whooping cough in the 70s (1970-1, 1974-5, 1978-9) and in 1982 many babies with severe infections were admitted to hospital, confirming the fact also apparent in the American survey that current vaccination programmes do not necessarily protect infants. The death rate, however, has fallen steadily during this time. In 1982, only 14 deaths were certified in Britain and in these, as in all deaths since 1964, most were in infants below 6 months of age or vulnerable in other ways. Only two were confirmed bacteriologically (only 4% since 1964), but other pathogens (parainfluenza and respiratory syncytial viruses, *Pseudomonas* spp) were isolated from four. In the 1982 outbreak there was a considerable overlap between notifications of whooping cough and various respiratory viral infections, as in 1977-9.¹⁸ Continuous surveillance of whooping cough, among other infections of childhood, in Glasgow since 1974 shows that outbreaks and severity are much more intense in areas of stress and deprivation, where spread of infection within and beyond households is expedited by large numbers of children in crowded homes, playgroups, and schools.

All of this goes to show that whooping cough may be almost as prevalent now as in the 1960s in some parts of Britain and, apparently, in the United States. It is still a distressing disease, a threat to life in some infants, and a problem badly in need of resolute attack in the clinical, epidemiological, and public health sectors. There is good reason in the data to believe that if deprivation and

overcrowding can be abated the incidence and severity of the disease will diminish, but until this happens a vaccine is needed. The data show that the present vaccine does not control outbreaks. I believe that it is insufficiently effective and too risky to be used in mass vaccination programmes, although I would agree that it should be available to parents and doctors who want to use it so long as they are advised of its limitations. This constraint should be applied also to international programmes and no fault compensation introduced for victims of adverse reactions.

GORDON T STEWART

Department of Community Medicine,
Ruchill Hospital,
Glasgow G20 9NB

¹ Miller DL, Ross EM. ABC of 1 to 7: whooping cough. *Br Med J* 1982;284:1874.

² Miller DL, Ross EM. Whooping cough: a shot in the dark. *Br Med J* 1983;286:1817.

³ Valman HB. ABC of 1 to 7: whooping cough. *Br Med J* 1982;284:886-7.

⁴ Preston NW. Toxicity of pertussis vaccine. *Br Med J* 1982;284:1817-8.

⁵ Public Health Laboratory Service. Effectiveness of whooping cough vaccines before 1968. *Br Med J* 1973;259:62.

⁶ Anonymous. Vaccination against whooping cough. *Br Med J* 1974;359:40.

⁷ Preston NW, Stanbridge TN. Efficacy of pertussis vaccines: a brighter horizon. *Br Med J* 1972;344:52.

⁸ Holt LB, Spasovic V. The role of surface antigens in the protective potency of Bordetella pertussis suspensions. *J Gen Microbiol* 1968;11:19-21.

⁹ WHO Immunization against whooping cough. *WHO Chronicle* 1975;29:365-7.

¹⁰ Griffith AH. Reactions after pertussis vaccines: a manufacturer's experiences and difficulties since 1964. *Br Med J* 1978;3:809-15.

¹¹ Miller CL, Fletcher WB. Severity of notified whooping cough. *Br Med J* 1976;3:117-9.

¹² Noah ND. Attack rates of notified whooping cough in immunised and unimmunised children. *Lancet* 1976;3:129-32.

¹³ Whooping cough vaccination: review of the evidence by the Joint Committee on Vaccination and Immunisation. London: HMSO, 1977.

¹⁴ Baylith WR, Stewart GT. Epidemiological evaluation of immunisation and other factors in the control of whooping cough. *Lancet* 1976;3:471-4.

¹⁵ Stewart GT. Vaccination against whooping cough: efficacy versus risks. *Lancet* 1977;3:234-7.

¹⁶ Royal College of General Practitioners. Effect of low pertussis vaccination uptake on a large community. *Br Med J* 1981;282:23-6.

¹⁷ Grob PH, Crowden MJ, Robbins JF. Effect of vaccination on severity and suspension of whooping cough. *Br Med J* 1981;282:1925-8.

¹⁸ Stewart GT. Whooping cough in relation to other childhood infections in 1977-79 in the UK. *J Epidemiol Comm Health* 1981;35:139-45.

¹⁹ Public Health Laboratory Service and 21 Area Health Authorities. Efficacy of pertussis vaccination in England. *Br Med J* 1982;285:357-9.

²⁰ Pertussis: supplementary surveillance. *Morb Mort Wkly Rep* 1982;31:334-6.

²¹ Pertussis—Maryland, 1982. *Morb Mort Wkly Rep* 1983;32:297-300.

TABLE I—Surveillance of whooping cough in Britain and USA

	Britain				USA	
	1978-9	1978-9	1978-9	1978-80	1978-81	1982
	Glamorgan ¹⁶	Guildford ¹⁷	Glasgow ¹⁸	21 areas ¹⁹	42 States ²⁰	Maryland ²¹
Children at risk (% of total population)	72 300 (20%)	57 700 (42%)	236 700 (23%)	437 758(†)	50 000 000 (estimate)	†
No of cases (per 1000 children)	2 295 (32)	658 (11)	3 150 (13)	11 776(27)	1 277 (0.026)	79†
No of cases <1 year of age (% of cases)	243 (11)	†	358 (11)	70 (6)	791 (62)	60 (76)
No of cases admitted to hospital (% of cases)	(64 (3))	5 (0.8)	240 (8)	171 (1.5)	740 (60)	45 (57)
No of cases with encephalopathy (% of cases)	2 (0.1)	0	0	†	5 (0.4)	1 (1.3)
No of deaths	3 (0.13)	0	0	†	7 (0.55)	0
Bacteriological confirmation	39%	33%	33%§	25%	72%†	56%
Secondary attack rate (household)	41%	36%	37%	50%	†	†§
Vaccine acceptance (diphtheria, tetanus, and pertussis vaccine three times)	50%	43%	56%	57%	95%	69-96%
Vaccine protection-contacts 1-5 years	59%	53%	51%	59%	†	†

*Surveillance of 21 areas by the Public Health Laboratory Service of balanced groups of children aged 0-6 receiving diphtheria, tetanus, and pertussis vaccine three times or diphtheria and tetanus vaccine three times.

†Not shown.

‡Mainly in cases admitted to hospital.

§Three out of four cases had received diphtheria, tetanus, and pertussis vaccine three times.

- ¹¹ National Childhood Encephalopathy Study. Report by the Committee on the Safety of Medicines and the Joint Committee on Vaccination and Immunisation 1981. *Whooping cough*. London: HMSO, 1981:79-169.
- ¹² Ehrengut W. Über Konvulsive reaktionen nach pertussis schutzimpfung. *Dtsch med Wochenschr* 1974;11:2273-6.
- ¹³ Barkin RM, Pichichero ME. DPT vaccine: reactivity of commercial products. *Pediatrics* 1979;63:256-9.
- ¹⁴ Hopkins RS. Reactions to DPT by lot and manufacturers: results of a survey in Montana. *International symposium on pertussis*, 1978. Washington DC: US Government Printing Office, 1979:300-3.
- ¹⁵ Hannik CA, Cohen H. Pertussis experiences in the Netherlands. *International symposium on pertussis*, 1978. Washington DC: US Government Printing Office, 1979:279.
- ¹⁶ Stewart GT. Toxicity of pertussis vaccine: frequency and probability of reactions. *J Epidemiol Comm Health* 1979;33:150-6.
- ¹⁷ Cody CL, Baraff CJ, Cherry JD, Marcy SM, Manclark CR. Nature and rates of reactions associated with DPT and DT immunization in infants. *Pediatrics* 1981;68:650-9.
- ¹⁸ Report of the Advisory Panel. The collection of data relating to adverse reactions to pertussis vaccine. In: Reports from the Committee on the Safety of Medicines and the Joint Committee on Vaccination and Immunisation. *Whooping cough*. London: HMSO, 1981:27-75.
- ¹⁹ Stewart GT, Wilson J. Pertussis vaccine and acute neurological illness in children. *Br Med J* 1981;282:1968-9.
- ²⁰ Pollock TM, Morris J. A 7-year study of disorders attributed to vaccination in the NW Thames Region. *Lancet* 1983;1:753-7.
- ²¹ Stewart GT. Whooping Cough in the United States and Britain. *N Engl J Med* 1983;308:464-5.
- ²² Rauber A, Bruner B, Feldman W, Nahmias A, Elisei W, McCoan J. Pertussis—Maine, Georgia (USA). *Mor Mor Wkly Rep* 1977;26:307-8.
- ²³ Premsky AL. Pertussis vaccination. *Dev Med Child Neurol* 1974;16:539-40.
- ²⁴ Lees AW. Ataxia and bronchectasis in pertussis. *Br Med J* 1950;1:138-41.
- ²⁵ Bland JM, Holland WW, Elliott A. The development of respiratory symptoms in a cohort of Kent schoolchildren. *Bull Phys Path Resp* 1974;10:699-715.

Mass hysteria among schoolchildren

SIR,—We appreciate Minerva's comments (2 April, p 1151) on the press coverage of the outbreak of mass hysteria among schoolchildren (mainly girls) in Boston, Massachusetts. Her remarks were particularly apt as they coincided with a similar episode that occurred in the West Bank during the end of March and the first week of April. Three hundred and twenty three schoolgirls in the Jenin district complained of symptoms resembling those in the Boston epidemic—abdominal pain, nausea, and dizziness. They were joined by other schoolgirls in the ensuing days.

The outbreak received the attention of the media and was reported variously: the girls had been "poisoned by an unidentified chemical substance," were suffering from a "mysterious malady" caused by some "environmental irritant," or were suffering from "the inhalation of some kind of gas." A yellow powder found on a few windowsills was suspected of containing hydrogen sulphide but was subsequently identified as pollen.

Two teams of foreign experts carried out extensive investigations. The report prepared by an Australian epidemiologist and an Italian toxicologist was presented at the World Health Organisation 36th annual conference in Geneva on 11 May. It concurred with those of the team from Atlanta and the Israeli Health Ministry and concluded that there was no evidence of environmental toxins and that the epidemic was induced by "anxiety." Unfortunately, the findings were not awarded the same publicity as the events preceding them.

We hope that Minerva's recommendations that "all journalists be taught the features of hysteria and a few other medical basics" have been noted and will be adopted.

JEAN E FISHMAN

Negev District,
Beer Sheva,
Israel

A Y JOSEPH

Soroka Medical Centre,
Beer Sheva,
Israel

Smoking, lung function, and body weight

SIR,—Dr T Khosla and Dr C R Lowe (14 May, p 1578) point out that one should not infer from our data (22 January, p 249) that lung function is unaffected if there is no overt airflow obstruction. We were aware of this, as can be seen from reference four of our paper, which presents the detailed lung function results of the studied subjects.¹

We showed that about half of the smokers without spirometrically detectable airflow obstruction (forced expiratory volume in one second (FEV₁)/vital capacity >66%) had small airways disease, defined as an increased slope of the nitrogen alveolar plateau or an increased closing capacity, or both. These smokers (n=32) had significantly lower values of FEV₁ (2.98±0.44 litres, 0.63±0.08 litres/m²) than smokers without small airways disease (n=36). The latter had values of FEV₁ (3.34±0.43 litres, 0.71±0.07 litres/m²), as well as of all other indices, that were indistinguishable from those of the asymptomatic non-smokers (3.52±0.51 litres, 0.72±0.08 litres/m²). We had not considered it necessary to discuss these aspects, which were beyond the scope of our paper and had already been presented elsewhere.

Dr Khosla and Dr Lowe also say that the ratio of FEV₁ over the forced vital capacity is a misleading indicator of impaired lung function since forced vital capacity is also decreased in smokers who "may not make an all out effort to expel the air in [their] lungs during measurements of forced vital capacity for fear of coughing." Both forced vital capacity and FEV₁ are derived from an expiratory manoeuvre during which more than half the volume is expelled at high flow rates. It is not surprising, therefore, that in patients with airway obstruction forced vital capacity is less than the slow vital capacity, which is recorded at low flow rates. This explains the loss of sensitivity of FEV₁/forced vital capacity compared with FEV₁. In our study we related FEV₁ to the slow vital capacity. Values of vital capacity were similar in our groups with and without airflow obstruction (4.29±0.64 litres and 4.32±0.59 litres respectively).

Finally, in contrast to what Dr A Keys writes (4 June, p 1883), our observation, fully confirmed by Dr F Kauffmann (16 April, p 1280), shows more than just "the inverse relation between smoking and relative weight," "simply [reflecting] compulsion for oral satisfaction." Our data support the idea that only those whose lungs are being adversely affected by smoking tend to lose weight.

B NEMERY
N E MOAVERO
L BRASSEUR
D C STĂNESCU

Cardiopulmonary Laboratory,
Cliniques Universitaires Saint-luc,
Brussels

¹ Nemery B, Moavero NE, Brasseur L, Stănescu DC. Significance of small airways disease in middle-aged smokers. *Am Rev Respir Dis* 1981;124:232-8.

Doctors should learn 100 words of Punjabi or Hindustani

SIR,—Some three years ago, having many Asian patients and not liking a 10 year old boy translating his mother's comments on gynaecological problems, I decided to learn some of their languages (4 June, p 1827). I was encouraged by a Hindu colleague and approached the community relations department at Newcastle Civic Centre. It was most helpful, and one of the officers lent me a dozen books of his own to read covering the religions, cultures, and customs of the various ethnic groups of the Indian subcontinent. He also provided me with a teach yourself Hindi book and a teacher, who came to my surgery and gave me lessons. In less than six months I was able to manage a surgery.

My colleague also arranged for me to be introduced to the president of the Sikh temple, and these people encouraged me and helped me to improve my pronunciation. I practised in odd bits of free time. I was working full time as a principal in general practice at the time and must admit to giving

up a lot of my free time for a few months. I must emphasise that I learnt only nouns and verbs and phrases for medical use—I would have been no use discussing the weather. The spin off is that the patients very much appreciate your efforts to meet them and understand their customs and will only too readily explain what is the right thing to do in the case of, for example, death. My patients were grateful and pleased that I had tried to reach them, and the good will generated was enormous. I am sure it was very good for the practice.

Asian community nurses also can give you help with colloquial phrases and feedback on problems in the practice. If one shows willing there is plenty of help to be had for free. I found a degree of friendship and kindness among Asians of all faiths that I had not known before; and the object of the exercise, the care of non-English speaking Asian women, became very much easier. They gave me trust which was not there before, presumably because I had become known on the Asian grapevine and because they knew that I was trying to treat them bearing their religious and cultural backgrounds in mind.

P A ELLIOTT

Weardale,
Co Durham

Hypnotics and anxiolytics

SIR,—Dr N Hockings and Dr Brian R Ballinger (18 June, p 1949) rightly say that the important difference among benzodiazepines is duration of action, but I think that their categorisation into short acting and all others does not quite reflect the present position and that it would be better to distinguish the very short acting drugs. Into this last category comes triazolam (Halcion), for which the authors quoted an elimination half life of five hours, but most publications give two to four hours—for example, Jochemsen *et al* now report a mean of 2.4 h.¹ Triazolam therefore would be better not loosely classed with lormetazepam (Noctamid), which has a half life of 10 hours, four times as long.

Last year, Dr K Morgan and I reported that triazolam 0.5 mg nightly for three weeks led to increasing daytime anxiety among middle aged poor sleepers.² Hostile letters were provoked in your columns. From the USA, however, Kales *et al* now confirm that during a second week of triazolam 0.5 mg nightly, "tension or anxiety increased above baseline."³ Like ourselves, they interpret the findings as daytime withdrawal phenomena. Moreover, Kales *et al* report early morning insomnia as a positive consequence of triazolam and its very rapid elimination—that is, wakefulness as a late night withdrawal phenomenon. Most doctors are familiar with morning insomnia, anxiety, and tension feelings provoked by the ingestion of large doses of another very rapidly metabolised hypnotic and anxiety relieving agent, alcohol.

In your journal in 1973 my colleagues and I wrote of the same phenomenon as a consequence of accustomed amylobarbitone sodium: a "latenight withdrawal rebound as the bedtime dose of drug is metabolised," providing a reason why patients may find "confirmation of their supposed need for a hypnotic at a time when the hypnotic itself is perpetuating the poor sleep."⁴

The very long acting, cumulative hypnotics like flurazepam (Dalmane), referred to by