6000 sets of notes which had complete information from birth or from 1948. Inadequate records will lead to further underestimation of the lifetime prevalence of this condition.

Whatever the methodological variations, epidemiological studies on epilepsy seem to be unanimous in supporting the observation that health care for this group is, as they say, "insouciant." Irregular attendance and poor surveillance seem to be the norm. We are currently working with local general practitioners to investigate ways in which the health care of people with epilepsy might be improved, and we are paying particular attention to this problem. Aspects which do not seem to have been covered by previous studies are the patient's own feeling of handicap, motivation to look for support, and satisfaction with the help already received. The personal interview advocated by Dr Goodridge and Dr Shorvon is one possible approach though there are other techniques, such as a properly conducted postal survey, which could allow such data to be collected. As far as we know, this approach has not been tested in patients with epilepsy.

We would agree with the comparisons drawn by Dr Goodridge and Dr Shorvon between diabetes and epilepsy in that both are common, chronic diseases requiring long term follow up. Rather than accepting that the care of people with epilepsy should be developed in regional specialist centres, however, it may be that the parallel with diabetes could be extended. As with patients with diabetes, general practitioners may feel that they wish to become more concerned with the long term care of patients with epilepsy with recourse to hospital services only for specific problems.

M H BURTON
RHYS WILLIAMS

University Department of
Community Medicine,
Addenbrooke's Hospital,
Cambridge CB2 2QQ

¹ Lloyd Jones A. Medical audit of the care of patients with epilepsy in one group practice. *J R Coll Gen Pract* 1980;30:396-400.

SIR,—The article by Dr D M G Goodridge and Dr S D Shorvon contains the sentence (p 646): "Valproate was the only other drug to have been used in more than 10% of patients and, perhaps because of its active marketing, appears to have been often chosen by the general practitioner as initial treatment in cases that might be considered inappropriate—for example, partial or symptomatic epilepsies."

Turnbull *et al*, comparing phenytoin and Epilim (sodium valproate) in the treatment of partial seizures, found no difference in the efficacy of the two drugs.¹ Covanis *et al* obtained complete control of seizures in 47% of partial epilepsies using sodium valproate.² Kugler makes the following statement referring to sodium valproate: "We have for a long time successfully treated patients with complex partial seizures" and later: "Its value appears to have been well established so that its use should certainly not be confined to the generalised epilepsies."³

There is therefore documented evidence of the value of valproate in the treatment of partial epilepsies.

R H GOSLING
Medical director

Labaz Sanofi UK Limited,
Wythenshawe,
Manchester M23 9NF

¹ Turnbull DM, Rawlins MD, Weightman D, Chadwick DW. A comparison of phenytoin and valproate in previously untreated adult epileptic patients. *J Neurol Neurosurg Psychiatry* 1982;45:55-9.

² Covanis A, Gupta AK, Jeavons PM. Sodium valproate: monotherapy and polytherapy. *Epilepsia* 1982;23:693-720.

³ Kugler J. Treatment of partial epilepsies. *Neurochirurgia (Stuttgart)* 1981;24, suppl:284.

SIR,—The article by Dr D M G Goodridge and Dr S D Shorvon (3 September, p 641) has not done much to resolve the controversy concerning the prevalence of epilepsy. Though epidemiological studies have shown wide variation in prevalence rates (3 to 6/1000), depending on the operational criteria for epilepsy and method of ascertainment, there is some degree of consistency regarding "active" cases.

Zielinski¹ drew attention to the fair agreement in several studies of the prevalence of active cases of epilepsy¹: 3.6 in Rochester,² 3.4 in Iceland,³ and 3.4 in England.⁴ Previous surveys in general practice in England and Wales have yielded prevalence rates quite close to those mentioned above: 3.3/1000 based on information from 106 general practices,⁵ and 4.2/1000 for "chronic epilepsy" in the survey of 14 general practices in south east England,⁶ though the authors gave 6.2/1000 for all cases, including subjects with single afebrile seizures; the rate of 4.2/1000 was also obtained by the research committee of the College of General Practitioners,⁷ excluding non-active cases.

Though Dr Goodridge and Dr Shorvon criticised previous investigations for "poorly defined diagnostic criteria," their definition of active epilepsy ("a seizure" within 24 hours of the survey) does not really conform to that generally accepted. We are forced to consider the basic argument, whether a single non-febrile seizure necessarily constitutes epilepsy, and if it is logical to include patients taking prophylactic anticonvulsant drugs after a single fit in epidemiological studies, merely on the grounds that they were taking antiepileptic medication when interviewed.

According to the criteria of Gunn and Fenton,⁸ quite similar to those adopted by Hopkins and Scambler,⁴ epilepsy is diagnosed if a subject presents a history of at least three attacks in any two year period, and the continuation of anticonvulsant treatment if the period did not immediately precede the time of examination. Prevalence rates of 5.3/1000 and 5.7/1000 for active cases⁹ (whether diagnosed or not) illustrate the methodological difficulties in research into epilepsy.

JAMES EDEH

King's College Hospital,
London SE5 9RS

¹ Zielinski JJ. Epidemiology. In: Laidlaw J, Richens A, eds. *A textbook of epilepsy*. Edinburgh, London, Melbourne, New York: Churchill Livingstone, 1982:16-33.

² Kurland LT. The incidence and prevalence of convulsive disorders in a small urban community. *Epilepsia* 1959;1:143-61.

³ Gudmundsson G. Epilepsy in Iceland. *Acta Neurol Scand* 1966;43, suppl 25:46-56.

⁴ Hopkins A, Scambler G. How doctors deal with epilepsy. *Lancet* 1977;1:183-6.

⁵ Logan WPD, Cushion AA. *Morbidity statistics from general practice. Vol 1 (General) GRO studies on medical and population subjects*. London: HMSO, 1958.

⁶ Pond DA, Bidwell BH, Stein L. A survey of epilepsy in fourteen general practices I. Demographic and medical data. *Psychiatry Neurologia Neurochirurgia* 1960;63:217-36.

⁷ Research Committee of the College of General Practitioners. A survey of the epilepsies in general practice. *Br Med J* 1960;ii:416-22.

⁸ Gunn J, Fenton GW. Epilepsy in prisons: a diagnostic survey. *Br Med J* 1969;iv:326-8.

⁹ Hauser WA. Epidemiology of epilepsy. In: Schoenberg BS, ed. *Advances in neurology. Vol 19. Neurological epidemiology: principles and clinical application*. New York: Raven Press, 1978.

Mineral content of the forearms of babies

SIR,—Though it may be true that the bone mineralisation of the fetal skeleton is not related to vitamin D deficiency in Asian women in Great Britain, Dr P Congdon and others (16 April, p 1233) have not convincingly documented this with photon absorptiometry. To our knowledge the use of ²⁴¹Am has not been shown to be either accurate or precise in the determination of bone mineral content in newborn infants. Given the low bone soft tissue contrast of ²⁴¹Am it is important that this documentation be given.

Dr Congdon and his colleagues found a correlation of only 0.46 between birth weight and "forearm" bone mineral content. Others have previously found a much higher correlation (0.89, 0.82) between bone mineral content (radius) and birth weight using ¹²⁵I.^{1,2} What is the reproducibility in infants using the technique of Dr Congdon? The use of arbitrary units is very confusing, and their conclusions do not seem justified until their technique has been validated. If there are in fact technical limitations of the ²⁴¹Am based measurements at the very low bone mineral content of newborn infants then the results of this interesting study must be questioned.

FRANK R GREER
RICHARD B MAZESS

Department of Pediatrics and Medical Physics,
University of Wisconsin Center for Health Sciences,
Madison,
Wisconsin, USA

¹ Minton SD, Steichen JJ, Tsang RC. Bone mineral content in term and preterm appropriate-for-gestational-age infants. *J Pediatr* 1979;95:1037-42.

² Greer FR, Lane J, Weiner S, Mazess RB. An accurate and reproducible absorptiometric technique for determining bone mineral content in newborn infants. *Pediatr Res* 1983;17:259-62.

Effect of daily oral omeprazole on 24 hour intragastric acidity

SIR,—Dr R P Walt and others (2 July, p 12) studied the effect of 30 mg omeprazole on intragastric acidity in patients with duodenal ulcers. We have examined the effect of 40 mg omeprazole, given in the morning for five days to eight healthy volunteers. Using similar methods to those of Dr Walt and his colleagues we measured intragastric pH at hourly intervals during a 24 hour period on two occasions: (a) baseline, before administration of omeprazole, and (b) on day 5 of omeprazole treatment.

Omeprazole had a dramatic effect on gastric acidity. On the baseline day the median pH was 1.9 (range 1.3-3.1 (SD 0.6)) and on day 5 the median pH was 5.0 (range 4.4-6.1 (SD 0.5)). The difference was highly significant ($p < 0.001$; Wilcoxon rank test). The mean hydrogen ion activity on each study day was 26.15 (SEM 6.38) mmol/l with placebo and fell to 1.11 (± 0.24) mmol/l with omeprazole (paired t test, $p < 0.001$), and the range in acid secretion during the 24 hours varied from 0.18 to 58.12 mmol/l with placebo to 0.002-6.76 mmol/l with omeprazole. These results in normal subjects are similar to those reported by Dr Walt and his colleagues in patients with ulcers.

The reduction in gastric acidity with omeprazole is substantially greater than that achieved with therapeutic doses of histamine H₂ receptor antagonists. It is therefore not

surprising that Gustavsson *et al* have found that omeprazole in doses of 20-60 mg a day is highly effective in healing duodenal ulcers.¹

PETER J PRICHARD
NEVILLE D YEOMANS
GEORGE W MIHALY
D BRIAN JONES
RICHARD A SMALLWOOD
WILLIAM J LOUIS

Gastroenterology Unit and
Department of
Clinical Pharmacology,
Austin Hospital,
Heidelberg 3084,
Victoria,
Australia

¹ Gustavsson S, Adami H, Loof L, Nybers A, Nyren O. Rapid healing of duodenal ulcers with omeprazole: double-blind dose-comparative trial. *Lancet* 1983; ii:124-5.

Parathyroid hormone and 25-hydroxyvitamin D concentrations in the elderly

SIR,—The conclusions drawn by Ms M M Petersen and others (20 August, p 521) are invalid, principally because the authors did not adequately define the renal function of their patients other than in a simplistic manner.

In the elderly, who often have a reduced muscle mass, plasma creatinine concentrations may be normal or minimally raised in the presence of a major reduction in glomerular filtration rate. It is unfortunately necessary, when dealing with any age group, but particularly the elderly, to measure creatinine clearance so as to avoid this well recognised pitfall, and this was done in one study.¹ Thus, in the elderly the skeletal muscle mass may be reduced to one quarter of normal, so the plasma creatinine concentration will be disproportionately low and may even be in the normal range despite the presence of advanced renal failure. Also, many workers have shown that in the elderly there is an age dependent reduction in glomerular filtration rate, and it is thus imperative not to assume that renal function is normal because of normal serum creatinine concentrations. Even in the so called normal elderly group (mean age 75.1 years) of Ms Petersen and her colleagues parathyroid hormone concentrations were lower than in either of the much older groups of patients (mean ages 80.9 and 83.3 years), perhaps supporting my contentions.

I would recommend therefore that Ms Petersen and colleagues repeat their study with 24 hour urine creatine clearances, and then perhaps more meaningful results will be obtained. Perhaps they will even agree with my observations.

G M BERLYNE

Downstate Medical Center,
State University of New York

¹ Berlyne GM, Ben Ari J, Kushelovsky A. The aetiology of senile osteoporosis; secondary hyperparathyroidism due to renal failure. *Q J Med* 1975;44: 505-21.

* * We sent a copy of this letter to the authors, who reply below.—ED, *BMJ*.

SIR,—Those of us who spend our lives working with the elderly are well aware of the pitfalls in defining renal function in terms of serum creatinine concentrations.¹ As we explained in our paper, that is why "impaired renal

function" was defined primarily by a raised blood urea concentration. In elderly patients such as we studied, blood urea values are highly correlated with glomerular filtration rate measured by a radioactive chromium labelled edetic acid technique.² Estimates of renal function which rely on urine collections are most unsatisfactory in elderly subjects; this is presumably the reason why Dr Berlyne and others could measure creatinine clearance in only nine of the 16 old people whom they studied.³

The correlation between circulating immunoreactive parathyroid concentrations and age which they reported was very strong ($r=0.92$), though based on only 26 subjects. We found a rather weak correlation of immunoreactive parathyroid hormone with age in our study of 119 subjects; as we state, circulating immunoreactive parathyroid hormone concentrations in our normal old subjects were lower than in normal young adults and were raised only in elderly inpatients (with or without "impaired renal function"). These contradictions cannot be explained away on the basis of age-related changes in body composition or renal function.

Our study showed what it was designed to detect—namely, an interrelation between circulating immunoreactive parathyroid hormone and 25-hydroxyvitamin D concentrations in sick and normal elderly people. Had we wished to test the hypothesis that age-related decline in renal function leads to increasing secretion of immunoreactive parathyroid hormone, we would have measured biologically active parathyroid hormone (not a portion of the molecule cleared by glomerular filtration) and glomerular filtration rate (using radiolabelled edetic acid, not urine collections) in an adequate number of normal subjects right across the adult age range. The study by Dr Berlyne and others³ fulfils none of these criteria.

R S BRIGGS
MERYL M PETERSEN
M R P HALL

Department of Geriatric Medicine,
Faculty of Medicine,
Southampton General Hospital,
Southampton SO9 4XY

¹ Petersen MM, Hall MRP, Briggs RS. Blood urea, serum creatinine and anthropometry in the elderly. *Clin Sci* 1983;65:13P.

² Denham MJ, Hodgkinson HM, Fisher M. Glomerular filtration rate in sick elderly inpatients. *Age Ageing* 1975;4:32-6.

³ Berlyne GM, Ben Ari J, Kushelovsky A. The aetiology of senile osteoporosis: secondary hyperparathyroidism due to renal failure. *Q J Med* 1975; 44:505-21.

Standardisation of oral anticoagulant treatment

SIR,—In his valuable and timely review (8 October, p 1000) Dr N K Shinton indicates that changing from the established British ratios to the international normalised ratios would present little disturbance as the values are so close.

The data on the international normalised ratios corresponding to British ratios provided by Dr Shinton were based on a preliminary calibration performed in this laboratory. Results with the international reference preparation were correlated with those of Manchester comparative reagent, the routine reagent in over 90% of British hospitals. The complete data from this calibration show that the agreement between Manchester reagent

and the international reference preparation is in fact even closer, as the accompanying table shows. Within the therapeutic range of 2.0 to

Comparison of British ratios and international normalised ratios

Clotting time (s)	British ratio	International normalised ratio
24.0	2.0	2.0
25.0	2.1	2.1
26.0	2.2	2.2
27.0	2.2	2.2
28.0	2.3	2.3
29.0	2.4	2.4
30.0	2.5	2.5
31.0	2.6	2.6
32.0	2.7	2.7
33.0	2.8	2.8
34.5	2.9	2.9
36.0	3.0	3.0
38.0	3.2	3.2
40.0	3.3	3.3
44.0	3.7	3.7
47.0	4.0	3.95
50-60	4.2-5.0	4.15-4.9

*Recommendation of British Anticoagulant Panel.³

4.0 ratio advocated by the British system for anticoagulant control, these values and international normalised ratios are interchangeable. Clinicians at home and abroad may therefore apply international normalised ratio to their patients stabilised on oral anticoagulants with the confidence in safety and effectiveness of this range derived from cumulative clinical experience and therapeutic trials.^{1,2}

L POLLER

Withington Hospital,
Manchester M20 8LR

¹ Taberner DA, Pollar L, Burslem RW, Jones JB. Oral anticoagulants controlled by the British comparative thromboplastin versus low dose heparin prophylaxis of deep vein thrombosis. *Br Med J* 1978;i:272-4.

² Hull R, Hirsh J, Jay R, *et al*. Different intensities of oral anticoagulation therapy in the treatment of proximal vein thrombosis. *N Engl J Med* 1982;307: 1676-81.

³ Blackburn EK. Long term anticoagulation therapy. *Prescriber's Journal* 1977;17:73-4.

Substitution of Maloprim for dapsone

SIR,—We have recently seen two patients undergoing treatment for dermatitis herpetiformis in whom Maloprim was inappropriately substituted for dapsone. The antimalarial preparation Maloprim contains 12.5 mg of pyrimethamine as well as 100 mg of dapsone and is normally prescribed in a dose of one tablet a week. Pyrimethamine is a diaminopyrimidine with a half life of about four days. As the dose of dapsone required to control dermatitis herpetiformis may be up to 400 mg a day considerable overdoses of pyrimethamine may be given with the risk of severe megaloblastic anaemia.

There appear to be two reasons for the substitution taking place in our patients. Firstly, Maloprim appears in *Monthly Index of Medical Specialties* but dapsone does not because there are no proprietary brands of the pure drug. Secondly, the entry under dapsone in the *British National Formulary* relates only to the treatment of leprosy and makes no mention of dermatitis herpetiformis. Attention was drawn to the problem of substitution two years ago,¹ and one of our patients had been warned of this possibility as a result. We hope that further difficulties may be avoided by appropriate changes in the *British National*