Final guidance issued on Herceptin after appeal rejected

Caroline White *London*

The National Institute for Health and Clinical Excellence (NICE) has issued its final guidance to English and Welsh health service trusts for the use of trastuzumab (Herceptin) for the treatment of early stage breast cancer in women with HER2 positive disease and free of heart disease.

The final guidance is based on an initial draft published earlier this year (*BMJ* 2006; 332:1409). It recommends that:

- Herceptin be provided at three weekly intervals for a year or until disease recurrence—whichever is sooner—after surgery, chemotherapy, and radiotherapy, if applicable
- Heart function be assessed before treatment starts and repeated quarterly in women

whose left ventricular ejection fraction drops 10%

- Herceptin be avoided in women with a left ventricular ejection fraction of 55% or less, heart disease, including arrhythmias, or poorly controlled high blood pressure.

The move, which comes three months after the drug was licensed for use in Europe, follows rejection of an appeal against the institute's draft proposals, made by Newbury and Community Primary Care Trust at the end of July.

The decision will affect about one in five of the 40 000 women diagnosed with breast cancer every year in the United Kingdom, according to figures from the cancer charity CancerBACUP.

Newbury and Community Primary Care Trust launched its

appeal on the grounds that the institute had "prepared guidance which is perverse in the light of evidence," one of the three qualifying criteria against which the institute can be publicly challenged on its decisions.

The trust claimed that several aspects of the proposed guidance were unclear. These included the institute's failure to specify exactly which patients would benefit most from treatment with the drug, given its cost and potential side effects, and the length of time patients should be treated after surgery, chemotherapy, or radiotherapy.

Implementation of the guidance "would not be in the best interests of the whole population for whose health care the PCT [primary care trust] has responsibility," it said.

But the appeal panel said that the institute had acted on the basis of the available evidence, intending its guidance to be prospective rather than retrospective, and had taken account of NHS resources. The appeal was rejected.

Richard Mills, the interim director of service improvement at Berkshire West Primary Care Trusts, said, "We welcomed the opportunity to submit our case, and we accept the ruling."

Derryn Borley, head of cancer support at CancerBACUP, said that she thought the decision might place some trusts under financial pressure.

"PCTs are far too small to be commissioning for these expensive drugs," she said. "Maybe cancer networks would be better placed to do this." □

Male circumcision is not a panacea for HIV prevention

Jocelyn Clark *Toronto*

Male circumcision was a hot topic at the 16th international conference on AIDS, on 13-18 August, in Toronto, accompanied by passionate debate on both sides.

Delegates heard from scientists that new HIV infections in men could be substantially reduced and million of lives saved if male circumcision were to be introduced. Critics argued that male circumcision will not be the quick fix for HIV prevention unless cultural beliefs are included in the equation. The World Health Organization and UNAIDS have so far refused to endorse male circumcision as a prevention tool until more evidence is produced.

Last year the first clinical trial of male circumcision was halted

prematurely because early results showed that circumcision gave men a 61% protective effect against HIV infection compared with men who weren't circumcised (*PLoS Medicine* 2005;2:e298). All eyes are on two other trials, one in Uganda and one in Kenya, which are expected to end in 2007.

New data from the Kenya trial were presented at the conference. Half of the 2800 previously uncircumcised men aged 18-24 years who were recruited to the study were circumcised. At the current midpoint of the trial, the overall incidence of HIV is 1.8%, below the 2.5% predicted in uncircumcised controls. Although the investigators are still blinded as to whether the men were circumcised or

A Masai Moran circumcision ceremony in Kenya

circumcision yields "a 50% protective effect."

Another presentation at the conference reported that the rate of new HIV infections was two thirds higher in uncircumcised men in a cohort of rural Kenyan men who had been followed for two years. Researchers from Yale University said that 52 000 lives would be saved if rates of circumcision in men were raised by just 20% a year for five years. A cost effectiveness study estimated that circumcision would cost about \$55 (£29; €43) per adult male, but that cost would be offset by savings of about \$2400 in future medical costs for every averted HIV infection. PH

not, Robert Bailey of the University of Illinois said that the incidence of HIV infection in all men in the study indicates that

See News Extra on bmj.com for a report from the 16th international conference on AIDS in Toronto. The report, *New Approaches to HIV Prevention: Accelerating Research and Ensuring Future Access*, is available at www.kff.org.