In brief

Access is granted to genetic database of flu viruses: The US Centers for Disease Control and Prevention have opened up access to a database of genetic blueprints for over 650 genes for influenza viruses. The move aims to encourage flu research by giving researchers around the world access to data on virus samples obtained in the United States.

Juice drinking may lower risk of Alzheimer's: Drinking three or more glasses of fruit or vegetable juice a week may lower the chances of developing Alzheimer's disease, say US researchers. A decade long study of the cognitive function of 1836 adults without dementia found that that regular juice drinkers were 76% less likely to develop the disease than those drinking one serving or less each week (*Am J Med* 2006;119:751-9).

Nurses too busy to help feed elderly inpatients: A survey of 500 nurses for the UK charity Age Concern shows that nine out of 10 said they were often too busy to help elderly hospital patients with eating. The survey is part of the charity's new campaign "Hungry to be Heard" (www.ageconcern.org.uk), which aims to end malnutrition among elderly inpatients.

NSAIDs may cut risk of enlarged prostate: Taking aspirin or other non-steroidal anti-inflammatory drugs daily may prevent or delay benign prostatic hyperplasia, claim US researchers in the *American Journal of Epidemiology* (doi: 10.1093/aje/kwj258). The 12 year study of almost 2500 middle aged men found that the risk of developing an enlarged prostate was halved among those taking the drugs daily.

Human Tissue Act comes into force: From 1 September 2006 all services and organisations in the United Kingdom that store or display human tissue must be licensed by the Human Tissue Authority, to comply with the new legislation. Consent must be given for use or storage of all body parts, organs, and tissues from living and dead people. See www.hta.gov.uk.

England's cancer plan needs an overhaul, think tank says

Michael Day *London*

The NHS's national cancer plan must be thoroughly overhauled to allow for an ageing population and rapidly emerging medical technology, the health think tank the King's Fund says.

Its new report on the future of cancer services in England, which was commissioned by Cancer Research UK, also warns that structural changes within the NHS—particularly the implementation of the payment by results system—will have a big effect on care.

However, it is mainly demographic change and medical advances that have done most to render the current plan—which

was launched in 2000—outdated, the report says.

With the proportion of people aged over 65 likely to grow from 16% of the population in 2004 to 23% by 2031 and the rapid development of new treatments, better screening, and earlier diagnosis, many more people will be developing and surviving cancer, say the authors, and the extra cost will have to be budgeted for.

Already, the report notes, the total number of new cases of cancer is increasing by 1.4% each year.

"The situation is changing very rapidly, and we need to

have this discussion now," said Rebecca Rosen, the lead author of the report and senior research fellow in health policy at the King's Fund.

In addition, Dr Rosen said, "a raft of changes" to the NHS in the past five years meant that changes to the plan were necessary. "Foundation hospitals, private providers, and payment by results all mean that we have to rethink the cancer networks," she said.

Dr Rosen said that payment by results, for example, might provide hospitals with financial incentives to continue treating patients in wards when they could be cared for effectively and more cheaply in smaller hospitals or in the community.

Future Trends and Challenges for Cancer Services in England is available at www.kingsfund.org.uk.

Incidence of childhood cancer rises

Roger Dobson *Abergavenny*

The overall incidence of childhood cancer in Europe has been rising at a rate of 1.1% a year on average, a new study shows.

The researchers found that from 1978 to 1997 the age standardised incidence increased from 120 cases per million children to 141 (*European Journal of Cancer* 2006;42:1961-1).

Although the rise may be partly the result of changes in diagnosis and registration of cases, the authors say that the size of the increase and other factors point to other causes.

"The patterns and magnitude of these increases suggest that other factors—for example, changes in lifestyle and in exposure to a variety of agents, have contributed to the increase in childhood cancer in the recent decades," write the authors, from the University of Mainz, the International Agency for Research on Cancer, and other centres.

The researchers used data from the automated childhood cancer information system project to look at trends in the incidence of cancer in children from 1978 to 1997. The data came

from 33 cancer registries in 15 European countries and involved almost 80 000 cases.

Their results show that over the 20 year period the incidence increased by an average of 1.1% a year and that in each five year period the incidence was higher than in the preceding one. The trend was seen in all geographical areas and in most of the disease groups, ranging from an average rise of 1.8% a year for soft tissue sarcomas to 0.6% for leukaemias. No change was seen in the incidence of bone tumours, hepatic tumours, and retinoblastoma.

The authors point out that the average increase was greater among girls than boys, although

the baseline cancer incidence was higher among boys.

The authors say there are probably many possible risk factors that are not well understood. Possibilities include increased mobility; changing exposure to a variety of artificial and natural agents; changes in lifestyle; maternal diet; and reproductive history, pregnancy, and perinatal factors.

"Changes in genetic factors cannot be excluded, due to improved medical care, which allows affected individuals reaching the reproductive age and possibly transferring predisposition to certain cancers," the authors wrote.