Where next for China?

Rising inequalities in health and wealth are biggest challenge

NG HAN GUAN/AP/REUTERS

EDITOR—I write with reference to the editorial by Liu.¹ Life expectancy in China doubled from 35 to 71 years between 1949 and 2001.² China's human development index has been rising steadily, and it ranked 85th among countries with an index of 0.755 in 2003.²

According to China's human development report, the national Gini coefficient for income distribution rose to 0.45 in 2002 from 0.3 in 1982.² The average income of the highest income tenth is 11 times that of the lowest tenth.²

Disparities in health indicators between urban and rural areas are widening. Life expectancy is 75.2 years for urban residents and 69.6 for rural residents.² The human development index is 0.816 in urban areas and 0.685 in rural areas. Rural infant mortality at 34% and maternal mortality at 61.9 per 100 000 births are nearly double that in urban areas (14% and 33.1/100 000, respectively). Rural areas have only 2.4 medical personnel per 1000 residents, compared with 5.2 per 1000 residents in urban areas.

Regional disparities are widespread, with the western regions faring the worst. Life expectancy for rural residents is less than 65 in Tibet, Yunnan, and Guizhou. Maternal mortality is also highest in the western regions. Tibet ranked lowest in terms of life expectancy (65.81 years), literacy (47%), and human development index (0.562).² The region has the lowest number of doctors and hospital beds in China.

Only 15% of the entire population of China can afford medical insurance.² Only

2% of rural migrant workers currently have unemployment insurance.² Coercive public health policies include forced abortions.³

China's ambitious aim of building a society in which people are not rich but have adequate food, clothing, and basic necessities for a decent life² will require addressing these widening disparities, prioritising health opportunities for all, and acknowledging the inherent, instrumental and constructive⁴ roles of freedom in achieving these goals.

Sonal Singh *instructor in medicine*
Department of Medicine, Wake Forest University,
Winston-Salem, NC 27157, USA
sosingh@wfbmc.edu

Competing interests: None declared.

- 1 Liu T. China's health challenges. *BMJ* 2006;333:365. (19 August.)
- 2 United Nations Development Program. *China human development report 2005*. http://hdr.undp.org/docs/reports/national/CPR_China/China_2005_en.pdf (accessed 17 Aug 2006).
- 3 Watts J. Chinese officials accused of forcing abortions in Shandong. *Lancet* 2005;366:1253.
- 4 Sen A. *Development as freedom*. Oxford: Oxford University Press, 1999.

Keep to the basic principles of critical appraisal

EDITOR—In the review by Liu et al of treatment of hyperbaric oxygen, several methodological issues need to be addressed.¹ Alternate allocation to treatment group, or methods based on patient characteristics such as date of birth, order of entry into the clinic, or day of clinic attendance are not reliably random. Such allocation sequences are predictable and reduce the guarantee that no potential subjects have been excluded by foreknowledge of the intervention.² These trials should be removed in a sensitivity analysis as well as the trials not reporting "random" in the methods section. Authors should be contacted for clarification if inadequate reporting is thought to be the problem.

Using simple randomisation methods often leads to the sample size being similar in the two groups and inadequate reporting of the baseline characteristics.³ Attempts should be made to compare the baseline characteristics of trials where allocation concealment is unclear or not used. Trials not reporting baseline characteristics should be removed in the sensitivity analysis. Trials that report adequately concealed allocation yield larger estimates of treatment effects ($P < 0.001$) than trials with inadequately concealed or unclear allocation. Odds ratios

can be exaggerated by 41% for inadequately concealed trials and by 30% for unclearly concealed trials (adjusted for other aspects of quality).⁴

If Chinese medical literature is to be a rich source of evidence then reporting of the basic principles of critical appraisal and analysing high versus low quality data remain important. Concentrate on the studies that are judged to be valid.

Carl J Heneghan *clinical research fellow*
Centre for Evidence-Based Medicine, Department
of Primary Health Care, University of Oxford,
Oxford OX3 7LF
carl.heneghan@dphpc.ox.ac.uk

Competing interests: None declared.

- 1 Liu Z, Xiong T, Meads C. Clinical effectiveness of treatment with hyperbaric oxygen for neonatal hypoxic-ischaemic encephalopathy: systematic review of Chinese literature. *BMJ* 2006;333:374. (19 August.)
- 2 Beller EM, Gebbski V, Keech AC. Randomisation in clinical trials. *Med J Aust* 2002;177:565-7.
- 3 Altman DG, Dore CJ. Randomisation and baseline comparisons in clinical trials. *Lancet* 1990;335:149-53.
- 4 Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 1995;273:408-12.

User fees increase China's health challenges

EDITOR—The editorial by Liu does not give adequate consideration to the policy option of health services at zero cost to users.¹ The evidence from China confirms previous evidence that introducing user fees for health care reduces use of health care and thereby adds to human misery.² For elderly people in China, medical cost (user fees) is the main reason for 40% of those not visiting a doctor and 75% of those not gaining admission to hospitals. Another argument against user fees is that it worsens health inequities.³ The human costs of prolonged illness at old age and the increase in long term health costs from delayed treatment are further disadvantages.

Even developed economies of Europe have steadfastly maintained some facet of a policy of zero user fees. In many, the only out of pocket expense is a partial payment for drugs.³ Developing countries, such as Sri Lanka and Cuba, are other examples where the governments have resisted charging user fees for health. This policy, together with women's education and extensive distribution of health services, has enabled good health (at least for infants and mothers) at relatively low cost.⁴

The evidence from China is another example which supports the idea that poorer countries should provide health care at zero cost to the users until proved otherwise. This should apply equally, at least

to the poorer regions of China, and removal of user fees should be a priority, using well established systematic methods.⁵

Saroj A Jayasinghe *locum consultant physician*
Barking, Havering and Redbridge Hospitals NHS
Trust, Romford RM7 0BE
adosaroj@hotmail.co.uk

Competing interests: None declared.

- 1 Liu T. China's health challenges. *BMJ* 2006;333:365. (19 August.)
- 2 Zhang T, Chen Y. Meeting the needs of elderly people in China. *BMJ* 2006;333:363-4. (19 August.)
- 3 Creese A. User fees. *BMJ* 1997;315:202-3.
- 4 Bhutta Z, Nundy S, Abbasi K. Is there hope for South Asia? *BMJ* 2004;328:777-8.
- 5 Gilson L, McIntyre D. Removing user fees for primary care in Africa: the need for careful action. *BMJ* 2005;331:762-5.

Birth ratio in India is also dwindling

EDITOR—Similar to the Chinese government,¹ the government of India also adopted a policy of two children per couple, which was later modified to the one child norm, to keep a check on the increasing population.

But until now the population is nowhere near being controlled, and the ratio seems to be increasingly imbalanced as the time passes. In many villages, marriages occur much before the legal age of consent. Five to six children are born to each couple. The general feeling is more kids, more hands, and more income. The scenario is no different in cities, where the total number of children born to a couple may be less, but at the cost of female feticide (in spite of a law against prenatal determination).

This scenario will not change for many years. The only hope seems to be women's education, because it is rightly said that by educating a man you educate a person but by educating a woman you educate an entire family.

Charu Chanana *registrar*
All India Institute of Medical Sciences, New Delhi
110029, India
charuchanana@rediffmail.com

Competing interests: None declared.

- 1 Ding QJ, Hesketh T. Family size, fertility preferences, and sex ratio in China in the era of the one child family policy: results from national family planning and reproductive health survey. *BMJ* 2006;333:371-3. (19 August.)

Rose tinted spectacles do not help

EDITOR—Potts overplays the benefits of China's one child policy and greatly underplays its drawbacks (late abortion, infanticide, and gross interference of an unelected government in the affairs of private citizens).¹ At one point he said that a falling birth rate offers a demographic dividend, as the economically productive proportion of the population grows more rapidly than the general population. As a doctor and an epidemiological simpleton, I would like this point to have been explained more thoroughly, especially as it seems to conflict with his later point that when birth rates fall rapidly, during their most productive years the younger generation needs to support a high ratio of old people.

When Potts came closest to acknowledging the political controversy that surrounds the one child policy, he couldn't resist a bit of Bush bashing. Why is the US government described as the "Bush administration," whereas its Chinese counterpart is called, not the "Hu administration" but simply "China"? At least Bush, unlike Hu, was elected by his citizens. And why is the US public criticism of the one child policy said to come from "a veritable media industry"? What about Chinese public criticism of the policy? Oops, sorry, I forgot.

Potts tells us that people in countries that fail to control their populations will either starve or be dependent on Western humanitarian aid. China, in contrast, "reaches out to help other countries." It was therefore surprising to read, a few pages further on in the same issue, that the UK government is to donate £30m (€44m; \$57m) to China to help its HIV/AIDS programme.²

Finally Potts praises the sacrifices the Chinese people made to implement the one child policy. I wish he had discussed the extent to which their sacrifice was voluntary. I wonder if the concept of "sacrifice" is too sanitised, and whether he had considered alternative words such as "anguish" or "despotism"?

Vaughan P Smith *general practitioner*
Taunton TA2 7SZ
VSmith1951@aol.com

Competing interests: None declared.

- 1 Potts M. China's one child policy. *BMJ* 2006; 333: 361-2. (19 August.)
- 2 In brief. UK donates aid to China. *BMJ* 2006;333: 7564:368. (19 August.)

Doctors' consciences are called on

EDITOR—I read the Editor's choice on doctors' consciences with increasing dismay and distress.¹ Considering that China has "arguably a cupboard full of skeletons" pays little regard to the continual abuses of human rights throughout China and especially in Tibet. Although a member of the United Nations, China has yet to ratify the covenant on civil and political rights (among others). This is not something that should be disregarded.

China may be trying to improve health care and openness, but that is of little help to the marginalised races and political opponents. Can forced abortions and sterilisations ever be justified?

Engaging with our Chinese colleagues and establishing links is a positive way forward for humanity, but we cannot blinker ourselves or consider it impolite to ask difficult questions.

Surely as doctors we should be defending all patients, the weak and the sick.

Kim D Miller *general practitioner*
Alness/Invergordon Medical Group, Rossshire
IV17 0UN
drkimmiller@hotmail.com

Competing interests: None declared.

- 1 Kale R. Changing China [Editor's choice]. *BMJ* 2006;333: 0-4777. (19 August.)

Spironolactone and risk of upper gastrointestinal events

Spironolactone may not be guilty

EDITOR—Verhamme et al present their data and case for warning patients on spironolactone about an increased risk of upper gastrointestinal bleeding.¹ They use an electronic database and conditional logistic regression analysis with adjustments. We do not believe their conclusion, however, and will not alter our clinical practice.

From a population of 300 000, they identified 523 cases with upper gastrointestinal events, of which only 13 were currently taking spironolactone. Of these 13, nearly 70% were also on agents known to cause ulcers. Only 30 of the control population were currently taking spironolactone, but these controls were less likely to be taking ulcer causing drugs. Similarly, the control patients overall had significantly less comorbidity than the case patients.

Despite the efforts of regression analysis, basing a conclusion on such a small number of patients is meaningless. So will we warn patients about a risk of bleeding—no, not due to spironolactone. But common sense says they are at increased risk of gastrointestinal events if they have multiple medical conditions and are taking ulcer causing agents.

Eugene Campbell *research fellow*
eugene.campbell@nottingham.ac.uk
Anthony Shonde *research fellow*
Stephen Foley *research fellow*
University of Nottingham, Nottingham NG7 2UH

Competing interests: None declared.

- 1 Verhamme K, Mosis G, Dieleman J, Stricker B, Sturkenboom M. Spironolactone and risk of upper gastrointestinal events: population based case-control study. *BMJ* 2006; 333:330.

Author's reply

EDITOR—With interest, we have read the rapid responses to our article and would like to reply to some of the questions and concerns that have been raised.

Campbell et al questioned the validity of our study and the conclusions we drew as the number of exposed cases was low and as cases had more comorbidity than controls. Low numbers, however, are a precision issue rather than an argument against validity. The fact that cases had more morbidity than controls is not surprising but can be adjusted for in the analysis. We disagree that conclusions based on small numbers are meaningless. Our study is not the first to report on this association; there have been case reports, a clinical trial, and recently a retrospective cohort study on the side effects of spironolactone in 762 patients has been published where it was shown that gastritis was reported in 2% of all patients.^{1 2}

Katia M Verhamme *senior researcher*
k.verhamme@erasmusmc.nl

Jeanne P Dieleman *senior researcher*
Bruno H Ch Stricker *professor of*
pharmacoepidemiology

Miriam C J M Sturkenboom *associate professor*
Department of Medical Informatics, Erasmus MC,
3000 DR Rotterdam, Netherlands,

Competing interests: None declared

- Weil J, Langman MJ, Wainwright P, Lawson DH, Rawlins M, Logan RF, et al. Peptic ulcer bleeding: accessory risk factors and interaction with non-steroidal anti-inflammatory drugs. *Gut* 2000;46:27-31.
- Williams E, Katholi R, Karambelas M. Use and side-effect profile of spironolactone in a private cardiologist's practice. *Clin Cardiol* 2006;29:149-53.

Association is unproved

EDITOR—The study reported by Verhamme et al has several noteworthy limitations.¹ Firstly, the mortality benefit conferred by spironolactone therapy in heart failure has been shown only among patients with severe disease² and hence, as the authors acknowledged, such patients are far more likely to be prescribed this therapy than patients with, for example, relatively asymptomatic echocardiographic evidence of impaired left ventricular function. Given the fact that the authors neither stated the criteria by which they defined heart failure, an important consideration in any primary care based study,³ nor reliably adjusted for its severity in the risk model, it is plausible that the observed higher risk of upper gastrointestinal bleeding among spironolactone users is explicable by a preponderance of severe heart failure in this group.

Secondly, the study made no attempt to examine or adjust for the confounding effect of *Helicobacter pylori* infection, a crucial player in the pathogenesis of peptic ulcer disease,⁴ leaving its findings open to the criticism that they may simply be due to a higher prevalence of *H. Pylori* among patients with gastroduodenal ulcers.

Thirdly, notwithstanding the exclusion from the study of people with oesophageal varices and alcoholism, it is still possible that it included some patients with chronic liver disease. In view of the well established increased risk of peptic ulcer disease among patients with cirrhosis,⁵ many of whom are regular consumers of spironolactone, it is conceivable that the observed association between this drug and upper gastrointestinal bleeding may well reflect a higher prevalence of cirrhosis among patients who bled while taking the drug, a possibility that was not taken into account in the risk adjustment.

Given the above shortcomings, the authors' conclusion, that spironolactone therapy increases the risk of gastroduodenal ulcers and upper gastrointestinal bleeding, remains unproved, and their recommendation to caution patients and doctors about this potential complication should therefore await verification in a more robust risk adjusted analysis.

Muhammad F Dawwas *clinical research fellow*
Clinical Effectiveness Unit, Royal College of Surgeons of England, London WC2A 3PE
dawwas@rcseng.ac.uk

Competing interests: None declared.

- Verhamme K, Mosis G, Dieleman J, Stricker B, Sturkenboom M. Spironolactone and risk of upper gastrointestinal events: population based case-control study. *BMJ* 2006;333:330-3. (12 August.)
- Pitt B, Zannad F, Remme WJ, Cody R, Castaigne A, Perez A, et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized aldosterone evaluation study investigators. *N Engl J Med* 1999;341:709-17.
- Sparrow N, Adlam D, Cowley A, Hampton JR. The diagnosis of heart failure in general practice: implications for the UK National Service Framework. *Eur J Heart Fail* 2003;5:349-54.

- Suerbaum S, Michetti P. *Helicobacter pylori* infection. *N Engl J Med* 2002;347:1175-86.
- Saad RJ, Chey WD. Peptic ulcer disease in patients with chronic liver disease: looking beyond bugs and drugs. *Gastrointest Endosc* 2005;62:357-9.

Causality remains unconvincing

EDITOR—I can draw only one certain conclusion from the study reported by Verhamme et al¹: it is unwise to ascribe causality to an association demonstrated by a case-control study. The authors have shown that people who get gastrointestinal bleeds are four to six times more likely to be taking spironolactone than those who don't get such bleeds. Does this mean that spironolactone causes gastrointestinal bleeding? If it does, then antacids (odds ratio 3.52) must also cause gastrointestinal bleeding. And proton pump inhibitors and H₂ antagonists (odds ratio 2.83) must be more likely to cause gastrointestinal bleeding than aspirin or anticoagulants (odds ratio 2.16 for both).

There are two possible explanations for the authors' findings: spironolactone causes gastrointestinal bleeding, and people who are given spironolactone are inherently more likely to get gastrointestinal bleeding.

The authors have tried to compensate for the known risk factors for gastrointestinal bleeding, but have they identified them all? In my own practice, spironolactone is used rarely, exclusively in heart failure, and particularly in difficult heart failure patients. Was there a difference in potassium values or creatinine values between cases and controls? Doesn't renal function influence the risk of gastrointestinal bleeding and affect the serum levels of a host of drugs?

Verhamme et al do not openly argue that there is a causal relation, but they imply it the last sentence of their paper. If I were to tell a patient "Did you know that people taking spironolactone are more likely to get bleeding ulcers?" they would assume that they could lower the risk by stopping the drug. They shouldn't. Nor should we.

There is, after all, an association between height and baldness (confounding factors, age and sex) but shaving your head does not make you taller. In fact, it makes you slightly shorter (personal observation).

Gruffydd P Jones *general practitioner*
Meddygfa Waunfawr, Caernarfon, Gwynedd LL55 4YY
gruffydd_jones@yahoo.se

Competing interests: None declared.

- Verhamme K, Mosis G, Dieleman J, Stricker B, Sturkenboom M. Spironolactone and risk of upper gastrointestinal events: population based case-control study. *BMJ* 2006;333:330-3.

Avoid delaying surgery in patients with severe ulcerative colitis

EDITOR—Collins and Rhodes, in their review of medical treatments for ulcerative colitis, cover surgical management options only very briefly, despite their acknowledgment that the overall colectomy rate is still about 25%.¹ Their comment that colec-

tomy may be needed if medical treatment does not produce a substantial response within seven to 10 days cannot go unchallenged.

Fifteen per cent of patients with ulcerative colitis will have a severe attack, requiring admission to hospital, at some time during their illness. These patients are traditionally treated with intravenous corticosteroids, with a response rate of 60%; the remaining 40% require colectomy, which essentially cures the patient. Persisting with medical treatment if the response is inadequate is dangerous. If surgery is delayed the patient's general condition may deteriorate and lead to a risk of multiple organ dysfunction syndrome,² which may adversely affect the outcome.

Scoring systems for ulcerative colitis are useful in stratifying the disease, with the aim of predicting the failure of medical therapy and identifying likely surgical candidates early. One of the most widely used is that of Travis et al,³ who showed that it could be predicted on day 3 that 85% of patients with more than eight stools on that day, or a stool frequency between three and eight together with a C-reactive protein level >45 mg/l, would require colectomy. Although it has often not been possible to demonstrate a clear significant relation between postoperative mortality and the time between admission and operation,⁴ it seems nonsensical to persist with medical treatment, which is unlikely to work after the third day, for up to 10 days, when surgery may offer cure and to delay it may be to the detriment of the patient. In cases of severe ulcerative colitis, most surgeons would advocate joint care by a physician and a surgeon in order to avoid any unnecessary delay in surgery.

Neil J Smart *specialist registrar general surgery*
North Bristol NHS Trust
drneilsmart@hotmail.com

Competing interests: None declared.

- Collins P, Rhodes J. Ulcerative colitis: diagnosis and management. *BMJ* 2006;333:340-3. (12 August.)
- Caprilli R, Latella G, Vernia P, Frieri G. Multiple organ dysfunction in ulcerative colitis. *Am J Gastroenterol* 2000;95:1258-62.
- Travis SP, Farrant JM, Ricketts C, Nolan DJ, Mortensen NM, Kettlewell MG, et al. Predicting outcome in severe ulcerative colitis. *Gut* 1996;38:905-10.
- Ritchie JK, Ritchie SM, McIntyre PB, Marks CG. Management of severe acute colitis in district hospitals. *J R Soc Med* 1984;77:465-71.

bmj.com

We select the letters for these pages from the rapid responses posted on bmj.com favouring those received within five days of publication of the article to which they refer.

Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.