Retinal is formed from apo-carotenoids in *Nostoc* sp. PCC7120: *in vitro* characterization of an apo-carotenoid oxygenase

Daniel SCHERZINGER*, Sandra RUCH*, Daniel P. KLOER†, Annegret WILDE‡ and Salim AL-BABILI*¹

*Albert-Ludwigs University of Freiburg, Institute of Biology II, Cell Biology, Schanzlestrasse 1, D-79104 Freiburg, Federal Republic of Germany, †Albert-Ludwigs University of Freiburg, Institute of Organic Chemistry and Biochemistry, Albertstrasse 1, D-79104 Freiburg, Federal Republic of Germany, and ‡Humboldt University Berlin, Institute of Biology, Plant Biochemistry, Chausseestrasse 117, D-10115 Berlin, Federal Republic of Germany

The sensory rhodopsin from *Anabaena* (*Nostoc*) sp. PCC7120 is the first cyanobacterial retinylidene protein identified. Here, we report on NosACO (*Nostoc* apo-carotenoid oxygenase), encoded by the ORF (open reading frame) *all4284*, as the candidate responsible for the formation of the required chromophore, retinal. In contrast with the enzymes from animals, NosACO converts β -apo-carotenals instead of β -carotene into retinal *in vitro*. The identity of the enzymatic products was proven by HPLC and gas chromatography–MS. NosACO exhibits a wide substrate specificity with respect to chain lengths and functional end-groups, converting β -apo-carotenals, (3*R*)-3-hydroxy- β -apo-carotenals and the corresponding alcohols into retinal and (3*R*)-3-hydroxyretinal respectively. However, kinetic analyses revealed very divergent K_m and V_{max} values. On the basis of the crystal

structure of SynACO (*Synechocystis* sp. PCC6803 apo-carotenoid oxygenase), a related enzyme showing similar enzymatic activity, we designed a homology model of the native NosACO. The deduced structure explains the absence of β -carotene-cleavage activity and indicates that NosACO is a monotopic membrane protein. Accordingly, NosACO could be readily reconstituted into liposomes. To localize SynACO *in vivo*, a *Synechocystis* knock-out strain was generated expressing SynACO as the sole carotenoid oxygenase. Western-blot analyses showed that the main portion of SynACO occurred in a membrane-bound form.

Key words: carotenoid cleavage, *Nostoc* and *Synechocystis* apo-carotenoid oxygenases (NosACO and SynACO), cyanobacteria, opsin, retinal.

INTRODUCTION

Retinal represents the chromophore of the widespread family of retinylidene proteins, commonly called rhodopsins. These light-responsive seven-helix transmembrane proteins are common in archaea, algae and animals. In addition, they have recently been discovered in eubacteria and fungi. The basic reaction catalysed by rhodopsins is the light-dependent *cis*–*trans* isomerization of retinal, attached in a Schiff-base linkage to a lysine residue in the seventh helix, across one of the double bonds in the polyene chain (for reviews, see [1,2]).

Cyanobacteria employ different receptors for sensing light (for a review, see [3]). It was shown that they contain phytochromes, cryptochromes and, as recently reported, rhodopsin [4]. There have been earlier indications for the occurrence of retinylidene receptors in cyanobacteria such as *Calothrix* [5] and *Leptolyngbya* [6], and direct evidence was delivered by Jung et al. [4], who identified a sensory rhodopsin from *Anabaena* (*Nostoc*) sp. PCC7120 (hereafter *Nostoc* sp. PCC7120). ASR (*Anabaena* sensory rhodopsin) has been a subject of comprehensive analyses, including its photochemical characteristics and structure at atomic level. ASR shows unique characteristics regarding its interacting transducer, its photocycle and the isomeric state of the bound chromophore on light and dark adaptation. In addition, it was shown that ASR exhibits light-induced reversible interconversion between 13-*cis*- and all-*trans*-retinal, leading to a wavelength-dependent isomer ratio [4,7–9]. Therefore ASR may act as a sensor for light-regulated colour-sensitive processes such as chromatic adaptation [10].

The chromophore of rhodopsins, retinal, is a C₂₀-isoprenoid that is usually synthesized as a cleavage product of carotenoids. In animals, retinal is formed through symmetrical oxidative cleavage of β -carotene (C₄₀) at the central C¹⁵=C^{15'} double bond. The corresponding enzyme 15,15'-BCO (β - β -carotene-15,15'-oxygenase), also named BCO I, is a member of the widespread carotenoid oxygenase family and has been identified from the fruitfly *Drosophila melanogaster* [11], chicken [12] and mammals [13,14]. Biochemical studies of human BCO I, expressed in insect cells and purified to homogeneity, suggested that at least one unsubstituted β -ionone ring half-site was compulsory for effective cleavage of the carotenoid substrate [15]. It has also been shown that the 15,15' BCO I acts as a mono-oxygenase [16]. However, recent investigations of AtCCD1 [*Arabidopsis thaliana* (thale cress) carotenoid cleavage dioxygenase 1] suggested a dioxygenase mechanism [17].

The identification of BCO I has become possible through its sequence homology with VP14 (viviparous14) from *Zea mays* (maize), the first carotenoid cleavage enzyme characterized. VP14 represents a non-haem iron oxygenase that catalyses the oxidative cleavage of 9-*cis*-violaxanthin and 9'-*cis*-neoxanthin, leading to xanthoxin, the precursor of the phytohormone ABA (abscisic acid) [18]. This cleavage reaction seems to be key regulatory in the biosynthesis of ABA [19–22]. Database information for sequenced genomes reveals the occurrence of homologous enzymes in all taxa (for reviews, see [23,24]), indicating that VP14 is a member of a widespread oxygenase family involved in different physiological processes. For instance, the genome of *A. thaliana* encodes nine putative carotenoid oxygenases [25]

Abbreviations used: ABA, abscisic acid; ASR, *Anabaena* sensory rhodopsin; 15,15'-BCO (also BCO I), β - β -carotene-15,15'-oxygenase; CCD1, carotenoid cleavage dioxygenase 1; DTT, dithiothreitol; EI, electron impact; GC-MS, gas chromatography–MS; NIST, National Institute of Standards and Technology (Gaithersburg, MD, U.S.A.); NosACO, *Nostoc* apo-carotenoid oxygenase; ORF, open reading frame; RPE65, retinal pigment epithelium 65; SynACO, *Synechocystis* apo-carotenoid oxygenase; VP14, viviparous14.

¹ To whom correspondence should be addressed (email salim.albabili@biologie.uni-freiburg.de).

catalysing different cleavage reactions [24]. The cloning of VP14 also led to the identification of carotenoid cleavage enzymes responsible for the synthesis of well known apo-carotenoid pigments, such as bixin in *Bixa orellana* (annatto or lipstick plant) [26] and saffron in *Crocus sativus* (saffron crocus) [27]. It was also shown that the VP14 homologue, CCD1, catalyses the cleavage of several carotenoids leading to the formation of C₁₃ volatile compounds like β -ionone in *A. thaliana* [28], *Lycopersicon esculentum* (tomato) [29] and *Petunia* (petunia) flowers [30]. Recent studies revealed that members of the carotenoid oxygenase family are involved in plant development mediating the synthesis of a yet-unknown carotenoid-derived compound that regulates apical dominance [31–34].

Recently, we have described SynACO, an apo-carotenoid cleavage oxygenase (ACO) from *Synechocystis* sp. PCC6803 (formerly named Diox1), as the first retinal-forming enzyme of eubacterial origin. In contrast with the related BCO I from animals, SynACO does not utilize β -carotene, but uses several carotenoid-cleavage products, named apo-carotenoids (e.g. 3-hydroxy-apo-carotenoids) to form retinal and retinal-like compounds [35]. As the first member of the widespread carotenoid oxygenase family, the crystal structure of SynACO was recently solved at 2.4 Å (1 Å = 0.1 nm) resolution, revealing the reaction-centre geometry and establishing a solid base for modelling of other family members [36].

The occurrence of a functional sensory rhodopsin in *Nostoc* sp. PCC7120 implies the ability of this cyanobacterium to synthesize the chromophore retinal. The genome of this filamentous cyanobacterium encodes three members of the carotenoid oxygenase family [ORFs (open reading frames) *all4284*, *all4895* and *all1106*] that may catalyse retinal synthesis. Here we report that the ORF *all4284*-encoded polypeptide, here named NosACO, is an apo-carotenoid-cleaving retinal-forming enzyme showing an enzymatic activity similar to that of SynACO from *Synechocystis* sp. PCC6803. In addition, reconstitution of NosACO and localization of SynACO indicated that both polypeptides are membrane-bound. The obtained experimental data were corroborated by a model of NosACO deduced from the crystal structure of SynACO. Our data indicate that the cleavage of apo-carotenoids to form retinal and retinal-like compounds is probably a widespread reaction in cyanobacteria.

EXPERIMENTAL

Cultures of cyanobacteria

Nostoc sp. PCC7120 cells were cultured in liquid BG11₀ medium [37] supplemented with 10 mM NaHCO₃. The *Synechocystis* sp. PCC6803 *SynDiox2* knock-out strain was grown in BG11 medium [37] containing 15 µg/ml chloramphenicol. Cultures were grown in Erlenmeyer flasks under constant illumination of 15 µmol of photons · s⁻¹ · m⁻² at 30 °C.

Cloning of *Nostoc* carotenoid oxygenases

The gene *all4284* coding for NosACO (accession number: NP_488324) was amplified using the primer pair NosACO-for: 5'-ATGCAAAGTTACAAATATCAACA-3' and NosACO-back: 5'-TTATACAAATACCTGGGAGTAA-3'. The ORFs *all4895* and *all1106*, encoding the other two carotenoid oxygenases from *Nostoc*, were amplified using the primer pairs ND2F/ND2B (5'-ATGCAGATAGTTGATAAAGGTCA-3'; 5'-TTATCTCTTCTCTGATTTCCATGT-3') and ND3F/ND3B (5'-ATGGTAAAAGATTCACCTCACTTTC-3'; 5'-TTAAATTCTCTTAAATTC-3'), respectively. The PCR reactions were carried out using approx. 100 ng of genomic DNA, 100 ng of each primer,

200 µM dNTPs and 1 µl Advantage[®] cDNA Polymerase Mix (BD Biosciences) in the buffer provided, as follows: 2 min initial denaturation at 94 °C followed by 32 cycles (30 s at 94 °C, 30 s at 56 °C, 2 min at 68 °C) and 10 min final polymerization at 68 °C. The obtained PCR products were purified using GFX[™] PCR DNA and Gel Band Purification Kit (Amersham Biosciences) and cloned into the pCR2.1[®]-TOPO[®] and pBAD/TOPO[®]-ThioFusion[™] vectors (Invitrogen). The nature of the products was verified by sequencing. The obtained pBAD/TOPO[®]-ThioFusion[™] vectors were used for the expression in carotenoid-accumulating *Escherichia coli* cells. The vectors pNosACO-CR2.1, pND2-CR2.1 and pND3-CR2.1 which carry *NosACO*, *NosDiox2* and *NosDiox3* were used for further cloning.

Protein expression and purification

NosACO was expressed as a glutathione-S-transferase (GST) fusion protein. For this purpose, *NosACO* was excised from pNosACO-CR2.1 using EcoRI, and ligated into EcoRI/ calf intestinal phosphatase-treated pGEX-5X-1 (Amersham Biosciences) to yield the vector pGEX-5X-NosACO. Subsequently, BL21 *E. coli* cells were transformed with pGEX-5X-NosACO, grown at 37 °C in 2 × YT medium (16 g/l tryptone, 10 g/l yeast extract and 5 g/l NaCl) and induced at an attenuation (*D*₆₀₀) of 0.5 with 0.2 mM isopropyl β -D-thiogalactoside. After incubation for additional 4 h at 28 °C, cells were harvested by centrifugation, resuspended in PBS and lysed in a French press. The fusion protein was then purified using glutathione-Sepharose 4B (Amersham Biosciences), and NosACO was released by overnight treatment with the protease Factor Xa in PBS containing 0.1 % (v/v) Triton X-100 at room temperature, according to the instructions of the manufacturer (Amersham Biosciences). For the reconstitution assays, NosACO was released in PBS containing 1 % (w/v) octyl β -glucoside. Purification steps and protein expression were analysed by SDS/PAGE.

Analyses of the membrane association

All steps were performed in 100 mM Bistris/HCl buffer, pH 6.8, containing 1 mM DTT (dithiothreitol). For preparation of liposomes, 1,2-diacyl-*sn*-glycero-3-phosphocholine from soya bean (*Glycine max*) (Sigma) was solubilized in chloroform/methanol (2:1, v/v); β -apo-8'-carotenol was then added to a final concentration of 30 µM. After drying, the buffer was added to obtain a final lipid concentration of 1 mg/ml. After incubation for 1 h on ice, liposomes were obtained by sonication on ice. A 720 µl portion of purified NosACO (about 1.2 µg/µl) in PBS containing 1 % (w/v) octyl β -glucoside was diluted (1:5) with the liposomes and incubated on ice for 10 min. Liposomes were pelleted at 100 000 g for 30 min, washed with 3.6 ml buffer, pelleted again, then resuspended in the same volume of buffer.

For the salt treatment, 1.8 ml of liposome suspension, containing NosACO and prepared as described above, was adjusted to a final concentration of 0.2 M KCl and incubated on ice for 10 min. Liposomes were then harvested by centrifugation at 100 000 g for 30 min, washed in 1.8 ml of buffer, pelleted again, then resuspended in 1.8 ml of buffer. Samples corresponding to equal portions of each fraction were subjected to SDS/PAGE and analysed by Western blotting using an ECL[™] Western Blotting Analysis System (Amersham Biosciences). NosACO was detected using mice polyclonal anti-SynACO antibodies raised against overexpressed and purified enzyme [36].

Synechocystis *SynDiox2* knock-out cultures were harvested, washed, then resuspended in PBS. The cells were disrupted by four passages through a French pressure cell at 18 000 lbf/in² (1 lbf/in² = 6.9 kPa). Cell debris was removed by repeated

centrifugation for 10 min at 4000 *g*. The supernatant obtained was then separated into a membrane and a cytoplasmic fraction by centrifugation for 4 h at 130 000 *g*. Membranes were washed with PBS, pelleted by centrifugation at 130 000 *g* for 2 h, then resuspended in PBS. Samples containing equal amounts of protein from each fraction were analysed by SDS/PAGE and Western blotting using the ECLTM Western Blotting Analysis System and mouse polyclonal anti-SynACO antibodies.

Enzyme assays

Substrates were purified as described in [35]. Synthetic apo-carotenals were kindly provided by BASF. The corresponding alcohols were obtained by reduction of the aldehydes with NaBH₄ in ethanol. Enzyme assays were performed as described in [26], with some modifications. Substrates were mixed with an ethanolic solution of octyl β -glucoside, dried using a vacuum centrifuge, and then resuspended in water. The final concentration of the apo-carotenoid substrates was 40 μ M; the concentration of octyl β -glucoside was 1 % (w/v). The incubation buffer contained 1 mM DTT (dithiothreitol) and 1 mg/ml catalase (Sigma) in 200 μ l of 100 mM Bistris/HCl, pH 6.8. Purified NosACO was then added to obtain a concentration of 100 ng/ μ l. Assays were incubated at 27 °C for 2 h. For GC-MS identification of the cleavage products the assay volume was 2 ml with incubation for 2.5 h. The assays were stopped by adding 1 vol. of acetone and extracted using petroleum benzene/diethyl ether (1:4, v/v). The analysis was performed as described below.

Analytical methods

Substrates and products were quantified spectrophotometrically at their individual $\lambda_{\max}$ values using absorption coefficients calculated from the $A^{1\%}_{1\text{cm}}$ as given by Barua and Olson [38] or as determined with the synthetic compounds (BASF). Protein concentration was determined using the Bio-Rad Protein Assay Kit. For HPLC, a Waters system equipped with a photodiode array detector (model 996) was used. The separation was performed using a C₃₀-reversed phase column (YMC Europe GmbH, Dinslaken, Germany) with the following solvent systems: B, methanol/*t*-butyl methyl ether/water (30:1:10, by vol.); A, methanol/*t*-butyl methyl ether (1:1, v/v). The column was developed at a flow rate of 1 ml/min with a gradient from 100 % B to 43 % B within 45 min, then to 0 % B within 1 min, maintaining the final conditions for another 14 min at a flow rate of 2 ml/min.

GC-MS analyses

To remove detergents present in the extracts, products were pre-purified by TLC using RP-18 F_{254s} plates (Merck). All plates were washed with chloroform prior to use. The plates were developed in methanol/water (100:1, v/v). Products were then scraped off and eluted with chloroform, evaporated, then redissolved in acetone for analyses. GC-MS analyses were performed as described in [35] using a Finnigan Trace DSQ mass spectrometer coupled to a Trace GC gas chromatograph. Identification of compounds was done by chromatographic comparison with the authentic references and by comparing the mass spectra with the NIST (National Institute of Standards and Technology) Mass Spectral Search Program Version 2.0.

Kinetic analyses

Measurements were carried out photometrically at 20 °C using a Shimadzu UV-2501PC spectrophotometer. The assays contained 100 mM Bistris/HCl, pH 6.8, 1 % (w/v) octyl β -glucoside, 1 mM DTT, 200 ng/ml catalase and purified NosACO at a concentration

of 30 ng/ μ l in a final volume of 700 μ l. The concentrations of the apo-carotenoid substrates used ranged from 5 to 80 μ M. The formation of the C₂₀ cleavage products was determined during an incubation of 5–15 min. Time intervals with maximum velocity were used for further calculations. The enzyme kinetics were calculated using the solver function of MS-Excel. This function fits the data to the Michaelis–Menten equation using non-linear regression analysis.

Homology modelling

A homology model of NosACO was built using the published coordinates of SynACO (PDB code 2BIX) as template. Model building and energy minimization was performed using SWISS-MODEL [39] in first approach mode. Solvent-accessible surfaces were generated with MSMS [40] software and analysed with the PISA [41] program.

Generation of the *Synechocystis* *SynDiox2* knock-out strain

A 557 bp fragment (*delta5*), representing a stretch of the 5'-non-coding region and a part of *SynDiox2*, and a 448 bp fragment (*delta3*), corresponding to the 3'-non-coding region and a part of *SynDiox2*, were amplified from 25 ng of *Synechocystis* sp. PCC6803 genomic DNA using Advantage[®] cDNA Polymerase Mix (BD Biosciences) in the buffer provided. The PCR reactions were performed using the primer pair I/II and III/IV for the amplification of *delta5* and *delta3*, respectively (I: 5'-GGGC-CGCTACTCGAGCCAGTCTG-3'; II: 5'-TCCTCTGCGTAAGC-TTGGGTTTGC-3'; III: 5'-ACCGGCGGAATTCCTTGTTTC-3'; IV: 5'-TGCTTAATGAGCTCTCCAGTTTCA-3'). Both fragments were then cloned sequentially into pKSCm2, a pBluescript KS with a *cat* (chloramphenicol transferase) gene cassette integrated into the EcoRV site, after XhoI/HindIII (for *delta5*) and EcoRI/XbaI (for *delta3*) digestion. The correctness of the resulting suicide vector pDiox2Knock was confirmed by DNA sequencing. pDiox2Knock, bearing a chloramphenicol resistance cassette between *delta5* and *delta3* in antisense orientation, was then transformed into wild-type cells of *Synechocystis* sp. PCC6803 as described by Ermakova et al. [42]. The absence of the wild-type allele in fully segregated mutants was verified by PCR using the primers Co1: 5'-AGTTTGTGTTGACCCAGACCGATGTA-3' and Co2: 5'-ATAGTTGACCATCCGTGGAGGTTT-3'.

RESULTS

β -Carotene is not cleaved by any of the three *Nostoc* carotenoid oxygenases

To determine the enzyme catalysing formation of retinal, the three *Nostoc* ORFs *all4284* (here *NosACO*), *all4895* (*NosDiox2*) and *all1106* (*NosDiox3*) encoding putative carotenoid oxygenases were cloned and expressed in β -carotene-accumulating *E. coli* cells. However, none of the expressed enzymes was capable of converting β -carotene, as determined by the absence of cleavage products on HPLC analyses (results not shown).

NosACO converts apo-carotenals into retinal

Since no cleavage activity was observed in carotenoid-accumulating *E. coli* cells, *in vitro* assays were performed using total lysate of cells expressing NosACO, NosDiox2 and NosDiox3 respectively. Subsequent analyses revealed that none of the expressed oxygenases is capable of β -carotene cleavage. Therefore apo-carotenals with different chain lengths were applied. Subsequent HPLC analyses showed that NosACO converted apocarotenoids with different chain lengths into a compound

Figure 1 HPLC analyses of the *in vitro* assay products obtained from β -apo-8'-carotenal (A), β -apo-10'-carotenal (B) and (3R)-3-hydroxy- β -apo-8'-carotenal (C) and the corresponding UV-visible spectra (I–III)

After 2.5 h incubation with purified NosACO (ACO), a striking cleavage activity was observed, indicated by colour changes compared with the controls (con). The reactions led to the formation of the products apo-8',15'-apo-carotenal (I) and retinal (II) from β -apo-8'-carotenal (A), retinal (II) from β -apo-10'-carotenal (B) and (3R)-3-hydroxyretinal (III) and apo-8',15'-apo-carotenal (I) from (3R)-3-hydroxy- β -apo-8'-carotenal (C). The structure of the products is given above the corresponding UV-visible spectra (see also Figures 2 and 3).

resembling retinal (C_{20}), whereas the products obtained from NosDiox2 and NosDiox3 were different in chain length depending on the substrates used (results not shown). This indicated that NosACO is the candidate for retinal formation. For a more comprehensive investigation of its enzymatic activity, NosACO was expressed as a GST fusion, purified using glutathione-Sepharose and released by the protease Factor Xa (see Supplementary Figure 1 at <http://www.BiochemJ.org/bj/398/bj3980361add.htm>). Here again, *in vitro* assays using the purified enzyme and major carotenoid compounds of *Nostoc*, e.g. β -carotene, echinenone and myxoxanthophyll, did not show any cleavage activity, whereas apo-carotenals were converted readily. As can be seen in Figure 1,

Figure 2 GC-MS analyses of the *in vitro* assay products

The products obtained from *in vitro* assays (Figure 1) were subjected to GC-MS analyses. The EI mass spectrum of product I was identical with that obtained with synthetic apo-8',15'-apo-carotenal ($C_{10}O_2H_{11}$) and showed the expected molecular mass of m/z 164. The EI mass spectrum of product II showed identity with the reference spectrum of retinal and the expected molecular ion of m/z 284. The EI mass spectrum of product III showed identity with the reference spectrum and the expected molecular ion of m/z 300 as well as the typical fragment of 162 m/z , corresponding to a mass of 300–138, indicative of the loss of the 3-hydroxy- β -ionone ring as described for zeaxanthin [55].

a striking activity was observed with β -apo-8'-carotenal which was converted by NosACO into a compound that resembled retinal in its UV-visible spectrum and its chromatographic behaviour on HPLC. To prove its identity, GC-MS analyses were carried out using retinal as a standard, which showed an identical retention time. The mass spectrum obtained was identical with published data [35,43] and with spectra in the NIST database, including the presence of the correct molecular ion of m/z 284 (Figure 2). Thus NosACO converted the C_{30} substrate into retinal (C_{20}). The

Substrate		Cleavage	
Chain length	β -Apo-4'-carotenal (C_{35})	n.d.
	β -Apo-8'-carotenal (C_{30})	++	
	β -Apo-10'-carotenal (C_{27})	++	
	β -Apo-12'-carotenal (C_{25})	n.d.	
Substitution of the β -ring	(3 <i>R</i>)-3-Hydroxy- β -apo-8'-carotenal	++
	(3 <i>R</i>)-3-Hydroxy- β -apo-12'-carotenal	+	
	Apo-8'-lycopenal	+	
	Apo-10'-lycopenal	++	
	Apo-12'-lycopenal	n.d.	
Alcohols	β -Apo-8'-carotenol	++
	(3 <i>R</i>)-3-Hydroxy- β -apo-8'-carotenol	++	
	Apo-8'-lycopenol	+

Figure 3 Structure of applied substrates and determined products (I–III)

The degree of cleavage was estimated by the ratio of cleavage product compared with the substrate. ++, Cleavage; +, low cleavage; n.d., not detected. The substrates β -apo-8'-carotenal (C_{30}), its corresponding alcohol and β -apo-10'-carotenal (C_{27}) were cleaved into retinal (C_{20} , **II**). (3*R*)-3-hydroxy- β -apo-8'-carotenal, its corresponding alcohol and (3*R*)-3-hydroxy- β -apo-12'-carotenal (C_{25}) were converted into (3*R*)-3-hydroxy-retinal (**III**). Apo-8',15'-apo-carotenodial (C_{10} , **I**) was produced from the cleavage of the β -apo-8'-carotenoids (C_{30}) as the second product. Substrates were cleaved at the $C^{15}=C^{15'}$ double bond, as indicated. With exception of apo-12'-lycopenals, the enzyme converted all applied apo-lycopenals into the corresponding C_{20} compound acycloretinal (not shown). The incubation with β -apo-4'-carotenal and β -apo-12'-carotenal did not lead to detectable products.

catalytic activity of NosACO should also lead to a second product(s) with a chain length of C_{10} or shorter. Indeed, apo-8',15'-apo-carotenodial (2,6-dimethylocta-2,4,6-trienedial, $C_{10}O_2H_{11}$) was detected and authenticated by HPLC (Figure 1) and GC-MS (Figure 2) analyses as a second product showing the chromatographic characteristics and the EI (electron impact) mass spectrum of the authentic reference compound (kindly provided by BASF). Taken together, the nature of the products identified demonstrates the cleavage of the used apo-carotenoids at the $C^{15}=C^{15'}$ double bond (Figure 3).

Substrate specificity of NosACO

The *Synechocystis* enzyme SynACO was shown to cleave a wide range of different apo-carotenoids [35]. To investigate the end-

group specificity of NosACO, (3*R*)-3-hydroxy- β -apo-8'-carotenol formally derived from zeaxanthin was used *in vitro* as described. As shown by HPLC analysis (Figure 1), this substrate was converted by the enzyme into (3*R*)-3-hydroxyretinal and apo-8',15'-apo-carotenodial. The nature of the C_{20} product obtained was further confirmed by GC-MS analysis (Figure 2), revealing a correct molecular ion of m/z 300 and a mass spectrum identical with that of the authentic reference compound (kindly provided by BASF). Further, we tested β -apo-8'-carotenol and the corresponding alcohols of the above-mentioned substrates. Here again, retinal or the corresponding C_{20} compounds were produced. To determine the chain-length specificity, we performed *in vitro* assays with β -apo-4'-carotenal (C_{35}), β -apo-10'-carotenal (C_{27}), β -apo-12'-carotenal (C_{25}) and (3*R*)-3-hydroxy- β -apo-12'-carotenal. The subsequent analysis showed that NosACO cleaved

Table 1 K_m and V_{max} values of NosACO for different substrates

Each value represents the average $\pm$ S.D. for three independent experiments.

Substrate	K_m (μ M)	V_{max} (nmol/min)
β -Apo-8'-carotenal	16.3 ± 1.6	1.0 ± 0.3
β -Apo-10'-carotenal	17.2 ± 1.8	0.3 ± 0.1
(3R)-3-Hydroxy- β -apo-8'-carotenal	2.1 ± 0.3	0.1 ± 0.04
β -Apo-8'-carotenol	24.5 ± 3.1	3.3 ± 1.7

β -apo-10'-carotenal and (3R)-3-hydroxy- β -apo-12'-carotenal forming retinal and (3R)-3-hydroxyretinal respectively (results not shown). In addition, the enzyme also cleaved apo-10'-lycopenal into acycloretinal (results not shown). In contrast, β -apo-12'-carotenal (C_{25}) and β -apo-4'-carotenal (C_{35}) were not cleaved. The substrates and the corresponding chemical structures are given in Figure 3.

Kinetic analyses

The data given above suggest a wide substrate specificity. However, the cleavage reactions exhibited very divergent initial reaction velocities depending on the substrate employed. For instance, apo-8'-lycopenal was clearly converted at a lower rate than was β -apo-8'-carotenal. Kinetic analyses were performed with the substrates β -apo-8'-carotenal, β -apo-10'-carotenal, β -apo-8'-carotenol and (3R)-3-hydroxy- β -apo-8'-carotenol exhibiting different chain lengths and functional groups. Table 1 gives the K_m and V_{max} values determined in the biphasic detergent-containing *in vitro* incubation system used. The lowest K_m values were obtained for (3R)-3-hydroxy- β -apo-8'-carotenol, followed by β -apo-8'-carotenal, β -apo-10'-carotenal and finally by β -apo-8'-carotenol. However, β -apo-8'-carotenal showed lower V_{max} values when compared with the corresponding alcohol, i.e., NosACO exhibited a higher affinity to aldehydes but converted alcohols more quickly. In addition, there was a significantly higher V_{max} with β -apo-8'-carotenal than with (3R)-3-hydroxy- β -apo-8'-carotenol, indicating that the enzyme converted unsubstituted

substrates more quickly than (3R)-3-hydroxy- β -derivatives. Although the determined K_m values for β -apo-8'-carotenal (C_{30}) and β -apo-10'-carotenal (C_{27}) were similar, the C_{30} substrate was converted much more quickly, as indicated by the V_{max} values.

Structure alignment with SynACO and modelling

The observed enzymatic activity of NosACO resembles that described for SynACO from *Synechocystis* [35]. In addition, both enzymes belong to the same family and share about 53 % identity at the amino acid level. The recently solved crystal structure of SynACO [36] allowed the construction of a homology model of NosACO. In the structure-based sequence alignment used, there are no gaps within secondary-structure elements (see Supplementary Figure 2 at <http://www.BiochemJ.org/bj/398/bj3980361add.htm>). Two small insertions of two residues each occur in NosACO as compared with SynACO. Both are located in a loop between β -strands 18 and 19 and were modelled with good stereochemistry. All other residues of the NosACO sequence were exactly superimposable on to the template. Given the considerable sequence identity between SynACO and NosACO, the resulting homology model is expected to be exact within a 1.5 Å root-mean-square deviation with respect to C^α positions [44]. All residues known to be important for substrate binding and catalysis in SynACO are found at corresponding positions in the NosACO homology model. NosACO consists of a single domain and adopts a β -propeller fold with seven propeller blades (Figure 4). Long loops on the top side of the β -propeller form a dome covering the active site, which is located in the centre of the protein at the propeller axis. The catalytic centre is formed by four histidine residues that are strictly conserved in all carotenoid cleavage oxygenases and co-ordinates a ferrous iron ion. Interestingly, all residues lining the hydrophobic substrate tunnel in SynACO, up to 10 Å away from the catalytic centre are either identical or conservatively replaced by residues with the same steric requirements in NosACO, giving rise to a substrate binding cavity with an almost identical shape (Figure 4). Assuming the same substrate binding geometry as was previously observed in SynACO [36], the homology model clearly predicts

Figure 4 Model of NosACO based on homology with SynACO

(A) Ribbon representation of NosACO. β -Strands and α -helices are coloured orange and green respectively. The eight residues forming the large hydrophobic patch at the top of the enzyme are shown in stick representation and labelled. The axis of the β -propeller runs vertically. (B) Surface representation showing the solvent-accessible surface in grey. The entrance to the substrate-binding tunnel is coloured blue and the hydrophobic patch of 889 Å² yellow. (C) Geometry of the substrate-binding tunnel. Surface representation of NosACO rotated by 90° along the propeller axis and cut through the middle of the enzyme. The substrate-binding tunnel runs horizontally and is shaded blue. The position of the catalytic iron atom is indicated in red. In order for cleavage to occur, the substrate must enter the tunnel far enough to bring the scissile double bond in close proximity to the catalytic iron. A constriction in the tunnel, marked by arrows, prevents entry of bulky substrates. (D) Close-up view of the constriction together with a β -ionone ring for size comparison. The constriction is too small to allow substrates carrying β -ionone rings on both termini such as β -carotene to enter the tunnel.

Figure 5 Western-blot analysis of the reconstitution assay

Total reconstitution assay containing soluble and liposome-bound NosACO (lane 1), isolated and washed liposomes (lane 2), the corresponding supernatant containing unreconstituted enzyme (lane 3), liposomes after treatment with 0.2 M KCl (lane 4) and the corresponding supernatant (lane 5). NosACO was detected using polyclonal anti-SynACO antibodies.

that C₄₀ substrates will not be cleaved by NosACO. On the surface of NosACO, a large non-polar patch consisting of leucine, tryptophan, phenylalanine and isoleucine residues lies at the same position as in SynACO (Figure 4), whereas the remaining surface of the molecule is predominantly polar. It was presumed that the corresponding region in SynACO dips into the membrane in order to provide access to the unpolar substrate. The hydrophobic patch in NosACO is even larger (889 Å² as compared to 800 Å² in SynACO), leading to the prediction that NosACO may also be a monotopic membrane protein.

NosACO and SynACO are membrane proteins

The lipophilic nature of the substrates as well as the predicted model indicate that NosACO is a membrane protein. Therefore we performed reconstitution assays into artificial membranes using the purified enzyme. NosACO could be readily reconstituted as shown in the Western-blot analysis (Figure 5). Additionally the membrane binding was stable upon high-salt treatment. These data coincide with the structural model and indicate that NosACO is a membrane-bound enzyme.

The genome of *Nostoc* encodes three members of the carotenoid oxygenase family which may not be distinguished by polyclonal antibodies. In contrast, *Synechocystis* contains only two members of this family (SynACO and ORF *slr648*, hereafter *SynDiox2*). To localize SynACO *in vivo*, we generated a knock-out strain by interrupting *SynDiox2*. Thus this strain encodes only SynACO and allows, therefore, the localization of SynACO by using polyclonal anti-SynACO antibodies, avoiding cross-reactions. As can be seen in Figure 6, only traces of SynACO could be detected in the cytoplasmic fraction, whereas the main portion of the enzyme occurred in a membrane-bound form.

DISCUSSION

The *Nostoc* sensory rhodopsin was the first of its kind found in cyanobacteria. Its occurrence implies the biosynthetic capacity for the chromophore, retinal. According to the current state of knowledge, retinal is formed either by a member of the widespread carotenoid oxygenase family (for reviews, see [45,46]) or by the very recently described enzyme Blh (bacteriorhodopsin-related-protein-like homologue) from marine proteobacteria [47] and its formerly identified halobacterial homologues [48]. Searching the genome of *Nostoc* for putative carotenoid oxygenases revealed the occurrence of three members of this protein family, namely

Figure 6 Localization of SynACO in a *Synechocystis SynDiox2* knock-out strain

(A) Coomassie Blue-stained SDS/polyacrylamide gel of membrane- (lane 1) and cytoplasmic-fraction proteins (lane 2). (B) The corresponding Western-blot analysis. The localization was performed using 25 µg of protein for each fraction.

ORFs *all4284*, *all4895* and *all1106*, whereas no sequence with significant similarity to *Blh* was found. This led us to suggest that retinal is formed through one of the encoded carotenoid oxygenases. However, *in vivo* and *in vitro* studies showed that none of these three enzymes is capable of cleaving β-carotene to yield retinal (residues not shown), excluding a central cleavage mechanism of β-carotene, as commonly found in animals. On the basis of our recent finding that the *Synechocystis* enzyme SynACO forms retinal and retinal-like compounds from apo-carotenoids and not from β-carotene [35], we assumed that *Nostoc* may utilize similar substrates for retinal formation. Accordingly, we performed *in vitro* enzymatic analyses using apo-carotenoids as substrates. Among the three carotenoid oxygenases, NosACO, encoded by the ORF *all4284*, was the one to form retinal and derivatives from apo-carotenals with different chain lengths, indicating its involvement in retinal supply. Notwithstanding the significance of the sequence homology between SynACO and NosACO (53 % identity), it could not necessarily be expected that both enzymes exhibit the same activity and, thus, that both cyanobacteria utilize apo-carotenals to form retinal. Generally, orthologous carotenoid oxygenases show more pronounced homology. For instance, the CCD1 from *A. thaliana* [28] and *L. esculentum* [29] share about 79 % identity at the amino acid level. The sequence divergence between SynACO and NosACO could be explained by the evolutionary distance between the unicellular *Synechocystis* sp. PCC6803 and the axenic nitrogen-fixing *Nostoc* sp. PCC7120.

To characterize the substrate specificity, *in vitro* studies were performed using the purified enzyme. NosACO cleaved β-apo-carotenals with different chain lengths, (3*R*)-3-hydroxy-β-apo-carotenals as well as the corresponding alcohols. Depending on the nature of the end groups, the cleavage led to the formation of retinal and (3*R*)-3-hydroxyretinal. However, the values

determined for K_m and V_{max} differed markedly (Table 1). Given that low substrate concentrations prevail *in vivo*, one would assume the preferred natural substrate to be a C₃-hydroxylated apo-carotenal with a C₂₇ or C₃₀ chain length. However, the identity of the natural substrate and the reaction leading to its formation are still to be elucidated. Like in *Synechocystis* [35], HPLC analyses of carotenoid extracts from *Nostoc* revealed the occurrence of small amounts of compounds which resembled (3R)-3-hydroxy- β -apo-carotenals in their UV-visible spectra as well as in their elution characteristics (results not shown), but these were too low in abundance to allow an investigation of their molecular structure. Such precursor molecules may be formed through the action of a different enzyme. The two additional members of the carotenoid oxygenase family found in the genome of *Nostoc* are good candidates. Although our preliminary analyses indicate that none of these enzymes cleaves β -carotene, it cannot be excluded that a (3R)-3-hydroxy substrate may be supplied by one of these oxygenases through an initial cleavage of myxoxanthophylls, which are common in *Nostoc* [49]. However, carotenoids may be also cleaved by enzymes such as peroxidases and lipoxygenases, which do not belong to the carotenoid oxygenase family. For instance, Zorn et al. [50] reported on a peroxidase from the strong-scented blewit mushroom (*Lepista irina*), which cleaved β -carotene to form certain flavour compounds. In addition, the genome of *Nostoc* encodes a putative lipoxygenase (ORF *all8020*) with unknown function. Alternatively, apo-carotenoids may arise from photo-destruction processes mediated by reactive oxygen species and increased under light or salt stress. It was reported that the transcript level of the homologous enzyme SynACO from *Synechocystis* was increased by 4-fold within 15 min after exposure to high light [51] and by 21-fold within 30 min after treatment with 684 mM NaCl [52]. Considering the similarity in the catalytic activity and the amino acid sequence, it appears conceivable that NosACO may be subjected to a similar regulation. In this case, NosACO may convert a non-enzymatically originated inhomogeneous population of apo-carotenoids into homogeneous C₂₀ compounds, namely retinal, retinol and its in-ring-hydroxylated derivatives.

The enzymatic activity of NosACO, the substrate specificity for apo-carotenoids and the kinetic characteristics resemble those of SynACO [35]. However, NosACO showed more specificity with respect to the substrate's chain lengths and did not convert β -apo-4'- and β -apo-12'-carotenal, which were both cleaved by SynACO [35].

The crystal structure of the latter enzyme explained and thus supported the experimentally shown absence of β -carotene-cleaving activity [35]. SynACO exhibits a substrate cavity with a bottleneck at the entrance arresting the β -ionone ring, thus placing the substrate into the correct position for retinal formation. In other words, the substrate enters the cavity with its open-chain end (Figure 4), since the β -ionone ring cannot pass. The NosACO model deduced from the structure-based alignment with SynACO showed a very similar geometry of the substrate cavity, explaining the observed absence of β -carotene-cleaving activity.

Consistent with the lipophilic nature of substrates and products, a further prediction drawn from the NosACO model is the attribute of being a monotopic membrane protein. In the model, the surface of NosACO exhibits a large non-polar patch consisting of leucine, tryptophan, phenylalanine and isoleucine residues at the same position as in SynACO. It was presumed that the corresponding region in SynACO dips into the membrane. The reconstitution experiments and localization studies shown here indicate that both enzymes are membrane proteins. In both experiments, only traces of soluble enzymes were detected. Similarly, the xanthoxin-forming enzyme PvNCED [*Phaseolus vulgaris*

(French bean) *nine-cis-epoxy-carotenoid dioxygenase*] was found only in the thylakoid fraction of the leaves [19]. In contrast, it was shown that only about 35 % of VP14 from maize is associated with the thylakoid membrane upon import. In addition, Western-blot analyses of maize chloroplasts revealed a stromal and a membrane-bound VP14 fraction [25]. It has been speculated that the degree of membrane association may be regulated in intact chloroplasts and that the soluble fraction is inactive [25], as reported for the phytoene desaturase from *Narcissus pseudo-narcissus* (daffodil) [53]. The membrane association of VP14 is mediated by an N-terminal amphipathic helix [25]. A more complicated pattern was reported for RPE65 (retinal pigment epithelium 65), an animal member of the carotenoid oxygenase family involved in the visual cycle. RPE65 occurs in a triply palmitoylated, membrane-associated (mRPE65) and in a soluble unpalmitoylated form (sRPE65). It appears that the palmitoylation determines whether RPE65 binds all-*trans*-retinyl ester (mRPE65) or all-*trans*-retinol (sRPE65), and thus, controls the ligand binding selectivity [54]. Taken together, it might be speculated that the soluble traces of SynACO may represent an inactive form.

The model also indicates that the product formed is deposited into membranes through a tunnel and not passed to a cytoplasmic protein. To test this prediction, we conducted *in vitro* assays using artificial membranes containing reconstituted NosACO. The subsequent analyses revealed clear retinal formation (results not shown). However, the amount of the product was low compared with the amounts obtained from the detergent-based assays and, therefore, did not allow a clear conclusion about the destination of retinal.

This work was supported by a grant (no. AL892-1) from the DFG (Deutsche Forschungsgemeinschaft) and by the HarvestPlus programme (www.harvestplus.org). We are indebted to Dr Peter Beyer (Center for Applied Biosciences, University of Freiburg, Freiburg, Germany) for valuable discussions. We thank Dr Hansgeorg Ernst (Fine Chemicals Research, BASF, Ludwigshafen, Germany) for providing the apo-carotenoids and Dr Andreas Seidler (Department of Plant Biochemistry, Faculty of Biology, Ruhr-University Bochum, Bochum, Germany) for his help in constructing the knock-out plasmids.

REFERENCES

- Spudich, J. L., Yang, C. S., Jung, K. H. and Spudich, E. N. (2000) Retinylidene proteins: structures and functions from archaea to humans. *Annu. Rev. Cell Dev. Biol.* **16**, 365–392.
- Zhai, Y., Heijne, W. H. M., Smith, D. W. and Saier, Jr, M. H. (2001) Homologues of archaeal rhodopsins in plants, animals and fungi: structural and functional predications for a putative fungal chaperone protein. *Biochim. Biophys. Acta* **1511**, 206–223.
- Mullineaux, C. W. (2001) How do cyanobacteria sense and respond to light? *Mol. Microbiol.* **41**, 965–971.
- Jung, K. H., Trivedi, V. D. and Spudich, J. L. (2003) Demonstration of a sensory rhodopsin in Eubacteria. *Mol. Microbiol.* **47**, 1513–1522.
- Hoff, W. D., Matthijs, H. C. P., Schubert, H., Crielgaard, W. and Hellingwerf, K. J. (1995) Rhodopsin(s) in eubacteria. *Biophys. Chem.* **56**, 193–199.
- Albertano, P., Barsant, L., Passarelli, V. and Gaultieri, P. (2000) A complex photoreceptive structure in the cyanobacterium *Leptolyngbya* sp. *Micron* **31**, 27–34.
- Sineshchekov, O. A. and Spudich, J. L. (2004) Light-induced intramolecular charge movements in microbial rhodopsins in intact *E. coli* cells. *Photochem. Photobiol. Sci.* **3**, 548–554.
- Sineshchekov, O. A., Trivedi, V. D., Sasaki, J. and Spudich, J. L. (2005) Photochromicity of *Anabaena* sensory rhodopsin, an atypical microbial receptor with *cis*-retinal light adapted form. *J. Biol. Chem.* **280**, 14663–14668.
- Vogele, I., Sineshchekov, O. A., Trivedi, V. D., Sasaki, J., Spudich, J. L. and Lueke, H. (2004) *Anabaena* sensory rhodopsin: a photochromic color sensor at 2.0 Å. *Science* **306**, 1390–1393.
- Grossman, A. R., Bhaya, D. and He, Q. (2001) Tracking the light environment by cyanobacteria and the dynamic nature of light-harvesting. *J. Biol. Chem.* **276**, 11449–11542.
- von Lintig, J. and Vogt, K. (2000) Filling the gap in vitamin A research. *J. Biol. Chem.* **275**, 11915–11920.

- 12 Wyss, A., Wirtz, G., Woggon, W., Brugger, R., Wyss, M., Friedlein, A., Bachmann, H. and Hunziker, W. (2000) Cloning and expression of β , β -carotene 15,15'-dioxygenase. *Biochem. Biophys. Res. Commun.* **271**, 334–336
- 13 Redmond, T. M., Gentileman, S., Duncan, T., Yu, S., Wiggert, B., Ganitt, E. and Cunningham, Jr, F. X. (2001) Expression, and substrate specificity of a mammalian β -carotene 15,15'-dioxygenase. *J. Biol. Chem.* **276**, 6560–6565
- 14 Yan, W., Jang, G. F., Haeseleer, F., Esumi, N., Chang, J., Kerrigan, M., Campochiaro, M., Campochiaro, P., Palczewski, K. and Zack, D. J. (2001) Cloning and characterization of a human β , β -carotene-15,15'-dioxygenase that is highly expressed in the retinal pigment epithelium. *Genomics* **72**, 193–202
- 15 Lindqvist, A. and Andersson, S. (2002) Biochemical properties of purified recombinant human β -carotene 15,15'-monooxygenase. *J. Biol. Chem.* **277**, 23942–23948
- 16 Leuenberger, M. G., Engeloch-Jarret, C. and Woggon, W. (2001) The reaction mechanism of the enzyme catalyzed central cleavage of β -carotene to retinal. *Angew. Chem.* **113**, 2684–2687
- 17 Schmidt, H., Kurtzer, R., Eisenreich, W. and Schwab, W. (2006) The carotenase AtCCD1 from *Arabidopsis thaliana* is a dioxygenase. *J. Biol. Chem.* **281**, 9845–9851
- 18 Schwartz, S. H., Tan, B. C., Gage, D. A., Zeevart, J. A. and McCarty, D. R. (1997) Specific oxidative cleavage of carotenoids by VP14 of maize. *Science* **276**, 1872–1874
- 19 Qin, X. and Zeevart, J. A. (1999) The 9-*cis*-epoxycarotenoid cleavage reaction is the key regulatory step of abscisic acid biosynthesis in water-stressed bean. *Proc. Natl. Acad. Sci. U.S.A.* **96**, 15354–15361
- 20 Thompson, A. J., Jackson, A. C., Symonds, R. C., Mulholland, B. J., Dadswell, A. R., Blake, P. S., Burbidge, A. and Taylor, I. B. (2000) Ectopic expression of a tomato 9-*cis*-epoxycarotenoid dioxygenase gene causes over-production of abscisic acid. *Plant J.* **23**, 363–374
- 21 Iuchi, S., Kobayashi, M., Tajiri, T., Naramoto, M., Seki, M., Kato, T., Tabata, S., Kakubari, Y., Yamaguchi-Shinozaki, K. and Shinozaki, K. (2001) Regulation of drought tolerance by gene manipulation of 9-*cis*-epoxycarotenoids dioxygenase, a key enzyme in abscisic acid biosynthesis in *Arabidopsis*. *Plant J.* **27**, 325–333
- 22 Qin, X. and Zeevart, J. A. (2002) Overexpression of a 9-*cis*-epoxycarotenoid dioxygenase gene in *Nicotiana plumbaginifolia* increases abscisic acid and phaseic acid levels and enhances drought tolerance. *Plant Physiol.* **128**, 544–551
- 23 Guiliano, G., Al-Babili, S. and von Lintig, J. (2003) Carotenoid oxygenases: cleave it or leave it. *Trends Plant Sci.* **8**, 145–149
- 24 Camara, B. and Bouvier, F. (2004) Oxidative remodeling of plastid carotenoids. *Arch. Biochem. Biophys.* **430**, 16–21
- 25 Tan, B. C., Joseph, L. M., Deng, W. T., Liu, L., Li, Q. B., Cline, K. and McCarty, D. R. (2003) Molecular characterization of the *Arabidopsis* 9-*cis* epoxycarotenoid dioxygenase gene family. *Plant J.* **35**, 44–56
- 26 Bouvier, F., Suire, C., Mutterer, J. and Camara, B. (2003) Oxidative remodeling of chromoplast carotenoids: identification of the carotenoid dioxygenase *CsCCD* and *CsZCD* genes involved in *Crocus* secondary metabolite biogenesis. *Plant Cell* **15**, 47–62
- 27 Bouvier, F., Dogbo, O. and Camara, B. (2003) Biosynthesis of the food and cosmetic plant pigment bixin (annatto). *Science* **300**, 2089–2091
- 28 Schwartz, S. H., Xiaoqiong, Q. and Zeevart, J. A. (2001) Characterization of a novel carotenoid cleavage dioxygenase from plants. *J. Biol. Chem.* **276**, 25208–25211
- 29 Simkin, A. J., Schwartz, S. H., Auldridge, M., Taylor, M. G. and Klee, H. J. (2004) The tomato carotenoid cleavage *dioxygenase 1* genes contribute to the formation of the flavor volatiles β -ionone, pseudoionone, and geranylacetone. *Plant J.* **40**, 882–892
- 30 Simkin, A. J., Underwood, B. A., Auldridge, M., Loucas, H. M., Shibuya, K., Schmelz, E., Clark, D. G. and Klee, H. J. (2004) Circadian regulation of the PhCCD1 carotenoid cleavage dioxygenase controls emission of β -ionone, a fragrance volatile of petunia flowers. *Plant Physiol.* **136**, 3504–3514
- 31 Ward, S. P. and Leyser, O. (2004) Shoot branching. *Curr. Opin. Plant Biol.* **7**, 73–78
- 32 Schwartz, S. H., Qin, X. and Loewen, M. C. (2004) The biochemical characterization of two carotenoid cleavage enzymes from *Arabidopsis* indicates that a carotenoid-derived compound inhibits lateral branching. *J. Biol. Chem.* **279**, 46940–46945
- 33 Booker, J., Auldridge, M., Wills, S., McCarty, D., Klee, H. and Leyser, O. (2004) MAX3/CCD7 is a carotenoid cleavage dioxygenase required for the synthesis of a novel plant signaling molecule. *Curr. Biol.* **14**, 1232–1238
- 34 Auldridge, M. E., Block, A., Vogel, J. T., Dabney-Smith, C., Mila, I., Bouzayen, M., Magallanes-Lundback, M., Dellapenna, D., McCarty, D. R. and Klee, H. J. (2006) Characterization of three members of the *Arabidopsis* carotenoid cleavage dioxygenase family demonstrates the divergent roles of this multifunctional enzyme family. *Plant J.* **45**, 982–993
- 35 Ruch, S., Beyer, P., Ernst, H. and Al-Babili, S. (2005) Retinal biosynthesis in eubacteria: *in vitro* characterization of a novel carotenoid oxygenase from *Synechocystis* sp. PCC6803. *Mol. Microbiol.* **55**, 1015–1024
- 36 Kloer, D. P., Ruch, S., Al-Babili, S., Beyer, P. and Schulz, G. E. (2005) The structure of a retinal-forming carotenoid oxygenase. *Science* **308**, 267–269
- 37 Rippka, R., Deruelles, J., Waterbury, J. B., Herdmann, M. and Stanier, R. T. (1979) Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *J. Gen. Microbiol.* **111**, 1–61
- 38 Barua, A. B. and Olson, J. A. (2000) β -Carotene is converted primarily to retinoids in rats *in vivo*. *J. Nutr.* **130**, 1996–2001
- 39 Schwede, T., Kopp, J., Guex, N. and Peitsch, M. C. (2003) SWISS-MODEL: an automated protein homology-modeling server. *Nucleic Acids Res.* **31**, 3381–3385
- 40 Krissinel, E. and Henrick, K. (2005) Detection of Protein Assemblies in Crystals. In *CompLife 2005* (Berthold, M. R., Glen, R., Diederichs, K., Kohlbacher, O. and Fischer, I., eds.), pp. 163–174, Springer-Verlag, Berlin and Heidelberg
- 41 Sanner, M. F., Olson, A. J. and Spehner, J. C. (1996) Reduced surface: an efficient way to compute molecular surfaces. *Biopolymers* **38**, 305–320
- 42 Ermakova, S. Y., Elanskaya, I. V., Kallies, K. U., Weihe, A., Börner, T. and Shestakov, S. V. (1993) Cloning and sequencing of mutant *psbB* genes of the cyanobacterium *Synechocystis* PCC6803. *Photosynth. Res.* **37**, 139–146
- 43 Robinson, K. R., Lorenzi, L., Ceccarelli, N. and Gaultieri, P. (1998) Retinal identification in *Pelvetia fastigiata*. *Biophys. Res. Commun.* **243**, 776–778
- 44 Chothia, C. and Lesk, A. M. (1986) The relation between the divergence of sequence and structure in proteins. *EMBO J.* **5**, 823–826
- 45 Moise, A. R., von Lintig, J. and Palczewski, K. (2005) Related enzymes solve evolutionarily recurrent problems in the metabolism of carotenoids. *Trends Plant Sci.* **10**, 178–186
- 46 Bouvier, F., Isner, J. C., Dogbo, O. and Camara, B. (2005) Oxidative tailoring of carotenoids: a prospect towards novel functions in plants. *Trends Plant Sci.* **10**, 187–194
- 47 Sabehi, G., Loy, A., Jung, K. H., Partha, R., Spudich, J. L., Isaacson, T., Hirschberg, J., Wagner, M. and Bjé, O. (2005) New insights into metabolic properties of marine bacteria encoding proteorhodopsins. *PLoS Biol.* **3**, e273
- 48 Peck, R. F., Echavarri-Erasun, C., Johnson, E. A., Ng, W. V., Kennedy, S. P., Hood, L., DasSarma, S. and Krebs, M. P. (2001) *brp* and *blh* are required for synthesis of the retinal cofactor of bacteriorhodopsin in *Halobacterium salinarum*. *J. Biol. Chem.* **276**, 5739–5744
- 49 Takaichi, S., Mochimaru, M., Maoka, T. and Katoh, H. (2005) Myxol and 4-ketomyxol 2'-fucosides, not rhamnosides, from *Anabaena* sp. PCC 7120 and *Nostoc punctiforme* PCC 73102, and proposal for the biosynthetic pathway of carotenoids. *Plant Cell Physiol.* **46**, 497–504
- 50 Zorn, H., Langhoff, S., Scheibner, M., Nimtz, M. and Berger, R. G. (2003) A peroxidase from *Lepista irina* cleaves β , β -carotene to flavor compounds. *Biol. Chem.* **384**, 1049–1056
- 51 Hihara, Y., Kamei, A., Kanehisa, M., Kaplan, A. and Ikeuchi, M. (2001) DNA microarray analysis of cyanobacterial gene expression during acclimation to high light. *Plant Cell* **13**, 793–806
- 52 Marin, K., Kanesaki, Y., Los, D. A., Murata, N., Suzuki, I. and Hagemann, M. (2004) Gene expression profiling reflects physiological processes in salt acclimation of *Synechocystis* sp. strain PCC 6803. *Plant Physiol.* **136**, 3290–3300
- 53 Al-Babili, S., von Lintig, J., Haubruck, H. and Beyer, P. (1996) A novel, soluble form of phytoene desaturase from *Narcissus pseudonarcissus* chromoplasts is Hsp70-complexed and competent for flavinylation, membrane association and enzymatic activation. *Plant J.* **9**, 601–612
- 54 Xue, L., Gollapalli, D. R., Maiti, P., Jahng, W. J. and Rando, R. R. (2004) A palmitoylation switch mechanism in the regulation of the visual cycle. *Cell* **117**, 761–771
- 55 Enzell, C. R. and Back, S. (1995) Mass spectrometry. In *Carotenoids* (Britton, G., Liaaen-Jensen, S. and Pfander, H., eds), vol. 1B, p. 264, Birkhäuser Verlag, Basel

Received 20 April 2006/1 June 2006; accepted 7 June 2006

Published as BJ Immediate Publication 7 June 2006, doi:10.1042/BJ20060592